*Proceedings and abstracts of the 6th International Symposium on Mineralogy
Cluj-Napoca – Romania, September 18-21, 2003*

Reviewers:

Univ. Prof. Dr. Volker Hoeck – „Paris Lodron” University of Salzburg (Austria)
Univ. Prof. Dr. Ioan Balintoni – „Babeş-Bolyai” University of Cluj-Napoca (Romania)
Univ. Prof. Dr. Lucreția Ghergari – „Babeş-Bolyai” University of Cluj-Napoca (Romania)
Univ. Prof. Dr. Corina Ionescu – „Babeş-Bolyai” University of Cluj-Napoca (Romania)
Univ. Prof. Dr. Bogdan Onac – „Babeş-Bolyai” University of Cluj-Napoca (Romania)
Univ. Prof. Dr. Șerban Vlad – „Babeş-Bolyai” University of Cluj-Napoca (Romania)
Dr. Alexandru Szakacs- Romanian Academy, Institute of Geodinamics, Bucharest (Romania)
Mrs. Tania Makarenko (for the English text) – „Babeş-Bolyai” University of Cluj-Napoca (Romania)

© 2003 Ionescu C. & Hoeck V. (Eds.). All rights reserved.

The authors are responsible for the content of their contributions.

**The editing and printing of this volume was supported by Grant 51/46/2002 (CNCSIS)
and the following sponsors:**

**Cluj-Napoca Municipal Council
S.C. Roșia Montană Gold Corporation S.A. Alba Iulia
S.C. Podvinalco S.A. Cluj-Napoca
S.C. Algo SRL Alba Iulia
S.C. Print Art S.R.L. Cluj-Napoca
S.C. Argisano S.R.L. Cluj-Napoca**

Computer editing: **Corina Ionescu & Alina Vesa**

**“Babeş-Bolyai” University of Cluj-Napoca
Studia Universitatis Babeş-Bolyai
24, Republicii Str.
Tel.: 0264-40.53.52; Fax: 0264-59.19.06
RO-3400 Cluj-Napoca, ROMANIA**

ISSN 1221-0803

*Mineralogical Society of
Romania*

*“Babeş-Bolyai” University of Cluj-Napoca
Faculty of Biology and Geology
Chair of Mineralogy
Romania*

Proceedings and abstracts

*of the **6th International Symposium on Mineralogy**
Cluj-Napoca — Romania, September 18-21, 2003
dedicated to Univ. Prof. Dr. Lucreția Ghergari on
the occasion of her 70th anniversary*

*Edited by:
Corina Ionescu and Volker Hoeck*

Acknowledgements

The Organizing Committee of the symposium take the opportunity of expressing their sincere gratitude to Mrs. Tania Makarenko, architect Romulus Zamfir, geol. engineer Horea Uioreanu, professor Ștefan Ionescu and engineer Mihai Păunescu from Cluj-Napoca, as well as to engineer Gary O'Connor and engineer Mircea Bocan from Alba Iulia, for their valuable help.

CONTENTS

Plenary session	
Ionescu, C., Onac, B.P.: Univ. Prof. Dr. Lucreția Ghergari: a life dedicated to science	
Balintoni, I.: Towards an improved model of the Laramian Transylvanides	
O'Connor, G., Szentesy, C., Breban, R.: Applied mineralogy and exploration techniques on the development of the Roșia Montană gold deposit (Roșia Montană, Romania)	
Proceedings and abstracts	
(in alphabetic order of the family name of the authors respectively of the first author indicated in the paper)	
Benea, M., Gorea, M.: The mineralogy and technological properties of some kaolins used in the ceramic industry	
Blănar, D.C., Domnișoru, D.: Mineral resources exploitation in the Brașov county (Romania): opportunities and threats to sustainable development	
Borovikova Yu.E.: Isomorphic substitutions in high and low vesuvianite structure	
Cauuet, B., Tămaș, C.G.: Dynamics of exploitation and types of mining workings in <i>Alburnus maior</i> ancient mining site (Roșia Montană, NW Romania)	
Damian, Gh., Damian, F.: Comparative study of the tetrahedrites from the Metaliferous Mts. and Baia Mare district (Romania) based on microprobe analyses	
Damian, Gh., Damian, F., Cook, J.N., Ciobanu, L.C.: Ag-sulphosalts in upper parts of the Baia Sprie deposit (Romania): microanalyses and implications for deposit zonality	
Gál, Á., Molnár, F., Ghergari, L., Szabó, Cs., Gatter, I.: Fluid inclusions and morphology of quartz from the Neogene epithermal deposits, Brad-Săcărâmb area (Transylvania, Romania)	
Gorea, M., Mariș, C.: The characteristics of quartz concentrate obtained by the flotation of the quartzo-feldspatic raw material from the Bedeciu deposit (Cluj district, Romania)	
Har, N.: Reaction coronas around quartz xenocrysts in the basaltic andesite from Detunata (Apusenii Mountains), Romania	
Har, N., Năstase, R.: Hercynite and magnetite in the hornfels xenoliths hosted by granodiorite from Valea Lungii (Cluj county, Romania)	
Hercot, O., Seghedi, I., Naud, J., Razvan Caracas, R.: Recent mineral deposition in the crater of the Ciomadul quaternary volcano - Harghita Mountains (Romania)	
Hîrtoanu, P., Udubaș, Gh.: Mineralogy of a "sacred monster": the Ditrău alkaline massif, East Carpathians (Romania)	
Iancu, G.O., Ionescu, C.: Types of chondrules from Romanian fallen ordinary chondrites	
Kouzmina, E.: The role of fluids during formation of the Svecofennian magmatic rocks (Russia)	
Laczkó, A.A.: The presence of tourmaline in the Harghita Mountains' volcanic structures (Romania)	
Marincea, Șt., Dumitraș, D.G., Diaconu, G.: Phosphates in the bat guano deposit from Grigore Decapolitul Cave (Căpățâni Mountains, South Carpathians), Romania	
Matovic, V., Vaskovic, N., Rosic, A.: Salt damages of some sandstone buildings in Belgrade (Serbia)	
Mârza, I., Pomârleanu, V.: Petrographic notes on the Praid salt breccia (Romania)	
Moazzen, M.: Chlorite-chloritoid-garnet equilibria and geothermometry in the Sanandaj-Sirjan metamorphic belt, Southern Iran	
Murariu, T., Rădășanu, S., Kasper, H.U., Schoenbeck, T.: Geochemical features of beryl from Voislova pegmatites (South Carpathians)	
Onac, B.P., Vereș, D.Ș., Kearns, J., Chirienco, M., Minuț, A., Breban, R.: Secondary minerals found in old mine galleries from Roșia Montană, Romania	
Onac B.P., White, W.B.: An unusual occurrence of berlinite (AlPO ₄) in the phosphate-bearing sediments from the Cioclovina Cave, Romania	

Onac, B.P., White, W.B., Hess, J.W.: The relationship between the mineralization of breccia pipes and mineral composition of speleothems: evidences from the Corkscrew Cave, Arizona (USA).....	
Pertlik, F.: Cubic CoTe ₂ : synthesis, structure determination and reflected light investigations	
Pintea, I.: Source regions of the melt and fluid phases in the porphyry Cu-Au-Mo deposits and others volcanic structures from Alpine Carpathian chain (Romania).....	
Pintea, I., Mosonyi, E., Bardocz, Z.: Textural features and melt inclusion types in andesites and basaltic andesites from the Eastern Carpathians (Călimani – Gurghiu – Harghita Mts.), Romania	
Pomârleanu, V., Mârza, I.: Fluid inclusions in salt (Transylvanian Basin, Romania)	
Pop, D., Bedeleian, I., Bedeleian, H.: XRD investigation of the clay fraction in some Romanian zeolitic volcanic tuffs and diatomites	
Pop, D., Stremțan, C.: The website of the Mineralogical Museum, Babeș-Bolyai University, Cluj-Napoca, Romania	
Radu, D.M.: Pyroxenite – marble interaction, a petrogenetic grid and a working example from the Inău crystalline island, Maramureș, Romania	
Rădășanu, S., Murariu, T., Kasper, H.U., Schoenbeck, T.: The geochemistry of apatite from Răzoare pegmatites (Romania).....	
Self, C.: The ontogeny of speleothems	
Simon, V., Ionescu, C., Dărăban, L.: Spectroscopic investigations of some obsidian archaeological artifacts	
Sreckovic-Batocanin, D., Nikolin, B., Vaskovic, N.: Pillow-lavas of the Povlen Mt. (Western Serbia)	
Stumbea, D.: Some geochemical remarks on basalts from Băița Bihor (Bihor Mts., Romania). Major elements	
Strutinski, C., Puște, A., Stan, R.: The metamorphic basement of the Romanian Carpathians: a discussion in the light of radiogenic K-Ar and ⁴⁰ Ar/ ³⁹ Ar dating	
Szakács, Al.: Mineral chemistry of the primary magmatic mineral assemblage of the „Dej tuff” (Romania).....	
Szakáll, S., Sajó, I.: Volborthite, a hydrous copper-vanadate from phosphate-bearing argillites in Dédestapolcsány, Uppony Mts. (N-Hungary)	
Tămaș, G.C., Costin, D.: Tetrahedrite – tennantite from the Roșia Montană gold-silver epithermal ore deposit, Apuseni Mountains (Romania).....	
Tămaș, G.C., Ghergari, L., Ionescu, C., Cauuet, B.: The relationships between breccia structures and ore veins. A case study from the Cetate Hill, Roșia Montană deposit (Romania).....	
Thanasuthipitak, P., Thanasuthipitak, T., Phetpu, K.: Mineral inclusions in ruby from Andilamena, Madagascar	
Vaskovic, N., Matovic, V., Sreckovic-Batocanin, D.: Petrology of garnet-amphibolite with white mica from Vranjska Banja Series (Serbian-Macedonian Massif, SE Serbia).....	
Viczián, I.: Mineralogy of Pliocene to Pleistocene pelitic sediments of the Great Hungarian Plain	
Viehmman, I., Ghergari, L.: The genesis of calcite rafts from caves – a point of view.....	
Vlad, Ș., Costin, D.: Mine-waste management	
Zugrăvescu, D., Polonic, G., Negoită V.: Low magnitudes of matrix stress related to aquathermal effect and montmorillonite dehydration in the Căldărușani-Tulnici geodynamic polygon, Romania	

Appendix: List of the papers published by Univ. Prof. Dr. Lucreția Ghergari.....

Index of authors

Advertisers

Plenary lectures

Towards an improved model of the Laramian Transylvanides

Ioan Balintoni¹

The Transylvanides which represent the uppermost group of Alpine tectonic units of the Apuseni Mountains originated from a Mesozoic rift located between the Preapulian and Getic cratons (Rădulescu & Săndulescu, 1973; Săndulescu, 1984; Balintoni, 1997). The term "Foreapulian Block" ("Preapulian bloc" in Romanian translation), was used by Săndulescu (1994), for the continental mass from which the Northern Apusenides or Inner Dacides (the Codru and Biharia Nappe Systems) have been sheared. The name "Getic Craton" was proposed by Balintoni (1994a) for the continental fragment located between the Transylvanian Rift and the External Carpathian Flysch Basin, from which proceeded the Getic crystalline. The Transylvanides were emplaced antithetically, during the Austrian and Laramian orogenic phases. During the compressional (Early Cretaceous) period, the rift basin evolved towards a foreland retroarc type basin, because it was installed on the upper plate sheared margin. If the Austrian Transylvanides (ATS) and the Mediterranean Apusenides are described as "in-sequence" tectonic units, the Laramian Transylvanides (LTS) are "out of sequence". In the Apuseni Mountains tectonic context, the Austrian orogenic phase is considered intra-Albian or around the Aptian-Albian boundary, the Mediterranean one as intra-Turonian (pre-Gosau) and the Laramian one as intra-Maastrichtian and close to the Maastrichtian end. This fact complicates the recognition of the Transylvanides, as well as their description and classification. Balintoni (1994, 1997) proposed a dual classification of the Transylvanides, with particular names for the Austrian and Laramian ones, because some parts of the ATS can be found again within several units of the LTS. According to latter classification of this author, the ATS include the Izvoarele, Valea Muntelui, Feneş, Colţul Trascăului, Bedeleu, Ardeu, Căbeşti, Căpânaş-Techereu and Bejan nappes, and the LTS comprise the Groşi, Criş-Bucium, Vulcan, Frasin, Metalliferous Mountains, Curechiu-Stăniş and Mureş nappes. Besides this, the Laramian Transylvanides transported also the post-Austrian sedimentary covers. Regarding the ATS and LTS many unsolved questions still persist, as it is for instance: the precise age for pre-Austrian and post-Austrian sedimentary formations; the correlation between the Austrian tectonic units enclosed by the Laramian nappes; the number of the LTS; the amplitude of the tectonic displacement; the relation between the Apuseni Mountains and the South Carpathians; the opening age of the Transylvanian Rift; the development of the magmatic component of the Transylvanides; the initial locale for the sedimentary and magmatic formations.

In the following we will analyse some actual issues of the relation between the ATS and the LTS and present an improved model.

I. The Bucium Unit: fact or myth?

The Bucium Unit was first described by Ianovici *et al.* (1976) as a part of the South Apuseni Mountains. These authors considered the South Apuseni Mountains as built up of some Early Cretaceous facial-structural units, arranged later by tectonic thrusting and folding, and they mentioned in the lowermost position, the Bucium Unit. According to them, the Early Cretaceous formations of the Bucium Unit are transgressively deposited upon the crystalline schists of the Highiş rise, which is formed by the Baia de Arieş and Muncel tectonic units. They are consisting of: micritic limestones, Tithonian-Neocomian in age; the Căbeşti Beds, Hauterivian-Aptian; the Valea Dosului Beds, Aptian; the Ponor Beds, Albian and the Pârâul Izvorului Beds, Late Albian-Cenomanian. In their upper part, the Pârâul Izvorului Beds grade into the Cenomanian Negrileasa conglomerates. The Pârâul Izvorului Beds unconformably overly the earlier formations, due to the Austrian orogenic movements.

Bleahu *et al.* (1981) confirm that the sedimentary deposits of the Bucium Unit constitute the cover of the Baia de Arieş and Muncel nappes, yet they partially modify the lithostratigraphy of these deposits.

Lupu (1983) considers the Bucium Unit as autochthonous, tectonically in a similar position as outlined by Bleahu *et al.* (1981).

Săndulescu (1984) regards the Bucium Unit as a sedimentary zone of the Baia de Arieş Nappe, and thus including it into the Internal Dacides or Northern Apusenides, together with the Vidolm Unit (see below).

Bordea (1992) conceives the „Bucium Nappe” as a Laramian tectonic unit overthrusting the Late Cretaceous cover of the crystalline towards the north. The trace of the overthrust is visible along the Hărăguş Hill-Medreşti Village. This author thinks that the relationship between the oldest sediments known in the Bucium Nappe and their basement (crystalline or ophiolites) could be tectonic because the intense deformation noticed within the black clays of the Ciuruleasa Formation, the lowermost lithostratigraphic unit of the Bucium Nappe. He is speaking about ophiolite basement for the reason that the Ciuruleasa and Valea Povernei formations contain ophiolite boulders, probably taken over from a such basement. The age of this tectonic contact is not discussed. Bordea ascribes the following formations to the Bucium Nappe: the Ciuruleasa Formation, probably Neocomian in age; the Valea Povernei Formation, Hauterivian-Early Aptian; a limy-schistous flysch, Aptian; the Gray Wildflysch Formation, Late Aptian-Early Albian; the Pârâul Izvorului Formation, Late Albian and the Negrileasa conglomerate, Cenomanian in age.

Balintoni (1997) follows the idea that the Bucium Nappe represents a Laramian tectonic unit, and underlines that its Early Cretaceous formations have to be described as part of the ATS, because they incorporate ophiolitic material, and this material could not be deposited directly on the continental crust of the Preapulian Craton.

Considering the above models and combining them with a careful study of the Blăjeni (Bordea, Constantinescu, 1975) and Abrud (Bordea *et al.*, 1979) sheets of the Geological Map of Romania, scale 1:50 000, the following comments can be made. The sequences older than the Pârâul Izvorului Formation of the Bucium Nappe sensu Bordea, probably belong to the Austrian Feneş Nappe, because the Early Cretaceous sedimentary suites of the two nappes can be paralleled. For example, Feneş type-beds crop out around Întregalde region, immediately westward of the boundary reverse fault of the Metalliferous Mountains Nappe. These beds are in a visible tectonic contact with the Baia de Arieş crystalline.

The tectonic line between the Medreşti Village and Hărăguş Hill, a Laramian overthrust according to Bordea (1992), can only be traced as an reverse fault on the Blăjeni and Abrud map sheets (scale 1:50 000). This is supported by the geometry of the tectonic line. The fact that in several places there is a continuation of the Maastrichtian sedimentation across the reverse fault line suggests an Early Maastrichtian or intra-Maastrichtian age for it (Săndulescu, 1984). The Late Cretaceous, pre-Maastrichtian sediments are missing on the southern compartment of the fault, indicating an erosion before the deposition of Maastrichtian strata. The exact age of the fault can be established only by new detailed paleontological investigations, because in some areas the Latest Maastrichtian sediments seem also to be cut off by some younger Laramian thrusts. In conclusion, the Bucium Nappe disappears because the formations older than the Pârâul Izvorului belong to the Austrian Feneş Nappe. The deposits younger than Pârâul Izvorului represent three post-tectonic cover units in a higher position: post-Austrian, Albian-Cenomanian in age; post-Mediterranean, Coniacian-Campanian in age (Gosau facies sediments); post-Early Laramian, Maastrichtian in age (probably Late Maastrichtian). The last two covers transgress also the Baia de Arieş crystalline. The Medreşti Village-Hărăguş Hill tectonic line constitutes a simple reverse fault, which grades in a thrust towards west, projecting farther into the Criş Nappe frontal line. The Metalliferous Mountains Nappe (Balintoni, 1997) is the first true Laramian tectonic unit in the southern part of the above discussed region (Fig. 2). Within this nappe, the Austrian Feneş Nappe is situated in the lowermost position.

II. Does the Vidolm Nappe exist?

The existence of a Laramian basement nappe in the northern Trascău Mountains was postulated for the first time by Russo-Săndulescu and Berza (1976). These authors delineated, at the Valea Muntelui headwaters, a volcano-sedimentary formation beneath a Baia de Arieş-type crystalline. The Baia de Arieş crystalline constituted the autochthon during the Austrian emplacement of the Transylvanides. During the Mediterranean orogenic phase, the Baia de Arieş crystalline formed the Baia de Arieş Nappe, the uppermost tectonic unit of the Biharia Nappe System (Northern Apusenides). In that time, the Baia de Arieş Nappe moved together with the Austrian Transylvanides, situated on its back. During the Laramian orogenic phase, the Baia de Arieş crystalline, as a component of the Baia

de Arieş Mediterranean Nappe, has been involved in the „out-of-sequence” tectonics, specific for this period in comparison with the previous orogenic phases. Thus, in this later deformation cycle, the Baia de Arieş crystalline thrust in places over the Transylvanides. The status of the crystalline outcrops known in the Transylvanides domain was confused before 1986, when Balintoni and Iancu have shown, that inside the Transylvanides, there is a tectonic unit built up of a crystalline sequence different of the Baia de Arieş crystalline. They called this nappe as „Colţul Trascăului”, and the constitutive crystalline as „Trascău Series”. This nappe is a Transylvanide, without connection with the Baia de Arieş Nappe. The volcano-sedimentary formation should be part of the banatitic volcanic suite, Late Cretaceous in age, which would indicate a Laramian age for the crystalline overthrusting. Russo-Săndulescu and Berza (1976) did not discuss the continuation of the basement overthrust, but they named the lower unit „Boierişte”, considering its outcropping as a tectonic window.

Fig. 1. Structural sketch of South Apuseni Mountains between Bucium and Blăjeni. 1- Baia de Arieş Nappe crystalline; 2 - Feneş Nappe; 3 – Late Aptian-Early Albian syn-tectonic formations; 4 - post-Austrian sedimentary cover (Late Albian-Cenomanian); 5 - post-Mediterranean sedimentary cover (Coniacian – Early Maastrichtian); 6 - post – Early Laramian sedimentary cover (Late Maastrichtian); 7 - Frasin Nappe; 8 - Vulcan Nappe; 9 - Tertiary magmatic and sedimentary rocks; 10 - banatites; 11 - intra-Maastrichtian overthrust; 12 - intra-Maastrichtian thrust; 13 - Latest Maastrichtian overthrust; 14 – The boundary between the Criş Nappe and the former Bucium Nappe; 15 – Unclear relationship. **B** – Feneş Nappe over the Baia de Arieş Crystalline, **C** – Feneş Nappe as part of the Criş Laramian Nappe, **MM** – Feneş Nappe as part of the Laramian Metaliferous Mountain Nappe.

Bleahu *et al.* (1981) assumed that the Boierişte tectonic window is part of the Feneş Nappe and assigned consequently the crystalline situated above the Boierişte, to the new Fundoaia Nappe, which can be traced close to the Mănăstirea Valley. In their hypothesis, the Fundoaia Nappe was formed by the Baia de Arieş-type crystalline. In that area, these authors distinguished, beneath the Fundoaia Nappe, the Hospea Nappe, also a new nappe, and they put together the Fundoaia and Hospea nappes within the Bedeleu Nappe System. This system is tectonically situated higher than the Feneş Nappe, and overthrust during the Laramian orogenic phase as a

single entity. Westward of the Bedeleu summit, Bleahu *et al.* (1981) have drawn two tectonic lines: a vertical fault limiting the extent of the Fundoaia Nappe, and a thrust, in the front of the Feneş Nappe. Lupu (1983) maintains Bleahu's *et al.* (1981) scheme, but he considers the frontal line of the Feneş Nappe as being an overthrust along its northern part.

Based on the structural sketch drawn by Russo-Săndulescu and Berza (1976), Săndulescu (1984) divided the Fundoaia Nappe *sensu* Bleahu in: a) the Fundoaia Nappe s.s., formed from ophiolites and crystalline limestones and b) the Vidolm Nappe, composed from the Baia de Arieş type crystalline overlying the volcano-sedimentary formation of the Boierişte tectonic window. This is a new concept, because within the Transylvanides domain two basement nappes (Fundoaia s.s. and Vidolm) appear. In this case, the Fundoaia Nappe s.s. is placed between the Vidolm Nappe and Bedeleu Nappe (or Trascău Nappe). According to Săndulescu (1984) the Feneş Nappe is in a higher position than the Vidolm Nappe, a correct vision, because he knew that the Baia de Arieş crystalline represented the autochthon for the Transylvanides, a fact neglected by Bleahu *et al.* (1981). The relation between the Fundoaia Nappe s.s. and the Feneş Nappe cannot be estimated because they do not come in contact one with each other. Towards the south, the Fundoaia Nappe s.s. ends before the onset of the Feneş Nappe. Westwards, the front of the Vidolm Nappe is traced somewhere beyond the Feneş Nappe boundary in the sense of Bleahu's *et al.* (1981). Additionally, Săndulescu suggests a possible subdivision of the Fundoaia Nappe s.s. into two units, because the metamorphosed limestones lie above unmetamorphosed ophiolites.

Balintoni & Iancu (1986) described the Colţu Trascăului Nappe (see above), formed by Middle Paleozoic metamorphics, emplaced above the Izvoarele and Valea Muntelui nappes. Mapping the Trascău Mountains region, the cited authors found that, beneath the Trascău Series crystalline and over the Baia de Arieş crystalline (as part of the Baia de Arieş Nappe), two types of formations were situated: Aptychus Beds and ophiolites. The very intricate relations between the two rock-suites have been solved tectonically, supposing that the ophiolites overthrust the Aptychus Beds. The Aptychus Beds have been included in the Izvoarele Nappe and the ophiolites in the Valea Muntelui Nappe. Ulterior researches (Balintoni, *unpublished data*) suggested that the Valea Muntelui and Izvoarele nappes can be paralleled with the Feneş Nappe. Balintoni and Iancu (1986) admitted that the Baia de Arieş Nappe has been intersected by a Laramian tectonic plane; its rear part, together with the Transylvanides on its back, overthrust the Boierişte volcano-sedimentary sequence located on the frontal part of the Baia de Arieş Nappe, together with the post-Mediterranean sediments. They named this Laramian tectonic unit the Lunca Arieşului Nappe. Its frontal line was traced approximately along the frontal course of the Feneş Nappe drawn by Bleahu *et al.* (1981) and Lupu (1983). The justification of this option was given by the supposition that alongside of this line, the bottom conglomerates of the Albian Ponor Formation were in an abnormal contact with Late Cretaceous deposits.

For tracing a Laramian tectonic plane, only two possibilities exist westward of the Bedeleu summit: immediately west of the summit, where the Baia de Arieş-type crystalline, carrying on its back the Albian conglomerates, thrusts in places the same Albian conglomerates; or along the foregoing reported contact, where the Albian conglomerates seemed to be situated over Late Cretaceous deposits. The thrust, which sometimes cuts off the conglomerates but in other places stays within the conglomerates, can be interpreted, at most, as an inverse fault, not as an overthrust. Consequently, only the second possibility remained for solving the issue of the Lunca Arieşului Nappe front position, because, apparently, along that line the older formations were situated above the younger ones.

With the occasion of the Poşaga sheet mapping, Lupu (in Balintoni *et al.*, 1987) showed that the relationship between the Albian and Late Cretaceous sediments is normal, transgressive and not tectonic.

The geological relations projected onto the Poşaga map sheet do not permit the tracing of a Laramian overthrust westward of the Bedeleu summit (Trascău). As a consequence of this situation, there is no way of advocating further on the existence of the Vidolm or Lunca Arieşului Nappe.

With respect to the conclusion that the Vidolm Nappe (or Lunca Arieşului) does not exist, the geological *enigma* of the volcano-sedimentary formation from the Boierişte area, remains to be understood. Thus, three explanations can be suggested, as follows:

1. The volcanic formation consists of the remnants of an explosive activity, similar with those known from the Baia de Arieş or Roşia Montană Neogene areas, containing polymictic breccias.

2. It represents some Quaternary deposits generated at a dolina-bottom, by drainage. The discharge probably took place along the contact between the Colţu Trascăului limestones and the Baia de Arieş Nappe crystalline.
3. A combination of these both hypotheses.

III. The significance of the Bozeş Nappe

There are two important issues to be clarified about the Bozeş Nappe, as it was originally defined by Bleahu *et al.* (1981):

1. Does it really represent a tectonic unit?
2. To which mountains does it belong: the Apuseni Mountains or the South Carpathians?

Ianovici *et al.* (1976) described the unit which was believed later to represent the sedimentary component of the Bozeş Nappe (Bleahu *et al.*, 1981), as „the Turonian? – Senonian from the south-eastern part of the Metalliferous Mountains”. Consequently, this succession should represent the post-tectonic cover of some Austrian tectonic units from the South Apuseni (Căpâlnaş-Techereu and Feneş units). On the other hand, these sediments transgress on the Rapolt crystalline and on the Sebeş Mountains crystalline, both belonging to the South Carpathians. Ianovici *et al.* (1976) included in the „Turonian?-Senonian from the south-eastern part of the Metalliferous Mountains” the Bobâlna, Geoagiu and Bozeş beds, the last one being considered as a 3000 m thick typical flysch. Bordea *et al.* (1978) called the three lithostratigraphic entities a „Cretaceous post tectonic cover”. Josefina Bordea (in Borcos *et al.*, 1981) assumes an overthrust in front of the Bozeş Beds.

Bleahu *et al.* (1981) denominated the Bobâlna, Geoagiu and Bozeş beds as „formations” and included them, together with the Rapolt crystalline, in the Laramian Bozeş Nappe. A questionable tectonic line was drawn westward of the Rapolt crystalline. In the general sketch of the tectonic units from the South Apuseni Mountains, the Bozeş Nappe was placed between the „Bedeleu Nappe System” and the Valea Mică-Galda Nappe. In the Bulbuci zone, in a tectonic window, outcrops of the Fenes and Valea Mica-Galda nappes occur beneath the Bozeş Nappe.

Lupu (1983) arranges the Bozeş Nappe between the „Bedeleu Nappe System” and the „Criş Nappe System”.

The conclusion which can be drawn from the last two models, is that the Bozeş Nappe is a tectonic element of the South Apuseni Mountains although the Rapolt crystalline evidently belongs to the South Carpathians.

Săndulescu (1984) considers the Bozeş Nappe as a cover unit, consisting of Late Cretaceous deposits. Towards the south, it is bordered by the South Transylvanian Fault, which separates it from the South Carpathians. In this way he tries to solve the contradiction existant in the previous models. On the tectonic sketch of the Meridional Apusenides by Săndulescu (1984), the South Transylvanian Fault passes towards southern part of the Transylvanian Depression, northward of the Rapolt crystalline. The mapping data does not support this hypothesis, because there are no any important E-W striking faults within the Bozes Beds (Bordea *et al.*, 1978).

Aware of this problem, Balintoni (1997) refers to the Mureş Laramian Nappe as a major tectonic unit, formed of the Bejan, Căbeşti and Căpâlnaş-Techereu Austrian subsidiary nappes, and includes in it also the Rapolt crystalline and Bozeş Beds. In this model, the frontal overthrust of the Bozeş Beds is prolonged westward into the Căpâlnaş-Techereu Laramian overthrust. Balintoni (1997) assumes that the pre-Laramian South Transylvanian Fault passes between the Bejan and the Căbeşti units, but does not cut off the Bozeş Beds, continuing eastwards beneath them. This model can be improved considering the following facts:

1. The Bozeş Beds transgress the Rapolt and the Sebeş Mountains crystalline. They are also very weakly deformed, similar to the Deva Beds, transgressing over the contact between the Poiana-Ruscă crystalline and Bejan Unit.
2. In contrast to the Deva and the Bozeş beds, the Late Cretaceous formations of the South Apuseni Mountains are strongly folded and faulted.
3. The amplitude of the South Apuseni Laramian thrusts and overthrusts diminishes towards the north, becoming vertical faults; at the northern end of the Trascău Mountains, eastern vergencies are observable too.

4. The age of the tectonic contact between the Căbeşti and the Bejan units is Laramian, as it cuts off the Turonian-Latest Cretaceous Deva Beds.

5. The vergences of the Laramian thrusts within the Transylvanian Depression are eastward oriented (Ciulavu, 1998).

Based on these observations, we may conclude:

1. The Bozeş Beds, together with the Rapolt crystalline, the Poiana Ruscă crystalline and the Bejan Unit form a single Laramian tectonic unit, being part of the South Carpathians. This unit can be called the Bozeş Nappe.

2. From Groşi till the north of Deva, the boundary of the Bozeş Nappe towards the Căbeşti Unit is almost vertical, representing a dextral strike-slip fault. At its western tip, between Burjuc and Groşi, Dinică *et al.* (1994) described strong mylonites developed in the sedimentary rocks of the Căbeşti Unit. These mylonites were misinterpreted as Paleozoic low grade metamorphics by Lupu *et al.* (1991).

3. The frontal boundary of the Bozeş Nappe does not represent the South Transylvanian Fault. The north-eastern prolongation of the Bozeş Nappe frontal line remains to be searched along the suture between the Preapulian and the Getic cratons. Its amplitude decreases towards north. Between Groşi and Deva this boundary can be defined as the South Apuseni Fault. Only between Deva and Alba Iulia it shows the features of an overthrust, because along this segment its direction changes. This changing in direction can be considered as a restraining bend along a strike-slip fault, the necessary condition for the appearance of oblique overthrusts.

4. The character of the South Apuseni Laramian deformations can be better explained if we accept a counterclockwise rotation of the Preapulian Craton around a pole situated next to the Preluca Massif. The paleomagnetic evidence for the supposed rotation is missing, but otherwise it is very difficult to constrain the gradual increasing of the Laramian thrust amplitude towards south-west, and also the west-east straight trace of the South Apuseni Fault.

5. Due to the separation of the Bozeş Nappe, previously a component of the Laramian Mureş Nappe (Balintoni, 1997), only the Căbeşti, Căpâlnaş-Techereu and Ardeu Austrian tectonic units remained to compose the Mureş Nappe.

6. The initial thrusting of the South Carpathian crystalline over the South Apuseni took place before the Latest Cretaceous times. During the Laramian orogenic phase, the South Carpathians thrust again over the South Apuseni Mountains, together with the Bozeş Beds.

IV. The Laramian Transylvanides

As we already showed, Ianovici *et al.* (1976), Bleahu *et al.* (1981), Lupu (1983), Săndulescu (1984) and Balintoni (1994, 1997) presented some general structural schemes of the Southern Apuseni Mountains, yet Balintoni (1994) made the first attempt to group the Transylvanides in an Austrian set and a Laramian one. In the latter he put together the Bucium, Frasin, Vulcan, Groşi, Criş, Feneş, Curechiu-Stănişia and Căpâlnaş-Techereu tectonic units.

In 1997, he modified the former scheme and noticed the following Laramian Transylvanides: Groşi, Criş-Bucium, Vulcan, Frasin, Metalliferous Mountains one, Curechiu-Stănişia and Mureş.

The data presented here, combined with the analysis of the Zlatna (Borcoş *et al.*, 1981) and Brad (Bordea & Borcoş, 1972) sheets of the Geological Map of Romania, scale 1:50 000, improved this scheme. In a new tectonic map (Fig. 2), the Laramian Transylvanides consist of the following nappes: Groşi Nappe; Criş Nappe; Vulcan Nappe; Frasin Nappe; Metalliferous Mountains Nappe; Mureş Nappe. In fact, the Vulcan and Frasin nappes exemplify no more than some tectonic „klippen”. The Metalliferous Mountains Nappe contains as Austrian Transylvanides, the Feneş, Colţul Trascăului and the Bedeleu nappes. The Mureş Nappe is formed from the Curechiu-Stănişia, Căpâlnaş-Techereu, Ardeu and Căbeşti Austrian Transylvanides. The relation between the Căpâlnaş-Techereu and Căbeşti units is still not completely understood. Parts of the Feneş Nappe preserved also under the Latest Cretaceous cover of the Baia de Arieş Nappe and within Criş Nappe body.

Fig. 2. The Laramian tectonic units of the South Apuseni Mountains: 1) Biharia Nappe System and its cover; 2) Groşi Nappe; 3) Criş Nappe; 4) Metalliferous Mountains Nappe; 5) Mureş Nappe; 6) Frasin Nappe; 7) Vulcan Nappe; 8) Bozeş Nappe (South Carpathians); 9) Tertiary Formations; 10) strike-slip movement.

In a palinspastic reconstruction of the Late Jurassic-Early Cretaceous sedimentation of the Transylvanian Rift, two big facial zones can be pointed out. One of them, represented by reef limestones installed on arc-type and MORB-type volcanics, is situated in a south-eastern position. The other one, constituted typically of Aptychus Beds, is located in a north-western position. The basement of the Aptychus Beds is not well-known, being expressed in places by volcanic agglomerates, lavas and radiolarites. They deposited off the arc, in deeper waters. The reef limestones crop out in the Bedeleu, Ardeu and Căpâlnaş-Techereu ATS. The Aptychus Beds are localized in the Feneş Nappe and they are also included as olistoliths in the Albian wildflysch sequence of the Groşi LTS. In the Criş LTS and the Curechiu-Stănişia ATS, Late Jurassic-Early Cretaceous micritic limestones, facially similar to the Aptychus beds, develop.

Considering the above information, we can suppose that, initially two big ATS were generated, an external one formed mainly of Aptychus Beds and micritic limestones, and an internal one built up of volcanics and reef limestones. First of them has been represented by the Feneş Nappe, emplaced on the Baia de Arieş crystalline. The other one overthrust the Feneş Nappe. During the Laramian orogenic phase, the Feneş Nappe was divided in several fragments. The frontal fragment remained on the Baia de Arieş crystalline, covered by the post-Mediterranean sediments. Another fragment is known at the base of the Metalliferous Mountains LTS; and another one is part of the Criş LTS.

It is not clear if the Curechiu-Stănişia ATS was a constituent of the Feneş Nappe, or formed a separate nappe or was a marginal piece of the upper big ATS.

The southern boundary of the Laramian Transylvanides runs along the northern front of the Bozeş Nappe.

Acknowledgements

Prof. V. Höck is acknowledged for his suggestions and observations which significantly improved the quality of the manuscript. My younger colleagues C. Balica and M. Cliveţi for solving the editorial questions. The drawer S. Moraru prepared the two sketches.

References

- Balintoni, I. (1994) Structure of the Apuseni Mountains. *Rom. Journ. Tect. Reg. Geol.*, **75**, Suppl. 2, 51-58.
- Balintoni, I. (1994a) The spatial and temporal classification of the Alpine geotectonic units of Romania in the light of plate tectonics. *Studia Univ. Babeş-Bolyai, Ser. Geol.*, **XXXIX**, 1-2, 7-20, Cluj-Napoca.
- Balintoni, I. (1997) The geotectonics of metamorphic terrains in Romania (in Romanian). *Edit. Carpatica*, 176 p., Cluj-Napoca.
- Balintoni, I., Iancu, V. (1986) Lithostratigraphic and tectonic units in the Trascău Mountains, North of Mănăstirea Valley, *D.S. Inst. Geol. Geofiz.* **70-71/5** (1983; 1984), 45-56, Bucureşti.
- Balintoni, I., Lupu, M., Iancu, V., Lazăr, C. (1986). Geological Map of Romania, scale 1:50 000, Poşaga sheet, *Inst. Geol. Geofiz.*, Bucureşti.
- Bleahu, M., Lupu, M., Patruşius, D., Bordea, S., Ştefan, A., Panin, S. (1981) The structure of the Apuseni Mountains. *Carp.-Balk. Geol. Assoc., XII Congr., Guide to excursion B3*, 107 p., *Inst. Geol. Geophys.* Bucureşti.
- Borcoş, M., Berbeleac, I., Bordea, S., Bordea, J., Mantea, G., Boştinescu, S. (1981) Geological map of Romania. Scale 1:50 000, Zlatna sheet. *Inst. Geol. Geofiz.* Bucureşti.
- Bordea, J., Berbeleac, I., Borcoş, M., Mantea, G., Stancu, J., Rogge-Ţăranu, E. (1978) Geological map of Romania, scale 1:50 000, Geoagiu Sheet. *Inst. Geol. Geofiz.*, Bucureşti.
- Bordea, S. (1992) The stratigraphy of the NeoJurassic and Cretaceous deposits from the western part of the Metaliferous Mts. *Ph.D. Thesis abstr.* 27 p., Univ. Al. I. Cuza, Iaşi.
- Bordea, S., Borcoş, M. (1972) Geological map of Romania, scale 1:50 000, Brad Sheet, *Inst. Geol.* Bucureşti
- Bordea, S., Constantinescu, R. (1975) Geological map of Romania, scale 1:50 000, Blăjeni Sheet. *Ins. Geol. Geofz.* Bucureşti.
- Bordea, S., Ştefan, A., Borcoş, M. (1979) Geological map of Romania, scale 1:50 000, Abrud Sheet. *Inst. Geol. Geofz.* Bucureşti.
- Ciulavu, D., (1998). Tertiary tectonics of the Transylvanian Basin. *Ph.D. Thesis*, 152 p., NSG 981105, Amsterdam
- Dinică, I., Pană, D., Conovici, M., Roşu, E., (1994). Dynamic metamorphism from the Mureş zone. *Anal. Univ. Bucureşti, Geol.*, **XLIII**, Abstr. vol.
- Ianovici, V., Borcoş, M., Bleahu, M., Patruşius, D., Lupu, M., Dimitrescu, R., Savu, H. (1976) The geology of the Apuseni Mts. (in Romanian). *Edit. Acad.*, 631 p., Bucureşti
- Lupu, M. (1983) The Mesozoic history of the South Apuseni Mountains. *An. Inst. Geol. Geofiz.*, **LX**, 115-124, Bucureşti
- Lupu, M., Marinescu, Fl., Rom, E., Nicolae, I., Mureşan, M. (1991) Geological map of Romania, scale 1:50 000, Lăpuş-Coştei Sheet. *Inst. Geol. Geofz.* Bucureşti.
- Rădulescu, D., Săndulescu, M. (1973) The plate tectonics and the geological structure of the Carpathians. *Tectonophysics*, **16**, 155-161, Amsterdam.
- Russo-Săndulescu, D., Berza, T. (1976) The Boierişte window from Valea Muntelui-Colţeşti (Trascău Mts.) (in Romanian) *D.S. Inst. Geol. Geofiz.*, **LXII**, 5 (1974-1975), 141-148, Bucureşti.
- Săndulescu, M., (1984). Geotectonics of Romania (in Romanian). *Edit. Tehn.*, 336 p., Bucureşti.

APPLIED MINERALOGY AND EXPLORATION TECHNIQUES ON THE DEVELOPMENT OF THE ROŞIA MONTANĂ GOLD DEPOSIT (ROŞIA MONTANĂ, ROMANIA)

Gary V. O'Connor², Cecilia Szentesy¹, Radu Breban¹

A number of applied mineralogy and exploration techniques have been used in the exploration and development of the Roşia Montană Gold deposit, Roşia Montană, Romania. The deposit is situated within the Apuseni Mountains region, approximately 65 kilometres west of the regional centre of Alba-Iulia and has been the focus of ongoing gold exploitation for over 2000 years (Fig. 1).

The exploration and development of the deposit has included the collection of 92,359 metres of drill samples from 695 underground and surface drill holes, including 28,439 metres of diamond drilling and 63,920 metres of reverse circulation drilling, in addition to 59,584 metres of underground channel sampling from the more than 140 kilometres of known underground development at Roşia Montană. These samples and underground access formed the source for much of the applied mineralogy and exploration techniques conducted.

The exploration program has included mineralogical studies on thin and polished sections. The use of fluid inclusion studies, XRD and microprobe analysis has also been applied to assist in the program. The geology and alteration, as well as structural setting have been modelled utilising Vulcan three-dimensional software. The same software has been used for the development of the resource estimation model and subsequent reserve model. The resource model has been calculated using ordinary kriging, multiple indicator kriging, inverse distance squared, and nearest neighbour techniques using 10 x 10m, 15 x 15 m and 20 x 20 x 10 m block sizes. The model and linked databases, using AcQuire and Micromine databases, also include all geological characteristics, geotechnical properties, metallurgical properties including recoveries, geophysical properties including hardness and abrasiveness, and geochemical properties including sulphur, sulphate and analysis of up to 47 elements, as well as the ABA and ARD properties of all rock types linked to the known mineralogy and geology.

The databases and models enable Roşia Montană Gold Corporation to know and understand the geology, mineralogy as well as the geochemical and geophysical characteristics of every block and thus rock type in the deposit.

Geology

The Roşia Montană deposit is hosted within rhyo-dacitic extrusive dome complexes and associated phreato-magmatic breccia complexes of Mid-Miocene age (13.5 Ma). The eruption of later andesitic flows and associated pyroclastics occurred in the area at 10.9 to 9.3 Ma. The magmatic rocks are hosted within Cretaceous age sediments, comprised dominantly of shales with subordinate sandstones, conglomerates and limestone units. Late basaltic andesite events have erupted at 7.4 Ma approximately 7 kilometres south east of Roşia Montană (Roşu *et al.*, 1997; Pécskay *et al.*, 1995).

Mineralogy

Gold mineralisation is associated with sulphides and sulpho-salts, including pyrite, sphalerite, galena, chalcopyrite, tetrahedrite (Ag-rich), proustite, argentite as well as tellurides (intermediate sulphidation). There is evidence for an early pyrite-arsenopyrite stage (low sulphidation). The Ag:Au ratio is highly variable, from low to high (associated with sulphides).

² S.C. Roşia Montană Gold Corporation S.A., 20, I.C. Brătianu, Str., RO-2500 Alba-Iulia, Romania. E-mail: <gary.oconnor@rmgc.ro>.

Fig. 1. Geology and location of the Roşia Montană deposit.

Gangue minerals are dominated by quartz, rhodochrosite, calcite, adularia and barite. Carbonate (rhodochrosite) plus base-metal sulphides decrease upwards and chinga veins (sulphidic siliceous rock flour) decrease with depth (Hedenquist, 2003).

Variable degrees of silicification, illite haloes, argillic zones and broad propylitic alteration are present also.

Applied Mineralogy

Based on fluid inclusion studies it has been modelled that the paleo-surface occurred at an elevation of approximately 1300 m asl. The current maximum elevation is 1046 m asl at Cârnic (Tămaş, 2002). Fragments of Cretaceous sediments as well as underlying Palaeozoic metamorphic rocks are present in the diatreme and hydrothermal breccias, in addition to dacitic fragments, indicating extensive vertical transport. Additionally wood fragments have been seen at depths of approximately 150 m below the current surface within hydrothermal breccias as well as at the current surface at Cârnic indicating large vertical convective movement within the breccias.

Fluid inclusion studies have indicated temperatures of deposition of 212-258^oC (average 230^oC) for quartz and 217-253^oC (average 238^oC) for carbonates, from very dilute (0.35-3.23 wt% NaCl) hydrothermal fluids (Leach & Hawke, 1997).

Exploration Techniques

The exploration of the Roşia Montană deposit has utilised the applied mineralogical studies to direct the exploration program in identifying gold mineralisation. The acquisition of data on the diamond drilling, on reverse circulation drilling and channel sampling program was conducted using pocket computers utilising "Field Marshall" software to digitally collect all geological and geotechnical data. This was down-loaded on a daily basis into "Micromine" and stored and validated in "AcQuire" data-bases. One metre samples were collected from all sampling media, with one in every 20 samples field duplicated. Samples are prepared to 98% -150 micron in LM-5 pulverisers and assayed for Au by a 50 g charge fire assay charge technique and Ag, Cu, Pb and Zn by AAS at the SGS managed laboratory in Gura Roşiei. Upon submission one "blind" internationally accredited standard is submitted for every 20 samples, plus on a monthly basis approximately 3% of all samples are sent for independent check analysis at laboratories in Australia and Canada. Total S, sulphide and sulphate analysis has been determined by Leco furnace and multi-element analysis by ICP-MS or ICP-AES and, where appropriate, by cold vapour AAS.

Over the course of the program to date, 5053 bulk density measurements have been made on selected drill core. Additionally all drill holes are surveyed both at surface and down hole, as have the underground drives. Aerial photography was used to produce 2 m contour maps and Landsat-TM, Aster and Spot satellite imagery has assisted in remote sensing analysis of the area. Geophysical techniques utilized have included aerial helicopter borne magnetics, the use of field magnetic susceptibility meters and ground induced polarisation and resistivity electrical methods as well as AMT surveys.

Three dimensional geology, alteration, mineralogical and resource models have utilised Vulcan software, which has also been used for generating digital terrain models. These models contain all available geological, geotechnical, mineralogical, metallurgical, geochemical, geophysical and physical characteristics of the deposit, which are used for modelling all aspects of the development of the deposit (Gossage & Barnes, 2002).

References

- Gossage, B., Barnes, J. (2002) Roşia Montană resource estimation-RSG Global. Independent report for RMGC, *unpubl.* Archives RMGC, 65 p + appendices.
- Hedenquist, J. (2003) Observations on the Roşia Montană gold deposit. Report for RMGC. 18 pages.
- Leach, T., Hawke, M. (1997) Petrologic study of Roşia Montană, Romania. Independent report for RMGC, *unpubl.* Archives RMGC, 15 p.
- Pécskay, Z., Edelstein, O., Seghedi, I., Szakács, A., Kovacs, M., Crihan, M., Bernad, A. (1995) K-Ar Datings of Neogene-Quaternary calc-alkaline volcanic rocks in Romania. *Acta Vulcanologica*, **7**, 2, 53-61.
- Roşu, E., Pécskay, Z., Ştefan, A., Popescu, Gh., Panaiotu, C., Panaiotu, C.E. (1997) The evolution of the Neogene volcanism in the Apuseni Mountains, Romania: Constraints from New K – Ar Data. *Geologica Carpathica*, **48**, 6, 353-359.
- Tămaş, C. (2002) "Breccia Pipe" structures associated with hydrothermal deposits from Romania (in Romanian). *Ph.D. Thesis*. "Babeş-Bolyai" University of Cluj-Napoca.

Proceedings and abstracts

THE MINERALOGY AND TECHNOLOGICAL PROPERTIES OF SOME KAOLINS USED IN THE CERAMIC INDUSTRY

Marcel Benea³, Maria Gorea¹

Three different kaolin types used in the ceramic industry (KO1 and MK2 kaoline from Ucraina and respectively BOJ kaoline from Bulgaria) were analysed by different methods in order to obtain a complete mineralogical and technological characterisation. Studies were carried out by using a combination of analyses of both, the bulk sample and the fine fraction. Following a well-established pre-treatment methodology (use of chemical reactivities, ultrasonic treatment, dispersion procedures), clay mineral concentration by centrifugation and sedimentation, oriented and random powder preparation, cation saturation, expansion/dehydration methods, a number of 12 X-ray diffraction (XRD) patterns⁴ were obtained for each sample. The interpretation of XRD patterns of the bulk samples, and of the fine clay fraction (< 2 µm) led to the identification of the following mineral phases: kaolinite, illite, quartz, and in a single one sample, also vermiculite. The semiquantitative mineralogical composition was determined based on peak intensities (mostly of lower orders) and peak areas corrected by various factors. The Hinckley crystallinity index was obtained measuring the heights of the (110) and (111) peaks. Scanning electron microscopy⁵ was also used in order to illustrate the identified mineral phases. For a better characterisation, the cation exchange capacity (CEC) and surface area (SA) were also measured by absorption of methylene blue. The particle size distribution was obtained using a laser granulometer⁶. The results of the above mentioned investigations are given in Table 1.

Table 1.

**Semiquantitative mineralogical composition (wt.%)
and physical features of kaolin raw materials**

Mineral/ Parameter	Kaolin type/sample number		
	KO1 / 4810	MK2 / 4811	BOJ / 4812
Kaolinite	89.00	69.85	82.95
Illite	6.00	19.50	10.40
Quartz	5.00	7.65	6.65
Vermiculite	-	3.00	-
CEC [mequ/100 g]	36.10	31.30	36.70
SA [m ² /g]	310	269	316
Hinckley index	0.44	0.87	1.35
Granulometry [%]:			
< 1 µm	15.36	2.64	11.15
< 20 µm	92.57	59.12	98.40
< 60 µm	100.00	98.39	100.00
< 100 µm	100.00	99.93	100.00

³ "Babeş-Bolyai" University of Cluj-Napoca, 1, Kogălniceanu Str., Dept. of Mineralogy, RO-3400 Cluj-Napoca, Romania. E-mail: <mbenea@bioge.ubbcluj.ro>, <mgorea@bioge.ubbcluj.ro>.

⁴ XRD: Philips PW 1710 diffractometer equipped with Bragg-Brentano geometry, copper anticathode (45kV, 40 mA), and graphite monochromator (University of Vienna, Austria).

⁵ SEM: Philips XL30 ESEM (University of Vienna, Austria).

⁶ Fritsch-Analysette 22 type Laser Granulometer (S.C. Faimar Baia Mare, Romania).

The main technological characteristics (dry and firing shrinkage, Pfefferkorn plasticity index, rheology, resistance, color after firing) of the ceramic masses prepared by using the analyzed kaolin types were also established.

There is a direct relationship between the mineralogical composition and particle size distribution of different kaolin types vs. the technological properties of the raw and fired ceramic products.

MINERAL RESOURCES EXPLOITATION IN THE BRAŞOV COUNTY (ROMANIA): OPPORTUNITIES AND THREATS TO SUSTAINABLE DEVELOPMENT

Doina Catrinel Blănaru⁷, Dan Domnişoru¹

The paper presents an alternative approach to the relation: “*mineral resources – environment protection*” vs “*economy – sustainable development*”. The regional strategies and policies, turned towards sustainable development by valorifying natural resources and especially regenerable mineral resources, may provide considerable income to local communities, without having a negative impact on the environment. Thus, the reevaluation of the mineral water resources having therapeutic effect from Braşov county (Romania), the measures taken for their protection as well as the studies undertaken for their rehabilitation and introduction in the touristic circuit are elements of the county strategies for a sustainable development. The threats of excessive and uncontrolled exploitation of the mineral resources both on the environment and on local communities, are presented also in the paper.

⁷Environment Protection Office of the Braşov County Council, Romania.

ISOMORPHIC SUBSTITUTIONS IN HIGH AND LOW VESUVIANITE STRUCTURE

Elena Yu. Borovikova⁸

Vesuvianite is an ortho-disilicate with complicated chemistry. Its general formula can be written as $X_{19}Y_{13}Z_{18}O_{68}T_{0-5}W_{10}$, where X is Ca and other large cations occupying 8-coordinate sites, Y are cations occupying 6 and 5-coordinate sites, as Al, Mg, Fe, Ti etc., Z are Si occupying 4-coordinate sites, T is B in 3 and 4-coordinate sites, and W are monovalent and divalent anions: OH⁻, F or O. Depending on atoms ordering in the structure, vesuvianite crystallizes in *P4/nnc* (high) or *P4/n* (low) space group. Low vesuvianite occurs in metamorphic rocks formed at low temperatures [$< 300^{\circ}\text{C}$, compared to those containing high vesuvianite ($400 - 800^{\circ}\text{C}$)].

A collection of more than 50 vesuvianite samples was investigated by a number of analytical methods such as electron microprobe analysis, IR-spectroscopy, Mössbauer spectroscopy and others. Vesuvianite samples were separated accordingly to their space groups, identified by IR-spectra.

On the basis of Mössbauer spectroscopy data, the schemes of high vesuvianite isomorphism are specified. It is shown that the

and

schemes are realized simultaneously and supplement each other. The scheme (1) plays the basic role, while the scheme (2) has a subordinated role.

Based on low vesuvianites IR-spectroscopy data, we suppose the ordering of different cations in the new Y(3a) and Y(3b) positions. The ordering of Y cations in two nonequivalent Y (3) sites may result in two ways: 1) Mg, Ti and Al occupy one of these sites, and Al, Fe³⁺ - the other one; 2) Mg and Al occupy one of Y(3) sites, and residual Al and high-valency cations occur in the other one. Thus, for charge-balance reasons, the OH-groups have to be substituted partly by O(11).

B can replace H in O(10) position, which is accompanied by the occurrence of new oxygen positions O(12) with triangular coordination around boron atoms. B can also occupy [O(11)₂ O(7)₂] tetrahedra. Thus, the positions of two atoms of hydrogen connected with O(11) should be vacant due to very short distance H - B. The IR-spectra of vesuvianites at $1300 - 1000 \text{ cm}^{-1}$ have shown that in the samples from rodingites the small quantity of B is triangular coordinated, replacing H in the O(10) position (1270 cm^{-1} vibration bond). In vesuvianites from skarns, at decreasing Al content, B starts to fill the tetrahedra (1110 cm^{-1} vibration band) besides triangles. The isomorphic scheme $\text{B} + (\text{Mg}, \text{Fe})^{2+} = 2\text{H} + \text{Al}$ is realized. B occurs in the tetrahedra of high Al and Mg, in practically Fe-free samples from the Talnakh skarns, generated by a rodingitization process. There the realized scheme of isomorphism is $\text{B} + \text{Mg} = 2\text{H} + \text{Fe}^{3+}$.

⁸ Lomonosov Moscow State University, 119992, GSP-2, MSU, A – 403, Russia. E-mail: <alena@geol.msu.ru>.

DYNAMICS OF EXPLOITATION AND TYPES OF MINING WORKINGS IN *ALBURNUS MAIOR* ANCIENT MINING SITE (ROŞIA MONTANĂ, NW ROMANIA)

Béatrice Cauuet⁹, Călin Tămaş¹⁰

At the beginning of the 2nd century A.D. a new economic space opened in the northeastern extremity of the Roman Empire. Rome acquired Dacia after the Trajanus's victorious campaign in 106 A.D. against Dacian king Decebal. The assets of this new territory were based on the presence of important metalliferous resources among which, gold and silver were the most important ones.

Located within the Apuseni Mountains, the Roşia Montană gold-silver ore district is one of the richest in Europe. The mining perimeter has about 6 km² and still remains extremely rich. In the ancient site of *Alburnus Maior*, located along the Roşia Valley, the mining activity developed mainly after the 2nd century A.D. According to some estimation, initially about 1000 t of gold and 3000 t of silver were concentrated within this ore deposit.

After twenty centuries of almost continuous exploitation of the Roşia Montană epithermal ore deposit centred on a phreatomagmatic breccia pipe structure (the Cetate breccia), the reserves would still be of more than 300 t of gold. These reserves were recently pointed out by S.C. Roşia Montană Gold Corporation S.A., a Romanian-Canadian mining company which is preparing an exploitation in a large open pit.

This mining project comprises the exploitation of the ore bodies hosted by the Cetate and Cărnic massifs, which surround the actual village. These two massifs, as well as other areas of the mining perimeter, contain important archaeological vestiges of the pre-Roman, Roman and medieval times. This is the reason why very important archaeological rescue diggings and studies were started before the initiation of the exploitation activity.

Thereby, the most important archaeological rescue programme from Romania, supervised by The Museum of National History of Romania (Bucharest). Developed since 2000 under the name of "National Programme *Alburnus Maior*". The aim is to identify and study the various archaeological sites from the surface (settlement areas, necropolis, sanctuaries, roads, workshops etc.) as well as underground vestiges, namely the various mining networks preserved in spite of modern mining re-openings. Several Romanian archaeological teams from various institutes, museums and universities are running diggings in the field at the surface, as well as an international team including French mining archaeologists, Romanian and German geologists specialized in the study of the underground and surface mining works.

After four years of survey (1999-2002) and three years of digging campaigns (2000 - 2002), the team of mining archaeologists engaged on the site is now able to present a first report regarding the dimension, the quality and the peculiarities of the mining vestiges. Beyond the wealth of this patrimony, the well-developed mining areas offer a coherent vision of the underground topography. It is centered upon the strategic choices of the miners regarding the shape of mining workings, their distribution, their connections and their relationships. These choices and the adoption of the typical plans of networks, of relatively standardized works (trapezoidal galleries, narrow stopings, chambers with pillars), associated to a rigorous follow-up of the mineralization markers (i.e. silicification), appear in an evident way in the reconstituted plans of the mine.

The plans of the ancient mining networks reveal a strategy of exploitation and a division of the mining space in a relatively regular shape in certain parts of the ore deposit. The topographic observations together with the chronological data supplied by the diggings emphasize a regular distribution of several exploitation zones in the Roman times, during 2nd and 3rd centuries. The exploitation zones are of comparable size and they are connected among them by major connection

⁹ Université Toulouse Le Mirail, 5, Allée Antonio Machado, 31520 Toulouse, France.

¹⁰ "Babeş-Bolyai" University of Cluj-Napoca, 1, Kogălniceanu Str., Dept. of Mineralogy, RO-3400 Cluj-Napoca, Romania. E-mail: <ctamas@bioge.ubbcluj.ro>.

ways, mainly inclined galleries dug into the barren rocks in order to facilitate the access, transport and the ventilation within the mining networks.

The ancient mining works were cross cut and sometimes enlarged by the modern mining re-openings. Since the 18th century, the archaeological artefacts were collected and preserved, i.e. tools and wooden equipments (ladders, propping, wheel of drainage), and especially waxed wooden tablets wearing written contracts, in Greek or Latin, between the miners and the mining contractors.

The Roşia Montană ore deposit is located in the Apuseni Mountains, part of the Carpathian chain. The Apuseni Mountains are divided into two major parts: the Northern and the Southern Apuseni Mountains; called also Metalliferous Mountains. The Tertiary calc-alkaline magmatism/volcanism from the Apuseni Mountains, tectonically controlled, generated porphyry copper systems and epithermal precious metals ore deposits, concentrated in the Metalliferous Mountains within the so-called Golden Quadrilateral.

The Roşia Montană gold-silver ore deposit is an epithermal one, according to its specific features, such as: the morphology of the ore bodies, gangue and ore mineralogy, types and spatial distribution of the hydrothermal alterations, as well as the geochemical association. The ore is centered on two dacite massifs known as Cetate and Cârnic Hills, remnants of a dacite dome (the Cetate dacite), and separated by a phreatomagmatic breccia pipe structure (the Cetate breccia). This phreatomagmatic breccia pipe structure controlled the genesis of the Roşia Montană ore deposit. At the ore deposit scale the genesis of the precious metals (gold, silver) ore bodies have been mainly controlled by a maar-diatreme environment. The most important ore body from the Roşia Montană ore deposit is the so-called Cetate breccia, which had a very high-grade ore. Besides this breccia pipe body, other smaller ore bodies were equally exploited: veins, stockworks, breccias, impregnations and placers.

The ancient exploitation focused especially on veins, breccias and stockworks. The study of the underground workings proved that the ancients followed different markers related to the occurrence of high ore grades, i.e. quartz veins, quartz-adularia veinlets, breccias (dykes or pockets), silicified zones, stockworks, as well as the occurrence of native gold mainly in open spaces or in quartz. The research and exploitation mining workings always focused in the highest ore grades of the ore deposit, using these markers. The most wanted ore bodies were flatly dipping and steeply dipping veins, and especially their intersections (the so-called "cross"), as well as the crosscutting areas among breccias and veins, which represent high-grade ore bodies.

The richness of the Roşia Montană ore deposit is due to the occurrence and superposition of various styles of mineralization. Thus, one may notice that within both Cetate and Cârnic dacite massifs there are high-grade mineralized structures (veins, breccias, stockworks), which were followed by the ancients from the surface by digging long descending galleries towards the "heart" of the massifs. Within the high-grade ore bodies the exploitation was developed on horizontal as well as on vertical by opening two or three close related mining levels (adits, stopings, wide rooms with or without pillars). Such well-defined exploitation areas were afterwards connected by inclined adits/planes. The underground mapping and sampling proved that the ancient exploitation stopped when the ore grades dropped or when the workings hit barren rocks.

During these diggings, most of the supposed ancient identified vestiges were finally dated from the Roman time. Radiocarbon dating obtained on wood fragments found in different mining networks emphasizes that the mining activity was already very important even from the pre-Roman (Dacian?) time in Rosia Montana. It then continued for a long time, then was revived and intensified under the control of the Roman administration.

The mining vestiges are very well preserved and present several peculiarities that allow us to identify them quickly. The ancient pre-Roman or Roman mining networks have systematically a trapezoidal section. The presence of lamp notches within adits, inclined planes or shafts are specific for Roman times. The mining workings have generally small dimensions, being always regularly calibrated and opened mostly by iron tools (pick and/or chisel). The silicified ore zones, with high hardness, were mined out by fire setting especially close to the surface where the natural ventilation was possible. The southern slope of the Cârnic massif as well as the southwestern part of the Cetate massif (the Găuri area) preserved fire-setting mining workings. The radiocarbon dating confirms the Roman age of these mining workings.

Among the different types of mining workings we may mention:

- Various types of drifts: short research drifts or long circulation drifts leading to different directions, or even exploitation drifts grouped next to each other in fan-shaped, star or chequer-work networks.
- Stopings or exploitation workings generally obtained by widening and deepening an older gallery.
- Upward or downward the galleries called “descending drifts”, dug from the surface or inside the massif between two isolated zones of exploitation. The floor of these galleries was often cut astairs.
- Chambers with pillars, the result of the exploitation by cross-cutting galleries within a limited space.
- Exploration and exploitation shafts with helical section, turning in right angle and containing stairs at the base. The vertical shafts between two levels are generally short and very rare. The ancient miners from the Roşia Montană preferred mostly inclined planes, with or without stairs as well as long or short descending drifts but not shafts.

The main mining workings at Roşia Montană remain the galleries. They are adapted to all the needs. In the current state of knowledge, few drainage systems have been observed, but the flooded levels situated in the deeper parts of the massifs have not been yet reached. The surveying should be pursued to help to find an answer to this important question.

Descending drifts connected to the surface should have done the ventilation of the deeper levels of the mine. The narrow sections of the galleries and their trapezoidal shape (wide base at the base) allowed to avoid mostly the wood propping of these small underground workings. In the high stopings, wood notches in the walls testify the wood propping placed transversely, necessary to support suspended wood floors. Despite the frequent occurrence of wood notches, the wood propping itself is not preserved in the upper part of the explored mining workings. On the other hand, in other parts of the site still not searched, as in the Orlea mining field wooden frames were found in place in the galleries and furthermore, in the Țarina mining field within a flooded shaft the wooden propping was recognized during the survey.

Using a division of the whole ore deposit into several concessions, the Roman State managed to develop the mining networks and of course to increase the production of precious metals. The archaeological discoveries suggest that the ancient mining networks represent in fact the final result of several decades of mining activity. The time span of Roman mining activity at *Alburnus Maior* lasted for about one century and a half, with a period of very intense exploitation of about half a century, which took place between the Roman conquest in 106 and the invasion of the Marcomans in 167. The Roman mining activity seems to succeed probably after a phase of pre-Roman (Dacian ?) mining one.

The study of the underground plans of the preserved mining networks allows to restore a well-defined spatial organization of the whole mine, as is the reflection of the economic planning regarding the production established by the Roman fiscal administration in order to improve the incomes for the Roman Empire. To study the Roşia Montana's mining workings, one needs to find their logic of development, conditioned by the nature of the ore deposit, by the type and the number of ancient mining developers, forced to work according to a program settled by laws.

COMPARATIVE STUDY OF THE TETRAHEDRITES FROM THE METALIFEROUS MTS. AND BAIJA MARE DISTRICT (ROMANIA) BASED ON MICROPROBE ANALYSES

Gheorghe Damian¹¹, Floarea Damian¹

Introduction

Tetrahedrites are the most frequent minerals in the Neogene hydrothermal mineralizations from Baia Mare district and of the Metaliferous Mts. The hydrothermal ore deposits from Baia Mare area and South Apuseni Mts. are related to Neogene magmatism.

In the Baia Mare district the metallogenetic activity would correspond to three distinct phases (Giuşcă *et al.*, 1973) during the Sarmatian-Pontian. The mineralizations have mainly base-metal character, subordinately gold-silver and copper occur. The upper parts of the copper and base-metal mineralizations have important quantities of gold and silver.

In the Metaliferous Mts. (South Apuseni Mts.) the metallogenetic activity is developed in two phases in the Badenian and the Sarmatian (Vlad & Borcoş, 1994). The mineralizations are epithermal mainly of gold-bearing type and subordinately gold/base-metal character. The ore deposits are related to the subvolcanic bodies and are represented by the porphyry and hydrothermal vein types. Tetrahedrites are dominantly present in the veins containing native gold, gold-telluride and base-metal $\pm$ gold.

Tetrahedrite occurrences

The most frequent occurrences of tetrahedrites in Baia Mare and Metaliferous Mts. are presented by Udubaşa *et al.* (1992). The majority of the investigated tetrahedrites from the Baia Mare area comprise Sb content. Tennantite have been identified in the eastern part of the Băiut and Căvnic district. The existence of tennantite has been proved in the copper mineralizations and in the upper zones of the veins. Tennantite is usually zoned, occurring together with tetrahedrite in parallel bands. Argentian tetrahedrites were identified at Ilba, Nistru, Herja, Băiut and Toroiaga. They are very frequent at Herja where the milimetric tetrahedral crystals associated with jamesonite (plumosite). These tetrahedrites have been identified only by microprobe analyses.

In the Metaliferous Mts. tetrahedrites associated with sulphides occur as grains smaller than 1 milimeter. Veinlets and tiny grains occur frequently in chalcopyrite, sphalerite and galena. Compact tetrahedrites masses are associated with silvanite at Săcărâmb.

The geochemistry of the tetrahedrites

Numerous tetrahedrite samples from several hydrothermal ore deposits from Baia Mare and Metaliferi Mts. have been investigated by microprobe analyses, (JCXA accelerating voltage of 25 kV ISIM Timişoara and Jeol SEM with EDS accelerating voltage 20 kV Université D'Aix Marseille III), (Table 1). Tetrahedrites from the Baia Mare district are rich in Sb while tetrahedrites from the Metaliferi Mts. contain a higher quantity in As. Some tetrahedrites from the copper mineralizations from the eastern Baia Mare district contain high amounts of As and they are zoned in respect to their As content. Besides the main elements (Cu, Sb, As, S), Zn and Fe reaching up to 7.69%, respectively 6.46%, were also determined in both areas. In Baia Mare district Zn is predominant in tetrahedrites and Fe is predominant in tennantite. In the tetrahedrites from the Metaliferous Mts. Zn is predominant compared with Fe. For the tetrahedrites from the Metaliferous Mts. a high content of Mn is characteristic. The presence of Mn in high quantities determines the separation of some Mn-rich tetrahedrite varieties.

¹¹ North University of Baia Mare, 62/A Victor Babeş Str., 4800 Baia Mare, Romania, <damgeo@univer.ubm.ro>.

All tetrahedrites contain Ag up to 2%. In some tennantite-tetrahedrite intermediate members from the gold-silver ore deposited in the Metaliferi Mts, the silver content reaches up to 11.85%. In the Baia Mare district the Ag-rich tetrahedrites contain up to 29.3% Ag. These tetrahedrites are specific for the base-metal ore deposits rich in Ag. The tennantites contain only small quantities of Ag, one exception is represented by the Tălagiu (Metaliferous Mts) tennantite-tetrahedrite intermediate member. The argentian tetrahedrites from the Baia Mare district always contain Fe in significant quantities as well as very small quantities of Zn.

An exception is an tetrahedrite from Băiut (Baia Mare district) which contains 21.3% Fe and also has high As content.

Table 1.

Chemical formula of the tetrahedrites based on microprobe analyses

Location	Chemical formula	Tetrahedrite members
Tarna	$\text{Cu}_{9.76}\text{Zn}_{1.34}\text{Fe}_{0.37}\text{Ag}_{0.12}\text{Sb}_{2.82}\text{S}_{13}$	Tetrahedrite with Zn
Ilba	$\text{Cu}_{6.29}\text{Zn}_{0.22}\text{Fe}_{1.63}\text{Ag}_{2.85}\text{Sb}_{3.77}\text{As}_{0.34}\text{S}_{13}$	Argentian tetrahedrite
Băiut	$\text{Cu}_{7.40}\text{Zn}_{0.92}\text{Fe}_{0.40}\text{Ag}_{1.50}\text{Sb}_{3.41}\text{As}_{0.93}\text{S}_{13}$	Argentian tetrahedrite
Cavnic-B	$\text{Cu}_{9.81}\text{Zn}_{1.98}\text{Fe}_{0.20}\text{Ag}_{0.10}\text{Sb}_{3.48}\text{As}_{0.85}\text{S}_{13}$	Tetrahedrite with Zn
Cavnic-B	$\text{Cu}_{10.64}\text{Zn}_{2.04}\text{Fe}_{0.25}\text{Ag}_{0.02}\text{Sb}_{3.51}\text{As}_{0.74}\text{S}_{13}$	Tetrahedrite with Zn
Kelemen Cavnic	$\text{Cu}_{10.06}\text{Zn}_{1.22}\text{Fe}_{0.07}\text{Ag}_{0.21}\text{Sb}_{3.54}\text{As}_{0.32}\text{S}_{13}$	Tetrahedrite with Zn
Roata	$\text{Cu}_{11.03}\text{Zn}_{0.42}\text{Fe}_{1.46}\text{Ag}_{0.06}\text{Sb}_{0.02}\text{As}_{5.41}\text{S}_{13}$	Tennantite
Robu	$\text{Cu}_{8.89}\text{Zn}_{1.12}\text{Fe}_{0.88}\text{Ag}_{1.78}\text{Sb}_{2.03}\text{As}_{0.60}\text{S}_{13}$	Argentian tetrahedrite
Văratec	$\text{Cu}_{12.87}\text{Fe}_{2.21}\text{Ag}_{0.34}\text{Sb}_{1.06}\text{As}_{4.23}\text{S}_{13}$	Tennantite
Cisma	$\text{Cu}_{10.51}\text{Zn}_{0.28}\text{Fe}_{1.77}\text{Ag}_{0.01}\text{Sb}_{3.20}\text{As}_{0.21}\text{S}_{13}$	Tetrahedrite with Fe
Cisma	$\text{Cu}_{8.66}\text{Zn}_{0.28}\text{Fe}_{0.96}\text{Ag}_{0.01}\text{Sb}_{3.01}\text{As}_{2.44}\text{S}_{13}$	Tetrahedrite-tennantite
Bandurița	$\text{Cu}_{6.58}\text{Zn}_{0.01}\text{Fe}_{5.06}\text{Ag}_{0.04}\text{Sb}_{0.23}\text{As}_{2.37}\text{S}_{13}$	Fe-bearing tennantite
Toroiaga	$\text{Cu}_{7.98}\text{Zn}_{0.70}\text{Fe}_{1.27}\text{Ag}_{2.41}\text{Sb}_{4.12}\text{S}_{13}$	Argentian tetrahedrite
Toroiaga	$\text{Cu}_{8.11}\text{Zn}_{0.56}\text{Fe}_{1.27}\text{Ag}_{2.41}\text{Sb}_{4.12}\text{S}_{13}$	Argentian tetrahedrite
Nistru	$\text{Cu}_{8.05}\text{Fe}_{2.06}\text{Ag}_{2.60}\text{Sb}_{4.88}\text{S}_{13}$	Argentian tetrahedrite
Herja 1	$\text{Cu}_{7.40}\text{Zn}_{0.92}\text{Fe}_{0.40}\text{Ag}_{1.50}\text{Sb}_{3.41}\text{As}_{0.93}\text{S}_{13}$	Argentian tetrahedrite
Herja 2	$\text{Cu}_{7.57}\text{Fe}_{1.82}\text{Ag}_{3.52}\text{Sb}_{4.24}\text{S}_{13}$	Argentian tetrahedrite
Herja 3	$\text{Cu}_{5.63}\text{Fe}_{2.21}\text{Ag}_{5.72}\text{Sb}_{4.74}\text{S}_{13}$	Freibergite
Herja 50	$\text{Cu}_{6.28}\text{Zn}_{0.20}\text{Fe}_{1.52}\text{Ag}_{3.45}\text{Sb}_{4.02}\text{As}_{0.01}\text{S}_{13}$	Argentian tetrahedrite
Herja 110	$\text{Cu}_{7.17}\text{Zn}_{0.23}\text{Fe}_{1.64}\text{Ag}_{2.60}\text{Sb}_{3.95}\text{As}_{0.03}\text{S}_{13}$	Argentian tetrahedrite
Băiut20	$\text{Cu}_{9.80}\text{Pb}_{0.004}\text{Zn}_{0.95}\text{Fe}_{1.06}\text{Ag}_{0.29}\text{Sb}_{2.68}\text{As}_{1.67}\text{S}_{13}$	Tetrahedrite-tennantite
Băiut 21	$\text{Cu}_{9.66}\text{Zn}_{0.73}\text{Fe}_{1.27}\text{Ag}_{0.22}\text{Sb}_{1.92}\text{As}_{2.17}\text{S}_{12.996}\text{Se}_{0.004}$	Tennantite-tetrahedrite
Băiut 22	$\text{Cu}_{9.65}\text{Zn}_{0.74}\text{Fe}_{1.28}\text{Ag}_{0.28}\text{Sb}_{1.84}\text{As}_{2.18}\text{S}_{12.988}\text{Se}_{0.012}$	Tennantite-tetrahedrite
Băiut 23	$\text{Cu}_{9.52}\text{Zn}_{1.12}\text{Fe}_{0.80}\text{Ag}_{0.36}\text{Sb}_{3.50}\text{As}_{0.67}\text{S}_{13}$	Tetrahedrite
Băiut 42	$\text{Cu}_{10.03}\text{Zn}_{1.00}\text{Fe}_{1.10}\text{Ag}_{0.10}\text{Sb}_{3.10}\text{As}_{1.038}\text{S}_{13}$	Tetrahedrite-tennantite
Băiut 43	$\text{Cu}_{9.77}\text{Zn}_{1.22}\text{Fe}_{0.77}\text{Ag}_{0.10}\text{Sb}_{3.11}\text{As}_{0.91}\text{S}_{19.988}\text{Se}_{0.012}$	Tetrahedrite-tennantite
Băiut 44	$\text{Cu}_{9.72}\text{Pb}_{0.004}\text{Zn}_{0.91}\text{Fe}_{1.03}\text{Ag}_{0.10}\text{Sb}_{2.88}\text{As}_{1.16}\text{S}_{12.988}\text{Se}_{0.012}$	Tetrahedrite-tennantite
Băiut 49	$\text{Cu}_{9.87}\text{Zn}_{1.29}\text{Fe}_{0.79}\text{Ag}_{0.06}\text{Sb}_{3.09}\text{As}_{1.12}\text{S}_{12.982}\text{Se}_{0.018}$	Tetrahedrite-tennantite
Băiut 50	$\text{Cu}_{9.78}\text{Pb}_{0.005}\text{Zn}_{1.08}\text{Fe}_{1.02}\text{Ag}_{0.05}\text{Sb}_{2.08}\text{As}_{2.04}\text{S}_{12.982}\text{Se}_{0.018}$	Tennantite-tetrahedrite
Băiut 56	$\text{Cu}_{9.97}\text{Zn}_{0.12}\text{Fe}_{1.87}\text{Ag}_{0.08}\text{Sb}_{2.12}\text{As}_{2.12}\text{S}_{13}$	Tennantite-tetrahedrite
Băiut 57	$\text{Cu}_{9.87}\text{Zn}_{1.19}\text{Fe}_{0.84}\text{Ag}_{0.05}\text{Sb}_{2.43}\text{As}_{1.75}\text{S}_{13}$	Tetrahedrite-tennantite
Barza-1	$\text{Cu}_{7.45}\text{Zn}_{1.65}\text{Fe}_{0.33}\text{Mn}_{0.95}\text{Ag}_{1.56}\text{Sb}_{3.89}\text{S}_{13}$	Tetrahedrite
Barza-2	$\text{Cu}_{7.57}\text{Zn}_{1.68}\text{Fe}_{0.48}\text{Ag}_{1.64}\text{Sb}_{3.84}\text{S}_{13}$	Argentian tetrahedrite
Barza-3	$\text{Cu}_{6.82}\text{Zn}_{0.98}\text{Fe}_{1.01}\text{Mn}_{1.31}\text{Ag}_{1.90}\text{Sb}_{3.93}\text{S}_{13}$	Argentian tetrahedrite
Tălagiu 1	$\text{Cu}_{8.91}\text{Zn}_{1.84}\text{Fe}_{0.34}\text{Ag}_{0.89}\text{Sb}_{3.06}\text{As}_{0.97}\text{S}_{13}$	Tetrahedrite-tennantite
Tălagiu 2	$\text{Cu}_{8.89}\text{Zn}_{1.81}\text{Fe}_{0.42}\text{Ag}_{0.92}\text{Sb}_{2.90}\text{As}_{1.06}\text{S}_{13}$	Tetrahedrite-tennantite
Coranda 1	$\text{Cu}_{8.70}\text{Zn}_{1.66}\text{Fe}_{0.27}\text{Mn}_{1.52}\text{Ag}_{0.23}\text{Sb}_{3.07}\text{As}_{0.91}\text{S}_{13}$	Tetrahedrite-tennantite
Coranda 2	$\text{Cu}_{9.58}\text{Zn}_{1.74}\text{Fe}_{0.21}\text{Mn}_{0.22}\text{Ag}_{0.19}\text{Sb}_{2.73}\text{As}_{1.40}\text{S}_{13}$	Tetrahedrite-tennantite
Coranda 3	$\text{Cu}_{8.32}\text{Zn}_{1.45}\text{Fe}_{0.26}\text{Mn}_{1.87}\text{Ag}_{0.24}\text{Sb}_{3.15}\text{As}_{0.80}\text{S}_{13}$	Tetrahedrite-tennantite
Săcărâmb	$\text{Cu}_{10.4}\text{Zn}_{1.53}\text{Ag}_{0.17}\text{Sb}_{1.12}\text{As}_{2.1}\text{S}_{12.86}\text{Se}_{0.14}$	Tennantite-tetrahedrite

Crystallochemistry of the tetrahedrites

The structural formula of the tetrahedrites is:

The structure of the tetrahedrites has been established by Pauling & Neuman (1934) and later confirmed by Wuench (1964). In all the analyzed samples (Table 1), the presence of up to two atoms of Fe and Zn is observed. These elements substitute Cu^{2+} . Only in the tennantite-tetrahedrite intermediate members, these elements substitute Cu^{2+} in approximately equal quantities. Zn substitutes Cu^{2+} in great quantities in the tetrahedrite end members, and Fe substitutes Cu^{2+} especially in the tennantites and in the argentian tetrahedrites. Because Fe and Zn do not exceed two atoms in the formula, it is obvious that they only substitute the Cu^{2+} . In the Metaliferi Mts. there are tetrahedrites with a high content in Mn. It is possible that this high content will also substitute Cu^+ because the sum of the bivalent elements (Zn, Fe, and Mn) is greater than the Cu^{2+} quantity.

Substitution of the Cu^+ by Ag is found only in tetrahedrite end members from Baia Mare and the Metaliferous Mts. Tetrahedrite-tennantite intermediate member from Tălagiu (Metaliferous Mts.) represents an exception. This fact is confirmed by the low content of silver in tennantite-tetrahedrite intermediate member in the gold-silver mineralizations from Săcărâmb. At low contents (under 2.5% Ag) the substitution of Cu^+ by Ag is achieved independently from the presence or absence of stibium. At high contents of Ag^+ , Fe^{2+} substitutes Cu^{2+} . In this case the substitution of Cu^+ which Ag^+ takes place at the same time with the substitution of Cu^{2+} with Fe^{2+} .

Different members of tetrahedrites were separated (Table 1). Most of the tetrahedrites are represented by the tetrahedrite-tennantite intermediate members. Most of the analyzed samples are tetrahedrite end members, because they contain $\text{Sb} > 3$ in formula (Table 1). Tetrahedrite end member is predominating in base metal ore deposits in the Baia Mare district. Some samples from Căvnic, Băiuţ and Săcărâmb correspond are tennantite, as they have more than 3 atoms of As in the formula. The tennantite is predominating in the metallogenetic field Băiuţ (in copper ore deposits) and in the Metaliferi Mts (Au-Ag-Te ore deposit).

The argentian tetrahedrites were identified at Ilba, Nistru, Herja, Băiuţ, Toroiaga, Gura Barza and Tălagiu. Only one tetrahedrite from Herja has $\text{Ag} > 4$ in the formula and can be named as freibergite. According to Riley (1974) the amount of 20% Ag delimits freibergite from the argentian tetrahedrite. In these cases, with the exception of two samples from Herja, which correspond to the freibergite, all the others can be assigned to argentian tetrahedrites.

Conclusions

The tetrahedrites from the base-metal mineralizations in the Baia Mare area and in the Metaliferous Mts. are represented by numerous members. It is for the first time that argentian tetrahedrite and freibergite have been proved to exist in Romanian occurrences, based on quantitative analytical data. The Mn tetrahedrite varieties from the Metaliferous Mts. are the first occurrences discovered in the world.

References

- Giuşcă, D., Borcoş, M., Lang, B., Stan, N. (1973) Neogene volcanism and metallogenesis in the Gutâi Mountains. *Excursion Guide IGR*, Bucharest 121 p.
- Pauling, L., Neuman, E.W. (1934) The crystal structure of binnite $(\text{Cu,Fe})_{12}\text{As}_4\text{S}_{13}$ and the chemical composition and structure of minerals of the tetrahedrite group. *Zeitschrift für Kristallographie* (Kristallgeometrie, Kristallphysik, Kristallchemie), **88**, 54-62.
- Riley, J.F. (1974) The Tetrahedrite - Freibergite Series, with reference to the Mount Isa Pb-Zn-Ag orebody. *Mineral Deposita*, **9**, 117-124.
- Udubaşa, G., Ilinca, Gh., Marincea, Şt., Săbău, Gh., Rădan, S. (1992) Minerals in Romania: State of the Art 1991. *Romanian Journal of Mineralogy*, **75**, 1-51.
- Vlad, Ş., Borcoş, M. (1994) Metallogenesis and plate tectonics in Romania, in Borcoş, M. & Vlad, Ş. (Eds.): Plate tectonics and metallogeny in the east Carpathians and Apuseni Mts, *Field Trip Guide* June 7-19, 1994, Geological Institute of Romania, Bucharest.
- Wuensch, B.J. (1964) The crystal structure of tetrahedrite, $\text{Cu}_{12}\text{Sb}_4\text{S}_{13}$. *Zeitschrift für Kristallographie* (Kristallgeometrie, Kristallphysik, Kristallchemie), **119**, 437-453.

Ag-SULPHOSALTS IN UPPER PARTS OF THE BAIJA SPRIE DEPOSIT (ROMANIA): MICROANALYSES AND IMPLICATIONS FOR DEPOSIT ZONALITY

Gheorghe Damian¹², Floarea Damian¹, Nigel J. Cook¹³, Cristina L. Ciobanu²

Introduction

The Baia Mare metallogenetic district is one of Europe's major metallogenic provinces and includes both gold and base-metal mineralization. Ores are associated with calc-alkaline Neogene volcanism. The calc-alkaline volcanism and associated metallogeny are related to an underlying pluton, (Borcoş et al., 1998). According to Giuşcă *et al.* (1973), metallogenetic activity corresponds to three distinct phases during the Sarmatian-Pontian.

The ore deposits in the Baia Mare district are mainly epithermal, of low sulphidation type. They are predominantly base-metal, but distinct Au-Ag and Cu types also occur. The higher levels of the Cu and base-metal ores contain important quantities of Au and Ag. The most important deposit in the district is Baia Sprie one.

Baia Sprie ore deposit – general data

This deposit is located along the regional E-W trending (Dragoş Vodă fault). The ore deposition is associated with the Jereapăn andesite eruptive phase (Giuşcă *et al.*, 1973) taking place within Pannonian (Edelstein *et al.*, 1992). The Principal Vein (5250 m in length) and the Nou vein are emplaced at the northern and southern limits of the andesite dyke respectively. Until 1972, the Baia Sprie ore deposit was known only in its western part, but exploration work proved its continuation eastwards of the Principal vein. The Pb-Zn mineralization includes some Au and Ag. Within the upper part of the deposit (Dealul Minei Hill), there are numerous branched veins and several stockwork-type ores, as well as some alignments parallel to the direction of the Main Vein.

The most comprehensive mineralogical studies on the Baia Sprie ore deposit are those by Tokody (1942), Manilici *et al.* (1965) for the Principal vein and Petruian *et al.* (1971) for the Nou vein. The Baia Sprie deposit is the type locality for andorite, felsöbányite, klebelsbergite, semseyite and szmikite, (Damian *et al.*, 1995). Despite the above studies, however, the occurrences of few sulphosalt minerals have been confirmed by microanalysis. We have studied samples from the upper part of the Principal vein and from branches of that vein, characterized by base-metal mineralization enriched in Au and Ag. Two typical parageneses appear in this zone: (i) sphalerite-galena-Ag-sulphosalts and (ii) pyrite-Ag-sulphosalts.

The mineralogy of Ag-sulphosalts

In the Pb- and Zn-ore zones fine-grained **acanthite**, associated with miargyrite and freibergite occurs (Fig. 1a).

Diaphorite occurs as µm-sized grains included within freibergite (Fig. 1b) and contains small quantities of Cu, Zn, and As, (Table 1). According to Tokody (1942), diaphorite appears associated with miargyrite.

Freieslebenite occurs as µm-sized grains including and associated with freibergite (Fig. 1b). In freieslebenite there are small impurities of Fe and As (Table 1).

Freibergite is mentioned from the Baia Sprie deposit for the first time in this study. It occurs associated both with sphalerite-galena and pyrite. The occurrence at Baia Sprie represents the second occurrence in the Baia Mare zone after Herja (Cook & Damian, 1997; Damian & Damian, 1999). It appears as sub-mm grains included in galena or associated with pyrite and other silver sulphosalts, (Fig. 1). Like freibergite from Herja the Baia Sprie one contains more Fe than Zn (Table 1).

¹² North University of Baia Mare, 62/A, V. Babeş Str., 4800 Baia Mare, Romania, E-mail: damgeo@ubm.ro

¹³ Geological Survey of Norway, N-7491 Trondheim, Norway.

In this case the double substitution of copper in the freibergite can be also noticed. In the freibergite structure Cu^{2+} is substituted by Fe^{2+} , and monovalent Cu^+ by Ag^+ . There is a direct correlation between the Fe and Ag contents.

Fig. 1. Back-scattered electron images of sulphosalt-rich assemblages, Baia Sprie deposit.

Abbreviations: Ach - acanthite, Tet - tetrahedrite, Mgy - miargyrite, Py - pyrite, Pgy - pyrargyrite, Pbs - polybasite, Prs - pearceite, Asp - arsenopyrite, Fsl - freiselebenite, Pyr - pyrargyrite-proustite, Dia - diaphorite, Qz - quartz.

Miargyrite occurs as submillimetric-sized grains associated with freibergite, pyrargyrite, polybasite and pyrite (Fig. 1a). The Baia Sprie miargyrite contains very minor As and the formula corresponds almost identically with the theoretical one.

According to Tokody (1942), **pyrargyrite** known from Baia Sprie, is associated with proustite, galena and sphalerite. We noticed very small grains of pyrargyrite associated with pearceite, miargyrite, freibergite and arsenopyrite (Fig. 1a).

Proustite is specific for the upper part from the western of Baia Sprie. In our samples, proustite occurs in the pyrite paragenesis associated with freibergite, diaphorite and freieslebenite. We note compositions from across the pyrargyrite-proustite join.

Pearcite-polybasite appears as a μm -sized grains associated with freibergite, pyragyrite, and miargyrite (Fig. 1a). This represents the first description of the mineral series from Baia Sprie. The formula corresponds with the theoretical one.

A further new occurrence for the deposit is **stephanite**, also confirmed by microanalysis.

Conclusions

Freibergite, stephanite and pearcite-polybasite represent the new occurrences in the Baia Sprie ore deposit. The presence of assemblages containing abundant Ag-sulphosalts in the upper part of the Baia Sprie deposit justifies the distinction made between gold and silver mineralization and further indicates the clear zonality of the Baia Sprie mineralization. The distribution of the mineralization in the Principal vein suggests its zonal character and the existence of the phases of mineralization. The presence of both Ag and Sb, in high quantities, within the hydrothermal solutions has determined the deposition of Ag-sulphosalts. The abundance of silver has also led to the appearance of freibergite rather than tetrahedrite (tennantite) as in the base-metal ore types. Formation of the minerals in this level has been realised in an open system after the boiling of the mineralising solutions.

Compositions of Pb- and Ag-bearing sulphosalts¹⁴, Baia Sprie ore deposit.

Mineral	Cu	Zn	Fe	Ag	As	Sb	Pb	S	Total	Formula [Fe, Zn, (Cu) ignored]
Pyrargyrite	0	0	0	57.74	0.78	22.68	0	18.92	100.12	Ag _{2.83} (As _{0.05} Sb _{0.99}) _{1.04} S _{3.12}
Pyrargyrite	0.12	0	0	61.21	0	20.89	0	17.76	99.98	(Cu _{0.01} Ag _{3.07}) _{3.08} Sb _{0.93} S _{2.99}
Pyrargyrite-proustite	0.08	0	0	59.42	6.90	12.39	0	20.53	99.32	(Cu _{0.01} Ag _{2.78}) _{2.79} (As _{0.51} Sb _{0.47}) _{0.98} S _{3.23}
Proustite	1.24	0	0	62.14	13.96	0.15	0	20.65	98.14	(Cu _{0.10} Ag _{2.83}) _{2.93} (As _{0.01} Sb _{0.91}) _{0.92} S _{3.16}
Proustite	0	0	0	63.53	13.54	1.17	0	19.96	98.20	Ag _{2.94} (As _{0.05} Sb _{0.90}) _{0.95} S _{3.11}
Diaphorite	0.13	0	0	22.56	0	27.49	31.12	18.45	99.75	Ag _{2.91} Pb _{2.07} Sb _{3.11} S _{7.93}
Diaphorite	0.26	0.38	0	22.05	0.21	26.69	32.03	18.96	100.58	Ag _{2.79} Pb _{2.11} (As _{0.04} Sb _{2.99}) _{3.03} S _{8.07}
Diaphorite	0.61	0	0	23.75	0	29.17	30.33	19.45	101.33	Ag _{2.94} Pb _{1.96} Sb _{2.99} S _{8.11}
Miargyrite	0	0	0	35.38	0	41.28	0	22.33	98.99	Ag _{0.96} Sb _{1.00} S _{2.04}
Miargyrite	0	0	0	36.64	0.17	40.98	0	21.57	99.36	Ag _{1.01} (As _{0.01} Sb _{1.00}) _{1.01} S _{1.99}
Freislebenite	0	0	1.06	18.48	0	23.36	41.29	17.92	102.11	Ag _{0.92} Pb _{1.07} Sb _{1.03} S _{2.99}
Freislebenite	0	0	0	20.03	0.10	24.05	40.17	17.23	101.58	Ag _{1.00} Pb _{1.04} (As _{0.01} Sb _{1.06}) _{1.07} S _{2.89}
Pearceite	0.39	0	0	75.41	6.14	0.34	0	15.70	97.98	(Cu _{0.14} Ag _{15.85}) _{15.98} (As _{1.86} Sb _{0.06}) _{1.92} S _{11.10}
Stephanite	0	0	0.49	65.01	0.07	17.13	0	15.69	98.38	Ag _{4.89} (As _{0.01} Sb _{1.14}) _{1.15} S _{3.97}
Acanthite	0.10	0	0	86.03	0	0	0	12.08	98.21	Ag _{2.04} S _{0.96}
Tetrahedrite	22.73	3.35	6.01	19.31	0.20	24.72	0	23.08	99.40	(Cu _{6.40} Ag _{3.20})(Fe _{1.93} Zn _{0.92})(As _{0.05} Sb _{3.63})S _{12.88}
Freibergite	15.81	2.57	5.45	26.42	0.19	24.95	0.48	22.16	98.03	(Cu _{4.71} Ag _{4.64})(Fe _{0.74} Zn _{1.85})(As _{0.05} Sb _{3.88})S _{13.09}

References

- Borcoş, M., Vlad, Ş., Udubaşa, Gh., Găbudeanu, B. (1998) Qualitative and Quantitative Metallogenetic Analysis of the Ore Genetic Units in Romania, *Rom. J. Mineral. Deposits*, **78**, Spec. Issue, 1-157.
- Cook, N.J., Damian, Gh. (1997) New data on plumosite and other sulphosalt minerals from the Herja hydrothermal vein deposit, Baia Mare district, Romania, *Geologica Carpathica*, **48**, 387-399.
- Damian, Gh., Nedelcu, L., Istvan, D. (1995) Two representative vein deposits (Au-Ag and Pb-Zn) related to Neogene volcanic structures. *Rom. J. Mineral.*, **77**, Suppl. 2, 45-63.
- Damian, Gh, Damian, FI (1999) New data of the freibergite and argentian tetrahedrite in hydrothermal veins from Baia Mare mining district. *Anal. Univ. Bucureşti*, **XLVIII**, 14-16.
- Edelstein, O., Bernard, A., Kovács, M., Crihan, M., Pecskay, Z. (1992) Preliminary date regarding the K-Ar ages of some eruptive rocks from Baia Mare Neogene volcanic zone. *Rev. Roum. Geol. Geogr. Geophys., Ser. Geol.*, **36**, 45-60.
- Giuşcă, D., Borcoş, M., Lang, B., Stan, N. (1973) Neogene volcanism and metallogenesis in the Gutâi Mountains. *Excursion Guide*, IGR Bucharest.
- Manilici, V., Giuşcă, D., Stipol, V. (1965) The study of the ore deposit from Baia Sprie (Baia Mare region) (in Romanian). *Mem.Com. Stat Geol.*, **VII**, 113 p.
- Petrulian, N., Steclaci, L., Oroveanu, F., Ştefan, H., Cioran, A. (1971) Mineralogical and geochemical study of the Nou vein from Baia Sprie (in Romanian), *St. Cerc. Geol., Geof., Geogr., Ser. Geol.*, **XVI**, 2, 2-16.
- Tokody, L. (1942) Minerals of Baia Sprie from geochemical view point (in Hungarian). *Math. t.-t. Ért.*, **61**, 324-386.

¹⁴ Analised in the Labotatory of the Geological Survey of Norway, N-7491 Trondheim, Norway

FLUID INCLUSIONS AND MORPHOLOGY OF QUARTZ FROM THE NEOGENE EPITHERMAL DEPOSITS, BRAD-SĂCĂRÂMB AREA (TRANSYLVANIA, ROMANIA)

Ágnes Gál¹⁵¹, Ferenc Molnár¹⁶², Lucreția Ghergari¹, Csaba Szabó¹⁷³, István Gatter²

The Brad-Săcărâmb basin is situated in the southwestern part of the Transylvanian Metaliferi Mountains and hosts several low-sulphidation type Au-base metal deposits. The host rocks of the mineralization are mainly Neogene andesites and dacites and subordinately Cretaceous to Neogene sandstones, shales and pelitic-psammitic sediments (Ivanovici *et al.*, 1969). The Neogene volcanic rocks were affected by two stages of hydrothermal alteration. The first stage is represented by propylitisation processes, the second stage is characterised by sericitisation, local adularitisation and argillitisation.

The main gangue minerals in the veins are quartz, Ca-Mn carbonates and barite. The predominant ore minerals are pyrite, galena, sphalerite, chalcopryite, stibnite, alabandine, realgar, tetrahedrite, buhornite, boulangerite and Au, Ag, Pb, Sb and Hg tellurides. The gold and silver-bearing minerals formed in association with galena and sphalerite. In some places, an early stage of Au-Ag and a later stage of Pb-Zn (with only little amounts of Au and Ag) mineralizations can be distinguished. Sulphosalts and chalcopryite, as well as carbonates are the latest minerals, following the earlier gold-bearing stages.

Quartz is a frequent hydrothermal mineral, which shows certain crystal forms under certain physico-chemical conditions (Molnár 1986, 1993; Hurai & Stersko 1987). In addition, fluid inclusions trapped during the growth of the quartz crystal can also be used for the estimation of the P-T-X conditions.

The crystal morphology and the fluid inclusions in quartz associated with the Pb-Zn (Au-Ag) stage of mineralization at the Săcărâmb, Coranda, Bocșa, Valea Morii Veche and the Carpen deposits were studied. Two different types of quartz were distinguished: a) vein quartz forming comb-textured aggregates with free crystal terminations pointing towards the center of the veins and b) euhedral quartz occurring in the vugs and cavities of the host rock of the veins. The crystal forms of the both types of quartz are: m prism (10^{-10}), positive r and negative z rhombohedra (10^{-11}), and rhombohedra with high indices {(h.0. $\bar{h}$.l), l (20^{-21}), l (02^{-21}), M (30^{-31}), 'M (03^{-31}), γ (40^{-41}), ' γ ($04\bar{4}1$), e ($50\bar{5}1$), 'e ($05\bar{5}1$), ξ ($60\bar{6}1$), ' ξ ($06\bar{6}1$), φ ($70\bar{7}1$), ' φ ($07\bar{7}1$), Ψ ($11.0.\bar{1}\bar{1}.1$), Ψ ($0.11.\bar{1}\bar{1}.1$)}. These rhombohedra with high indices appear as stripes forming a staggered habit.

The quartz crystals were cut parallel to the "c" axis; double polished thin sections were prepared for the fluid inclusion study. No clearly defined growth zoning was observed in the quartz crystals. A spider web structure, typical for the epithermal quartz crystals (Bodnar *et al.*, 1985) was recognized. Liquid-rich and vapor-rich primary aqueous fluid inclusions occur inside the crystals, as the growth of the crystals took place during the boiling of the hydrothermal fluids. Microthermometry was carried out on the liquid-rich two-phase fluid inclusions only. The first melting temperatures were noticed between -18 and -22°C, indicating a NaCl - dominated aqueous brine. The melting of the last ice crystal occurred in the range from -4.1 to -0.3°C apparently corresponding to a salinity between 0.5 and 6.6 wt% NaCl eqv. (Bodnar, 1993). The minimum homogenization temperatures of the liquid phase for the vein quartz are 250 and 260°C (the Săcărâmb-Bernard level and the Valea Morii Veche, respectively). The fluid inclusions from the vein quartz show the lowest values of the average apparent salinity: 1.9 – 1.4 NaCl eqv. wt%. The minimum temperatures of the

¹⁵ "Babeş-Bolyai" University of Cluj-Napoca, 1, Kogălniceanu Str., Dept. of Mineralogy, RO-3400 Cluj-Napoca, Romania. E-mail: <galagi@bioge.ubbcluj.ro>.

¹⁶ Department of Mineralogy, Eötvös Loránd University, Pázmány sétány 1/c, 1117 Budapest, Hungary.

¹⁷ Department of Petrology and Geochemistry, Eötvös Loránd University, Pázmány sétány 1/c, 1117 Budapest, Hungary.

homogenization of the liquid phase for the euhedral quartz from the wall rocks are ranging from 180 to 230°C, while the salinities are between 2.6 and 4.5 wt% NaCl .

The frequency distribution of the crystal forms shows no difference between the comb-textured quartz aggregates or the vein quartz. The step habit is the result of the variability of the fluid chemistry during the crystallization.

Although the textural features of the fluid inclusions suggest that both, the vein quartz and the comb-textured aggregates are the result of a boiling system, in the T_h/T_{mice} (Homogenization temperature vs. Ice melting temperature) diagram only the vein quartz points on boiling processes.

According to Haas diagram (1971), the calculated pressure ranges a wide domain, between 5 and 72 bar.

References

- Bodnar, R.J. (1993) Revised equation and table for freezing-point depression of H₂O-salt fluid inclusions. *Geochimica and Cosmochimica Acta*, **57**, 683-684.
- Bodnar R.J., Reynolds, T.J., Kuehn, C.A. (1985) Fluid inclusion systematic in epithermal system. *Rev. in Econ. Geol.*, **2**, Geology and geochemistry of epithermal systems, 73-88.
- Haas, J.L. (1971) The effect of Salinity on the Maximum Thermal Gradient of a Hydrothermal System at Hydrostatic Pressure, *Ec. Geol.*, **66**, 940-946.
- Hurai, V., Stresko, V. (1987) Correlation between quartz crystal morphology and composition of fluid inclusions as inferred from fissures in Central Slovakia (Czechoslovakia), *Chemical Geology*, **61**, 225-239.
- Ianovici, V., Giuşcă, D., Ghiţulescu, T. P., Borcoş, M., Lupu, M., Bleahu, M., Savu, H. (1969) The geological evolution of the Metaliferous Mts. (in Romanian). *Ed. Acad.*, 741 p., Bucureşti.
- Molnár, F. (1986) Morphologic and genetic research on quartz crystals relating to Paleogene-Neogene ore genesis (in Hungarian, with English abstract). *MSc. Thesis*, Dept. of Mineralogy, "Eötvös Lorand" University, Budapest.
- Molnár, F. (1993) Genesis of epithermal mineralization of Tokaj Mts. on the basis of fluid inclusion studies (in Hungarian, with English abstract). *Ph.D. Thesis*, Dept. of Mineralogy, "Eötvös Lorand" University, Budapest.

THE CHARACTERISTICS OF QUARTZ CONCENTRATE OBTAINED BY THE FLOTATION OF THE QUARTZO-FELDSPATIC RAW MATERIAL FROM BEDECIU DEPOSIT (CLUJ DISTRICT, ROMANIA)

Maria Gorea¹⁸, Cristina Mariş¹⁹

Quartz is one of the most important industrial raw materials used for ceramics, glass, mortar, etc. Its suitability and technological qualities are directly connected to the chemical composition (SiO_2 and iron oxides). The technology of the feldspar processing applied in the case of the quartzo-feldspatic rocks from the Bedeciu deposit involves a stage of quartz flotation and separation. This paper focuses on the technological, chemical, mineralogical and particle size distribution of quartz separated by flotation with the object of using it in various industrial products.

The Bedeciu deposit is located at the border between two major structural units, the Gilău Mountains and the Huedin Basin (included within the Transylvanian Depression), in the northern part of the contact area between intrusive rocks (the Muntele Mare granite) and crystalline schists of the metamorphic Someş series.

The main stages of the preparation technology consist of coarse (< 60 mm) and fine grinding (< 0.9 mm), followed by the 1st flotation stage (for the separation of micas and Fe-containing minerals) and the 2nd flotation stage (for the separation of feldspars). The drying, deferrisation, and packaging close the preparation cycle.

The raw material as well as the concentrated products i.g. micas, feldspar and quartz were chemically²⁰ and mineralogically²¹ analysed. The chemical composition of the raw material, of micas resulted after the 1st flotation stage, of feldspar obtained after the 2nd flotation stage and of quartz concentrate is presented in table 1. The mineralogical composition of the raw material and of quartz-concentrate is presented in table 2.

Table 1.

Chemical composition of the raw material and mineral-concentrates.

Sample (product)	Chemical composition [% oxides]							
	SiO_2	Al_2O_3	Fe_2O_3	Na_2O	K_2O	CaO	MgO	L.O.I.
Raw material	75.05	14.78	0.64	3.75	3.50	0.56	0.25	0.87
Micas concentrate	52.88	29.50	2.50	1.00	8.60	0.63	0.45	4.11
Feldspar concentrate	71.62	16.45	0.19	6.00	4.40	0.46	0.10	0.44
Quartz concentrate	92.28	2.74	0.15	1.3	1.30	0.46	0.41	0.43

Table 2.

Semiquantitative mineralogical composition of the raw material and quartz-concentrate.

Sample	Mineralogical composition [%]						
	Micas	Na-feldspar	K-feldspar	Ca-feldspar	Quartz	Limonite	Kaolinite
Raw material	7.10	36.63	18.17	1.64	33.85	0.68	1.72
Quartz concentrate	1.52	12.68	5.18	0.79	77.22	0.16	1.72

¹⁸“Babeş-Bolyai” University of Cluj-Napoca, 1, Kogălniceanu Str., Dept. of Mineralogy, RO-3400 Cluj-Napoca, Romania. E-mail: <mgorea@bioge.ubbcluj.ro>.

¹⁹ S.C. Cominex Nemetalifere S.A. – 1, Principală Str., 3452 Căpuş, Romania.

²⁰ Classical method; Chem. Lab. of S.C. Cominex Nemetalifere S.A., Căpuş.

²¹ Dron 2-type X-Ray powder diffractometer (Lab. of S.C. Procema S.A. Cluj-Napoca).

The distribution of the particle size of quartz obtained after flotation (intermediate product) and quartz concentrate obtained after drying and deferrisation (final product) is given in table 3.

Table 3.

Particle size distribution²² of the quartz concentrates.

Sample	Particle size distribution [%]				
	+ 500 µm	+ 250 µm	+ 200 µm	+ 90 µm	+ 40 µm
Intermediate product of quartz concentrate	0.50	18.00	15.00	56.00	8.00
Final product of quartz concentrate	0.50	8.00	53.00	25.00	2.50

According to the granulometry and chemical composition, especially Fe₂O₃ content, quartz concentrate obtained by the flotation of quartzo-feldspatic rocks from the Bedeciu deposit can be used as raw material for optical or crystal glass, for ceramics or for preparing dry mortars as well as other building materials.

²² Fritsch-Analysette 22 type Laser Granulometer (SC Faimar Baia Mare).

REACTION CORONAS AROUND QUARTZ XENOCRYSTS IN THE BASALTIC ANDESITE FROM DETUNATA (APUSENI MOUNTAINS), ROMANIA

Nicolae Har²³

Reaction coronas consisting of Al-poor pyroxene, tridymite and glass are developed around quartz xenocrysts of sedimentary origin hosted in olivine - bearing basaltic andesites from Detunata (Apuseni Mts. - Romania). The quartz xenocrysts reacted with the basaltic melt. The outer parts of the reaction coronas consist of a glassy matrix containing prismatic augite crystals developed perpendicular to the interface between the coronas and basaltic host rock. The inner parts of the coronas consist of isolated crystals of augite, fragments of quartz and tridymite crystals in a glassy groundmass. Tridymite is also present in fractures of the quartz xenocrysts. Microprobeanalyses and Raman spectroscopy were used to identify the composition of the coronas pyroxenes, glass, and silica polymorphs. The coronas pyroxenes have a homogeneous composition, typical for augite. The glass is highly siliceous ($\text{SiO}_2 = 72.0 - 76.8 \text{ wt.}\%$) as the result of quartz dissolution, and high in alkalis ($\text{Na}_2\text{O} = 1.38 - 3.22 \text{ wt.}\%$; $\text{K}_2\text{O} = 4.72 - 6.23 \text{ wt.}\%$) and aluminum ($\text{Al}_2\text{O}_3 = 9.31 - 12.18 \text{ wt.}\%$). Tabular, twinned crystals of tridymite show a high content of alkalis ($\text{Na}_2\text{O} + \text{K}_2\text{O} = 0.30 - 0.39 \text{ wt.}\%$) and aluminum ($\text{Al}_2\text{O}_3 = 0.54 - 0.86 \text{ wt.}\%$). Sodium and potassium ions appear to fill structural channels and voids in the tridymite structure, charge balanced by the substitution of Al for Si. The Raman spectra of tridymite show the most representative peaks, of 403 and 422 cm^{-1} .

The diffusion processes and temperature were the controlling factors in the genesis of the newly-formed minerals in the reaction zones. Si^{4+} diffused from the quartz lattice, while Ca^{2+} , Mg^{2+} , Na^+ and K^+ migrated from the basaltic-andesitic melt towards the coronas.

²³ "Babeş-Bolyai" University of Cluj-Napoca, 1, Kogălniceanu Str., Dept. of Mineralogy, RO-3400 Cluj-Napoca, Romania. E-mail: <har@bioge.ubbcluj.ro>.

HERCYNITE AND MAGNETITE IN THE HORNFELS XENOLITHS HOSTED BY GRANODIORITE FROM VALEA LUNGII (CLUJ COUNTY, ROMANIA)

Nicolae Har²⁴, Ramona Năstase¹

Isolated intermediate magmatic bodies of Upper Cretaceous – Paleogene age crops out in the Valea Drăganului – Valea Lungii area (northern part of the Apuseni Mountains, Romania). The petrography is mainly granodioritic, typical for hipabyssic facies. They contain frequently xenoliths from the basement of the area. In the intrusive granodioritic body from Valea Lungii two different types of xenoliths are present: of magmatic (porphyritic granodiorite) and metamorphic (gneiss) origin. The xenoliths are variable in size (up to 30 cm in diameter) and usually dark in color. The study is focused on the xenoliths of metamorphic origin, that have undergone pyrometamorphic transformations under the influence of the host melt. The resulted hornfels preserves the texture of the initial gneiss. Two different varieties of hornfels are present: spinel andalusite and spinel sillimanite hornfels. Both include different amounts of K-feldspar, biotite, phlogopite, magnetite and hercynite.

Based on the mineralogical assemblages of the spinel hornfelses the following remarks can be made:

- the xenoliths' external zone are dominated by sillimanite, associated with K-feldspar, biotite, phlogopite, magnetite and hercynite;
- in the xenoliths' inner part, andalusite is present and sillimanite is missing;
- biotite is associated mainly with hercynite. The Mg – Fe biotite becomes more Mg – rich (phlogopite) due to temperature increasing and thus, the phlogopite is associated with magnetite resulted from the biotite alteration;
- the Al - silicate polymorphs (andalusite and sillimanite) indicate thermal conditions of 650 – 750°C and a low pressure, typical for the pyroxene hornfels facies;
- Al –silicates polymorphs are the result of the reaction between muscovite and quartz content of the gneiss:

- the absence of muscovite and quartz in the hornfels suggest that the above reaction took place;
- the presence of magnetite and hercynite in the resulted hornfels can be explained due to the transformation of the Mg-Fe rich biotite into a Mg-rich biotite (phlogopite), at high temperature.

²⁴ "Babeş-Bolyai" University of Cluj-Napoca, 1, Kogălniceanu Str., Dept. of Mineralogy, RO-3400 Cluj-Napoca, Romania. E-mail: <har@bioge.ubbcluj.ro>.

RECENT MINERAL DEPOSITION IN THE CRATER OF THE CIOMADUL QUATERNARY VOLCANO - HARGHITA MOUNTAINS (ROMANIA)

Olivier Hercot¹²⁵, Ioan Seghedi²²⁶, Jean Naud¹, Razvan Caracas¹

At the margin of Sf. Ana crater, on the Ciomadul volcano, representing the morphological mark of the youngest volcano in south-eastern Europe (ca. 30 ka), a recent deposition of sulphates and carbonate has been found. The deposition drapes a cavity of a supporting wall on the right side of the road, which leads into the interior of the crater. The detected sulphates are **thenardite** [Na_2SO_4] and **aphthitalite** [$\text{K}_3\text{Na}_2(\text{SO}_4)_2$], rarely found in natural conditions. A rare carbonate, **nahcolite** [NaHCO_3] was also detected associated with the sulphates. Since the mineral assemblage is intimate and has low crystallinity it was not possible to isolate each mineral phase for detailed analyses. Further collection and observation will be done in the future for better characterisation of this recent mineral deposition.

²⁵ Université Catholique de Louvain, Departament de Geologie, Faculte des Sciences, Belgique.

²⁶ Institute of Geodynamics, 19-21, J.-L. Calderon Str., Bucharest 70201, Romania. E-mail: <seghedi@geodin.ro>.

MINERALOGY OF A “SACRED MONSTER”: THE DITRĂU ALKALINE MASSIF, EAST CARPATHIANS (ROMANIA)

Paulina Hirtopanu²⁷, Gheorghe Udubaşa¹

Unique in the whole area of the Carpathians, the Ditrău alkaline massif (DAM) displays both a great variety of contrasting lithologies and a big number of mineral species. Rocks from ultrabasite to high alkaline types, as well as granites, contact metamorphic assemblages, porphyry-style ores (Aurora), veins with sulphides, oxides and various REE minerals are known to be concentrated on a relatively small area, of only about 100 sq.km.

Due to contrasting host lithologies and various mineral forming processes the mineralogy of DAM is very complex. Nearly all the mineral classes have here representatives, including rare and very rare mineral species such as zirconosilicates, radioactive carbonates, sapphirine, Bi-sulphosalts, nearly all the species of the pyrochlore and perovskite groups etc, some of them forming single occurrences in Romania, e.g. hemusite, sederholmite, tungstenite, mosesite, senaite, srilankite, minrecordite, rosenbuschite, eudyalite, lovozerite, hiortdahlite, donpeacorite, sadanagaite, vishnevite, loparite etc. (Hirtopanu, 1998). The total number of mineral species is about 250 thus reaching the status of a “sacred monster” (Udubaşa, 1994). Nevertheless, there are both very frequent minerals such as nepheline, cancrinite, sodalite, various feldspars (including Ba-feldspars), amphiboles (K-pargasite, sadanagaite, magnesiosadanagaite, eckermannite, nyböite, ferrinyböite, richterite, Li-fluorarfvedsonite, Li-fluorkataphorite, Li-ferriECKermannite, etc.) and very rare or “minor minerals” such as isocubanite, ferromolybdate, strüverite, wöhlerite, zirkelite, betafite, ytrobetafite, willemseite, huttonite, cheralite, astrophyllite, aenigmatite, durangite etc. (Hirtopanu, 1998).

The knowledge of mineralogy of DAM began early in the XIX century, undoubtedly with the more frequent minerals, i.e. nepheline (called eleolite due to its greasy appearance, giving the rock name “eleolite syenite” or “ditroite”; Zirkel, 1866), sodalite, cancrinite, haüyne, etc. Later on other less frequent minerals were described, such as baddeleyite, zircon, bastnäsite, parisite, various complex oxides such as pyrochlore (group), fersmite, ixiolite, tantalite; pyrophanite; rhönite, wöhlerite; monazite and xenotime; mackinawite, bismuthinite, molybdenite, etc.

Finally some other mineral species (about 100) were added to the list; first of all the zirconosilicates (rosenbuschite and baratovite, catapleiite, eudyalite, låvenite, kupletskite), sapphirine (as big grains of centimetre size associated with corundum, spinel and cordierite), a series of REE bearing carbonates (synchysite, cordylite, hydroxylbastnäsite-(Ce), hydroxylbastnäsite-(Nd), hydroxylbastnäsite-(La), yttrparisite), goyazite, a number of rare sulphides and sulphosalts such as geerite, lautite, sederholmite, kesterite, gallite, briartite, arsenosulvanite, tungstenite etc. (Hirtopanu, 2000, *unpubl.*; Hirtopanu *et al.*, 2003). Thus there is an appreciable increase of mineral number from 25 in 1966 and 85 in 1977 to about 250 nowadays. Further increase is expected as the alkaline massifs worldwide permanently provide about one quarter of the new minerals proposed to the CNMMN of IMA.

References

- Hirtopanu, P. (1998) Grant report Romanian Academy, Ditrău. *IGR Archives*, Bucharest.
Hirtopanu, P. (2000) Report GIR Bucharest, Ditrău. *IGR Archives*, Bucharest.
Hirtopanu *et al.* (2003) Some old and new minerals from the Ditrău alkaline massif and their genetical implication. Abstr. vol. „*Petrology-Global Context*” (*Anniv. symp. dedicated to Acad. Dan Rădulescu*), Univ. Bucharest, p. 29.
Udubaşa, G. (1994) Sacred monsters of mineral occurrences in Romania: past, present, future. *Anal. Univ. Bucureşti*, **XLIII**, Suppl., Abstr. vol., 34-35.
Zirkel, F. (1866) *Lehrbuch der Petrographie*, Bonn, 595 p.

²⁷ Geological Institute of Romania, Bucharest, Romania. E-mail: <Paulina.Hirtopanu@igr.ro>.

TYPE OF CHONDRULES FROM ROMANIAN FALLEN ORDINARY CHONDRITES

Gabriel Ovidiu Iancu²⁸, Corina Ionescu²⁹

The chondrites (stony meteorites) are believed to represent the oldest rocks in the solar system and they are characterized by the presence of small mineral spheres called *chondrules*. The chondrule texture represents one of the easier criteria used for the petrologic-chemical classification of all chondrite main categories (O -ordinary, E-enstatite, C-primitive/carbonaceous chondrites, Rumuruti -type and Kakangari -type meteorites).

Based on the metal content and mineralogical composition, the ordinary chondrites are subdivided into three distinct groups, designated as H, L, and LL chondrites. The H-chondrites have high iron content, about 25-31% (including 12-21% metallic iron), the L-chondrites have lower iron content, ranging from 20 up to 25% (with 5-10% metallic iron) and the LL-chondrites contain 19-22% iron (with only 2-3% metallic iron).

Six petrographic types of chondrites, numbered from 1 to 6, are recognized. The ordinary chondrites show four petrographic types (numbered from 3 to 6), indicating progressive stages of thermal metamorphism.

Based on texture, Van Schmus & Wood (1967) proposed 4 categories of chondrules for each chondrite petrographic type: very sharply defined chondrules (type 3; e.g. H3, L3, LL3), well-defined chondrules (type 4; i.g. H4, L4, LL4), readily distinguished chondrules (type 5; e.g. H5, L5, LL5) and poorly defined chondrules (type 6; i.g. H6, L6, LL6). A fifth category (type 7), naming a melted type, was added (e.g. H7, L7, LL7).

Up to now, 7 fallen ordinary chondrites have been registered and investigated in Romania and various scientists previously classified them as follows: Mezö-Mădăraş (syn. Mădăraş) – L3, Ohaba – H5, Kakowa (syn. Cacova) – L6, Zsadany (syn. Jădani) – H5, Mocs (syn. Mociu) – L5-6, Sopot – H4, Tăuţi – L6. The single found ordinary chondrite, Tuzla, was classified as L6 type (Stanciu & Stoicovici, 1943; Graham *et al.*, 1985; Miura *et al.*, 1995; Iancu, 2003).

In order to determine the type and size of chondrules from Romanian fallen meteorites, samples from some Romanian and Austrian museums were examined with optical polarizing microscope as well as with JEOL JSM-5400 and JSM-6400 scanning electron microscopes.

In H3, L3 and LL3 chondrites, the chondrules are very sharply defined and closely packed. According to Norton (2002), within each chondrite group the chondrules can vary considerably in size. In the ordinary chondrites, the chondrules range from few microns to one centimeter. For the H, L and LL groups, the 300 µm, 500 µm, and 600 µm diameter are prevailing, with an average size of about 450 µm for all three groups. The mean diameter tends to vary with the type of chondrules. Most of the chondrules of H, L and LL classes show remarkable mineral consistency, being composed of olivine or/and pyroxene. Although they present apparently a bewildering array of textural forms, close examination reveals, acc. to Gooding & Keil, (1981) only seven basic textural types, with some variation within each type:

- PO - porphyritic olivine;
- PP - porphyritic pyroxene;
- POP - porphyritic olivine-pyroxene;
- RP - radial pyroxene;
- BO - barred olivine;
- C – microcrystalline;
- GOP - granular olivine-pyroxene.

²⁸ University "Al. I. Cuza" of Iasi, Department of Mineralogy and Geochemistry, 20 A Carol I Blv., RO-6600 Iaşi, Romania, E-mail: <gabiiiancu@geo.uaic.ro>.

²⁹ University "Babes-Bolyai" of Cluj-Napoca, Romania, Department of Mineralogy, 1 Kogălniceanu Str., RO-3400 Cluj-Napoca, E-mail: <corinai@bioge.ubbcluj.ro>.

The microscopic study in thin sections revealed the presence of the following types of chondrules in the Romanian fallen chondrites:

1. Mezö-Mădăraş (syn. Mădăraş) meteorite: very sharply defined PO, POP, RP, BO and GOP chondrules, ranging in size from 300 µm up to 600 µm;
2. Ohaba meteorite: readily distinguished PO, RP, BO (sometimes polysomatic) and GOP chondrules, ranging in size from 300 µm to up 1200 µm;
3. Kakowa (syn. Cacova) meteorite: poorly defined PO, RP and BO chondrules, ranging in size from 300 µm up to 600 µm;
4. Zsadany (syn. Jădani) meteorite: readily distinguished PO, POP, RP and BO (sometimes polysomatic) chondrules, ranging in size from 250 µm up to 750 µm;
5. Mocs (syn. Mociu) meteorite: readily or poorly defined PO, RP and BO chondrules, ranging in size from 250 µm up to 700 µm;
6. Tăuţi meteorite: poorly defined PO, RP and BO chondrules, ranging in size from 300 µm up to 600 µm;
7. Sopot meteorite: well defined or readily distinguished PO, POP, RP and BO (sometimes polysomatic) chondrules, ranging in size from 200 µm up to 600 µm.

References

- Gooding, J.L., Keil, K. (1981) Relative abundances of chondrule primary textural types in ordinary chondrites and their bearing on conditions of chondrule formation, *Meteoritics*, **16**, 17-43.
- Graham, A.L., Bevan, A.W.R., Hutchison R. (1985) Catalogue of meteorites; with special reference to those represented in the collection of the British Museum (Natural History), 4th ed., *British Museum (Natural History)*, London, 460.
- Iancu, O.G. (2003) Stages of shock metamorphism reached by the meteorites fallen in Romania. In Abstr. vol. „Petrology-Global Context”, Anniversary Symposium dedicated to Acad. Dan Rădulescu on his 75th Anniversary, p. 30, Bucharest.
- Miura Y., Iancu, O.G., Iancu, G., Yanai, K., Haramura, H. (1995) Reexamination of Mocs and Tauti chondritic meteorites; classification with shock degree, *Proc. NIPR Symp. Antarct. Meteorites*, **8**, Tokyo: 153-166.
- Norton, O.R. (2002) The Cambridge encyclopaedia of meteorites, *Cambridge Univ. Press*, 354.
- Stanciu, V., Stoicovici, E. (1943) Romanian meteorites (in Romanian). *Rev. Muz. Mineral. Geol. Univ. Cluj la Timişoara*, **VII** (1-2), 3-34.
- Van Schmus, W.R., Wood, J.A. (1967) A chemical-petrologic classification for the chondritic meteorites, *Geochimica Cosmochimica Acta*, **31**, 747-765.

THE ROLE OF FLUIDS DURING FORMATION OF THE SVECOFENNIAN MAGMATIC ROCKS (RUSSIA)

Elena Kouzmina³⁰

The Tervu massif is one of the Late Svecofennian magmatic massifs situated near the Archean–Proterozoic boundary within the Fennoscandian shield (The North-West Ladoga Region, Russia). Features of the fluid regime during the crystallization of the two feldspars granites of the Tervu massif's have been studied. Based on thermo- and cryometric methods, five types of fluid inclusions were identified in quartz crystals, as follows:

- 1) inclusions of H₂O (mono, two-phase);
- 2) low mineralized-aqueous mono- and two-phase inclusions (up to 1-3 wt. % of NaCl, KCl);
- 3) highly mineralized-aqueous (Na⁺, K⁺, Ca²⁺, Mg²⁺ salts, i.g. chlorides) inclusions (two- or poly-phase);
- 4) high density CO₂-inclusions (mono-, two-phase);
- 5) CH₄-N₂ inclusions (mono-phase).

The frequency of the fluid inclusions shows an apparent heterogeneity of the different parts of the massif. Also, the concentration and composition of the fluid inclusions in different parts of the massif are variable. The wide dispersion of the distribution of the H₂O and CO₂ inclusions is typical. All the data obtained on the fluid inclusions of the Tervu massif indicate that the equilibrium of the earliest inclusions changed later, during late-metamorphic events, and, probably, new supplies of metamorphic fluids were introduced in the granitic system.

³⁰ Saint Petersburg State University, Institute of Precambrian Geology and Geochronology. St. Petersburg, Russia. E-mail: <ek@ek6762.spb.edu>.

THE PRESENCE OF TOURMALINE IN THE HARGHITA MOUNTAINS' VOLCANIC STRUCTURES (ROMANIA)

Attila-Albert Laczkó³¹

Based on currently knowledges, the tourmalinization processes in the Harghita Mountain's Neogen-Quaternary volcanic arc have an important bearing in the major volcanic structures, such as Stâncă, Ostoros, Ivó - Cocoizaş, Vârgheiş and Luci-Lazu volcanic structures (Fig. 1).

Fig. 1. – Volcanological map of the Harghita Mountains (according to Szakács & Seghedi, 1995, with modifications). Legend: 1. Quaternary swamp or lake deposits; 2 - Tertiary postvolcanic and synvolcanic sediments; 3. Tertiary prevolcanic molasse sediments of the Transylvanian basin; 4. Cretaceous-Tertiary sediments of the Flysch zone of the Eastern Carpathians; 5. Late Paleozoic-Cretaceous sediments of the Eastern Carpathians; 6. Precambrian-Paleozoic metamorphic and plutonic rocks of the Crystalline-Mesozoic Zone of the Eastern Carpathians; 7. Neck; 8. Crater; 9. Caldera-like depressions; 10. Collapse calderas (caldera fault); 11. Porphyritic intrusive rocks; 12. Fine porphyritic intrusive rocks; 13. Volcanic core complexes; 14. Extrusive domes; 15. Lava flows; 16. Pyroclastic cone; 17. Stratovolcanic cone; 18. Effusive cone; 19. Coarse pyroclastic rocks – proximal facies; 20. Mudflow, debris avalanche, debris flow and ephemeral stream epiclastic volcanic rocks. Volcanic edifices and areas: 1. Răchitiş; 2. Stâncă-Ostoros; 3. Ivó-Cocoizaş; 4. Vârgheiş; 5. Harom-Jigodin-Şumuleu; 6. Luci-Lazu; 7. Cucu; 8. Tirco; 9. Pilişca; 10. Ciomadul; 11. Bicsad-Malnaş volcanic field.

Apart from the volcanic structures mentioned above, tourmaline is believed to occur in many other areas affected by the hydrothermal-metasomatic processes. That fact is based on the high boron concentrations in these areas, between 200-3000 ppm to 8000 ppm value (Tekerő

³¹ S.C. "Geolex" S.A. – Miercurea Ciuc, Romania. E-mail: <laczkoati@yahoo.com>.

crater on the north-eastern part of the Luci-Lazu volcanic structure; the area between the riverhead of the Köveş stream and the Cormoş stream; the area between the Mic stream and the Harom stream; the intracraterial zone of the Cucu volcano – Moţoi *et al.*, 1975; Rădulescu *et al.*, 1984).

In the relevant professional literature tourmaline is considered as being an indicator of the "porphyry copper" type (Cu - Mo +/- Au) mineralizations (Vlad, 1983). The presence of the tourmalinization in the Harghita Mountains' Neogen-Quaternary volcanic rocks does not indicate in all cases the existence of the "porphyry copper" mineralizations. Only in the Ostoros, Ivó - Cociozaş and possibly Vârghiş volcanic structures such mineralizations have been identified. The other mineralized zones in the Harghita Mountains differ from this type of mineralization.

The tourmalinization processes were located in the intensely tectonized zones (Stanciu, 1976; Tănăsescu, 1978; Boboş, 1995; Laczkó *et al.*, 2003). In all the cases the existence of tourmaline was supposed first based on the presence of a boron anomaly (hundreds and thousands ppm), and later this assumption was confirmed by mineralogical and X-ray diffractions analyses.

As a rule, tourmaline is present as rosettes, fans, granular aggregates, acicular crystals and microcrystals and in general is associated with quartz (Figs. 2, 3). As a function of the contents of metallic ions in the hydrothermal fluids, tourmalines are represented by schörlite (Fe-rich tourmaline – Stanciu, 1976; Tănăsescu, 1978), dravite (Mg-rich tourmaline - Laczkó *et al.*, 2003) and by elbaite (Li-rich tourmaline – Boboş, 1995).

Fig. 2. – Tourmaline in the Stâncă structure.

a) in the intrusive body (F1 Drill); b) in breccia's cement (F2 Drill). 1N.

The widespread presence of the tourmalinization within the framework of syn-magmatic processes in the Harghita Mountain's Neogen-Quaternary volcanic rocks, suggest that the boron was one of the main volatile substances which played an important role in the mineralization processes in these volcanic structures. The X-ray powder diffraction analyses data³² from the Ivó-Cociozaş and Luci-Lazu (Sântimbru-Băi) structures (Fig. 4) show the presence of dravite (Mg-rich tourmaline) in these areas (Laczkó *et al.*, 2003).

Fig. 3. Tourmaline from Sântimbru-Băi. a) brecciated andesite with quartz-dravite cement – N +; b) breccia matrix consisting of quartz (colourless), dravite (acicular crystals) and pyrite (black)– 1 N.

³² X-Ray Powder Diffractometer type DRON-3; Cu anticatode, radiation K_α with $\lambda = 1,54051 \text{ \AA}$.

Fig. 4. – The X-ray powder diffraction spectra of tourmaline samples from a) the Sântimbru-Băi quarry, ground level 0 and b) the Sântimbru-Băi quarry, ground level 1. D – dravite, Q – quartz.

References

- Boboş, I. (1995) Kaolinite deposits from the Harghita Mts. (in Romanian). *Ph.D. Thesis*, "Babeş-Bolyai" University of Cluj-Napoca, 204 p.
- Laczko, A., Ghergari, L., Tóth, A. (2003) Pollution sources in the area of Hg-ore deposit from Sântimbru Băi, Harghita county. *Environment & Progres Sympos. Vol.* (Petrescu, I. – ed.), 2003, "Babeş-Bolyai" University, Cluj-Napoca, 303-307.
- Moţoi, Gr., Szakács, Al., Setel, A., Vărăşmaş, N., Setel, M., Tănăsescu, L. (1975) Report. *Unpubl.*, 62 p. S.C. "GEOLEX" S.A. Archives, Miercurea Ciuc.
- Rădulescu, D., Peltz, S., Stanciu, C., Seghedi, I., Szakács, Al., Udrescu, C., Bratosin, I., Tănăsescu, A., Vâjdea, E., Grabari, G., Stoian, M., Popescu, Fl., Ionescu, Fl., Popescu, L., Niculin, M., Scurtu, Fl., Moldoveanu, M. (1984) Report. *Unpubl.*, 102 p. S.C. "GEOLEX" S.A. Archives, Miercurea Ciuc.
- Stanciu C. (1976) Hydrothermal alterations in the Ostoros volcano (Drill 3) from the Harghita Mts.(in Romanian). *D.S. Inst. Geol. LXII/1*, 199-213, Bucureşti.
- Szakács, Al., Seghedi, I. (1995) Time-space evolution of Neogen-Quaternary volcanism in the Călimani-Gurghiu-Harghita volcanic chain. *Rom. Jour. Stratigraphy. 76*, 24 p. Bucureşti.
- Tănăsescu, L. (1978) Data upon the presence of tourmaline and fluorite in the Neogene volcanics from the Harghita Mts (in Romanian). *D.S. Inst. Geol. LXIV* (1976-1977), 37-41, Bucureşti.
- Vlad, Ş.N. (1983) The geology of the porphyry copper deposits (in Romanian). *Ed. Acad. Rom.* 156 p., Bucureşti.

PHOSPHATES IN THE BAT GUANO DEPOSIT FROM GRIGORE DECAPOLITUL CAVE (CĂPĂȚÂNII MOUNTAINS, SOUTH CARPATHIANS, ROMANIA)

Ștefan Marincea³³, Delia-Georgeta Dumitraș¹, Gabriel Diaconu²

“Peștera Liliecilor de la Mănăstirea Bistrița” (the Cave of the Bats from Bistrița Monastery) is located in the right slope of the gorges of the homonymous river (Căpățânii Mountains, South Carpathians), at about 800 m upstream from the monastery. The cave, referred to hereafter as “Grigore Decapolitul Cave”, is famous due to a small church located in the biggest room of the cave, opened at about 80 m upwards the valley bed. This church hosts the relics of the Saint Grigore Decapolitul, who gave the second name of the cave. The cave galleries are developed in Upper Jurassic micritic limestones with calcarenite intralayers. Including the divergent galleries, the cave has 400 m in length.

A small (25 m x 15 m x 5 m) room, located in the terminal part of the cave, hosts an active bat colony. Consequently and thus an important bat guano deposit can be observed on the floor. Hydroxylapatite, brushite, taranakite, ardealite, leucophosphite, gypsum and calcite mainly compose the guano mass. At the basis of the guano deposit, near its limit with the *terra rossa* mass, small amounts of illite (the 2M₁ polytype), interstratified kaolinite-illite, quartz and amorphous iron sesquioxides are included in the guano. The iron sesquioxides are probably the same that gave the red ferruginous staining to the *terra rossa*.

The purpose of the present study is to obtain additional data on the phosphate and sulfate species in the deposit, using inductively coupled plasma - atomic emission spectrometry (ICP-AES), ion chromatography, scanning electron microscopy, X-ray powder diffraction (XRD) and infrared absorption spectrometry. All analytical facilities, procedures and experimental details are similar to those already described by Marincea *et al.* (2002) and Marincea & Dumitraș (2003).

Hydroxylapatite occurs as cream coloured to orange crusts of dull appearance. SEM images show that these crusts are mainly composed by spherical aggregates of up to 100 μm in size. In their turn, the aggregates are composed by thin, roughly hexagonal blades of hydroxylapatite, flattened on (0001). Individual crystals are up to 0.2 μm thick and 10 μm across. The crystal masses are highly fractured, and weathering products (probably brushite, but also some gel-like, iron-bearing phases) may be observed within the fractures.

The indices of refraction, measured in monochromatic light ($\lambda = 589$ nm), on thin, platy aggregates, taken off from a representative sample (PGD 1 B) are $\omega = 1.646(1)$ and $\varepsilon = 1.640(1)$. The density of the same sample, measured by sink-float in methylene iodide diluted with toluene, at 25°C, is 3.15(1) g/cm³, which compares well with the calculated value $D_x = 3.155$ g/cm³.

An ICP-AES analysis of a representative sample (PGD 1 B - with sulfate measured by ion chromatography and water in hydroxyl groups deduced for charge balance) gave (in wt.%): CaO = 54.75, MnO = 0.14, MgO = 0.29, FeO = 0.45, K₂O = 0.10, Na₂O = 0.12, P₂O₅ = 41.54, SO₃ = 0.92, H₂O = 1.69. The corresponding chemical-structural formula, calculated on the basis of 6 (P + S) and 26 (O,OH) per formula unit (*pfu*), is:

The (Sr, Ba, REE)-for-Ca substitutions are minor: the analyzed sample contains low Sr (102 ppm), Ba (70.9 ppm), Yb (0.899 ppm), Gd (2.04 ppm), Er (0.838 ppm), Dy (2.57 ppm), Y (16 ppm), Eu (0.265 ppm), La (8.82 ppm), Ce (2.37 ppm) and Nd (6.5 ppm).

The infrared spectrum of the same sample gave, however, a pattern typical for a water-bearing carbonate-hydroxylapatite. The spectrum contains OH stretching (3412 cm⁻¹) and librational (625 cm⁻¹) bands of molecular water, together with bands assumable to CO₃ groups (ν_3 1466 cm⁻¹, ν_3' ~1450 cm⁻¹, ν_2 873 cm⁻¹). Bands materializing vibrational modes of the PO₄ groups were found at 1080 cm⁻¹ (ν_3), 1042 cm⁻¹ (ν_3'), 960 cm⁻¹ (ν_1), 604 cm⁻¹ (ν_4), 564 cm⁻¹ (ν_4') and 470 cm⁻¹ (ν_2).

³³ Geological Institute of Romania, Bucharest, RO-73844, Romania, E-mail: <marincea@igr.ro>.

The unit cell parameters for 10 representative samples, as refined by least squares from XRD data, are given in Table 1.

Table 1.
Unit cell parameters of hydroxylapatite (Grigore Decapolitul Cave).

Sample	a (Å)	c (Å)	V (Å ³)	n ⁽¹⁾	N ⁽²⁾
PGD 1 A	9.425(2)	6.878(2)	529.2(2)	6	32
PGD 1 B	9.431(3)	6.867(3)	529.0(4)	3	30
PGD 5 B	9.433(2)	6.879(3)	530.1(2)	4	31
PGD 5 C	9.419(3)	6.895(3)	529.8(3)	4	40
PGD 6 A	9.422(3)	6.881(4)	529.1(4)	3	53
PGD 6 C	9.420(2)	6.873(1)	528.2(2)	10	32
PGD 8 B	9.431(3)	6.872(5)	529.3(4)	4	29
PGD 9 B	9.423(2)	6.876(3)	528.8(2)	4	46
PGD 9 C	9.431(2)	6.874(3)	529.5(2)	6	40
PGD 11 B	9.430(2)	6.877(3)	529.6(3)	4	40

* Monochromatized Cu K α radiation (λ = 1.54056 Å); (1) number of least squares refining cycles; (2) number of reflections in the 2 θ range 10 - 90° used for refinement.

The mean unit cell parameters, taken as average of the ten sets of values in Table 1, are $a = 9.427(5)$ Å $c = 6.877(7)$ Å and $V = 529.3(5)$ Å³.

Brushite generally occurs as bright white coatings, with silky luster, on the hydroxylapatite crusts, or as thin layers, nodules or lenses enclosed by the guano mass. SEM study shows that the mineral forms compact aggregates of finely lamellar crystals, flattened on (010) and elongated toward a direction that may be [101] or [102]; they are up to 0.1 μ m thick, 5 μ m wide and 10 μ m long. Radial sprays of very thin plates, disposed on hydroxylapatite, were also found.

An ICP-AES – ion chromatography analysis of a representative sample (PGD 2 C) yielded (in wt.%): CaO = 32.36, MnO = 0.03, MgO = 0.05, FeO = 0.09, K₂O = 0.07, Na₂O = 0.01, P₂O₅ = 40.89, SO₃ = 0.38, H₂O (as calculated for charge balance) = 26.12.

This composition, normalized on the basis of 1 (P + S) *pfu*, leads to the chemical-structural formula: [Ca_{0.993}Mn_{0.001}Mg_{0.002}Fe²⁺_{0.002}K_{0.003}Na_{0.001}](HPO₄)_{0.992}(SO₄)_{0.008}·2H₂O.

As well as in the case of hydroxylapatite, the (Sr, Ba, REE) for Ca substitutions are insignificant; the analyzed sample contains minor Sr (45.8 ppm), Ba (70.1 ppm), Y (2.03 ppm) and La (2.13 ppm).

The refined unit cell parameters of the sample, based on 60 reflections between 7.5 and 1.1 Å, for which an unambiguous indexing was possible, are $a = 5.802(3)$ Å, $b = 15.181(8)$ Å, $c = 6.235(3)$ Å and $\beta = 116.15(3)^\circ$. With $Z = 4$, the calculated density for the formula given before is $D_x = 2.318$, which is close to the value measured by sink-float in methylene iodide diluted with toluene [$D = 2.32(1)$]. The indices of refraction of the same sample, measured as maximum and minimum values on platy clusters of crystals are $n_{\min} = \alpha = 1.540(2)$ and $n_{\max} = \beta = 1.546(3)$, γ (calculated for $2V_c = 86^\circ$) = 1.551. Calculation of the Gladstone-Dale relationship using the constants of Mandarino (1981) yields superior compatibility (compatibility index = - 0.012).

Ardealite occurs as earthy, white-yellow aggregates, deposited on hydroxylapatite crusts or as nests and small veins in the brushite - hydroxylapatite masses. The unit cell parameters, calculated by least squares refinement from 63 reflections taken off from the most representative diffraction pattern (sample PGD 8 A), are: $a = 5.714(3)$ Å, $b = 31.002(13)$ Å, $c = 6.255(3)$ Å and $\beta = 117.16(3)^\circ$.

Leucophosphite sparsely occurs as very fine-grained (up to 20 μ m in length and 10 μ m in width) reddish-brown, platy crystals, typically associated with 2 M₁ illite and low (alpha) quartz. The mineral was found at the basis of the guano mass, near the contact with *terra rossa*. The unit cell parameters calculated after 5 cycles of least-squares refinement from the X-ray powder data (46 reflections in the d range of 4.75 to 1.11 Å) are $a = 9.762(4)$ Å, $b = 9.667(3)$ Å, $c = 9.748(4)$ Å and $\beta = 102.28(2)^\circ$. Wet-chemical tests on impure material helped to discriminate between leucophosphite and spheniscidite, which have basically the same XRD pattern.

Taranakite occurs as crusts or nodular masses composed by ill formed, almost warty, crystal groups, and, more frequently, as earthy masses of chalky appearance. SEM examination shows that the mineral forms clusters of roughly hexagonal plates, up to 20 µm across.

The cell parameters of 7 selected samples, as refined by least-squares analysis of the X-ray powder data, are given in Table 2.

Table 2.

Unit cell parameters of taranakite (Grigore Decapolitul Cave).

Sample	<i>a</i> (Å)	<i>c</i> (Å)	<i>V</i> (Å ³)	n ⁽¹⁾	N ⁽²⁾
PGD 6 A	8.701(1)	95.04(2)	6231.0(13)	8	54
PGD 6 B	8.694(1)	94.98(2)	6217.6(17)	5	76
PGD 7 A	8.692(1)	94.92(1)	6210.7(8)	8	63
PGD 9 C	8.694(1)	94.96(1)	6215.1(11)	10	77
PGD 10 A	8.698(7)	94.93(1)	6219.9(9)	10	70
PGD 10 C	8.710(2)	95.04(4)	6244.7(27)	8	58
PGD 11 A	8.697(1)	95.08(2)	6228.0(13)	6	85

* Monochromatized Cu Kα radiation (λ = 1.54056 Å); (1) number of least squares refining cycles; (2) number of reflections in the 2θ range 5 - 65° used for refinement.

The mean unit cell parameters, taken as average of the values in Table 2, are *a* = 8,698(6) Å, *b* = 94,99(6) Å and *V* = 6224(11) Å³. Both values are lower than those calculated by Marinacea & Dumitraş (2003) for synthetic (NH₄)₃Al₅(HPO₄)₆(PO₄)₂·18H₂O [*a* = 8.726(6) Å and *c* = 96.17(9) Å] and K₃Al₅(HPO₄)₆(PO₄)₂·18H₂O [*a* = 8.706(4) Å and *c* = 95.94(8) Å], which is probably due to both Fe³⁺-for-Al and (Na, Ca, Mn, Mg, Fe²⁺)-for-K substitutions.

Gypsum forms radial or parallel aggregates of minute bladed crystals up to 15 µm in length, which currently show parallel growth along the *b** axis. Macroscopically, the mineral occurs as snow-white aggregates deposited at the surface of the guano mass or as small (up to 5 cm across) nests or lenses enclosed by guano. The cell parameters of a representative sample (PGD 4 A), calculated by least-squares refinement on the basis of 69 X-ray powder reflections, are *a* = 5.640(3) Å, *b* = 15.243(6) Å, *c* = 6.501(3) Å, β = 118.18(2)°.

References

- Mandarino, J.A. (1981) The Gladstone-Dale relationship. IV. The compatibility concept and its application. *Can. Mineral.*, **19**, 441-450.
- Marinacea, Ş., Dumitraş, D. (2003) The occurrence of taranakite in the "dry" Cioclovina Cave (Sureanu Mountains, Romania). *N. Jb. Mineral., Mh.*, **3**, 127-144.
- Marinacea, Ş., Dumitraş, D., Gibert, R. (2002) Tinsleyite in the "dry" Cioclovina Cave (Sureanu Mountains, Romania): the second world occurrence. *Eur. Jour. Miner.*, **14**, 157-164.

SALT DAMAGES OF SOME SANDSTONE BUILDINGS IN BELGRADE (SERBIA)

Vesna Matovic³⁴, Nada Vaskovic¹, Aleksandra Rosic¹

Introduction

Sandstone has been a traditional building material in Serbia since medieval times. It was used extensively in building monuments, sculptures and especially in Serbian parochial architecture.

In the first half of the twentieth century, and later (1955-1970), sandstone was used as a building material as irregular or regular slabs or blocks as well as for stone panels on façades of numerous buildings of Belgrade. Sandstone buildings are exposed to both external and internal weathering processes. The interaction of these processes depends on the nature of the environment and the degree to which both, the features of the building and the location of the stone within the building, affect this. Through the normal processes of weathering/thawing, a natural breakdown of sandstone occurs. This is intensified by salt crystallization, and the internal processes involved in the mineralogical conversion.

Unfortunately, in the last twenty years, the acceleration of sandstone decay in buildings of Belgrade has been noticed. The common damages types are as follows: exfoliation, granular disintegration, peeling, black crust formation and efflorescence.

Soluble salts are important agents in the stone deterioration. They concentrate as efflorescence on the surface and as sub-efflorescence close beyond the surface of the stone. In both cases, soluble salts occur in most places where deterioration occurs. They are very mobile, moving along the surface as well as in and out of the surface. These movements occur periodically i.e. depending on the seasons and the atmospheric conditions (Amoroso & Fassina, 1983). The sources of the soluble salts are various, including the geogenic sources (the stone itself, rain water, groundwater etc) and sources related to the anthropogenic activities and products (atmospheric pollution, mortars etc; Winkler, 1994).

The crystallization of soluble salts is a widespread weathering process on more observed sandstone buildings in Belgrade. The aim of this paper is to represent morphology of salt weathering and its mineral composition on the following buildings: St. Marco Church, Riverbank Pillars of Brankos Bridge and the Monument "Oslobodjocima Beograda 1944".

The mineralogical composition of powdered salt samples was obtained with a "Philips" PW1051 diffractometer under the following experimental conditions: $V_g=1^\circ 20/\text{min}$, $V_p=10\text{mm}/\text{min}$ and $R/C=500/\text{l}$ using $\text{CuK}\alpha$ radiation, tube conditions 45 kV and 25 mA.

Physical properties of stone samples were measured in the Laboratory of stone and aggregates at the Institute of highway in Belgrade.

History and architecture of some objects in Belgrade

The *Branko's Bridge* is an important utilitarian structure connecting the old part of the Belgrade on the right bank of the river Sava with the new part of the city, situated on its left bank. The old decorated pillars represent an extraordinary union of modern engineering and old architecture. Some of them are still used for communication, as stairways. Pillars built from 1931 to 1933 are made of ferro-concrete, while stone paneling was made of Bele Vode sandstone. They were constructed in Roman-Byzantine style adorned by numerous architectural elements and sculptural decoration (Kadijevic, 1996). Cornices, rosettes and consoles ornament the upper parts of the façades.

The *St. Marco Church* was built between 1931 and 1939, and represents the first building of the Byzantine style in our country made of ferro-concrete. Roofs are also made of ferro-concrete and covered with cooper, walls are of cement-mortar. The church façades were made of the sandstone from Bele Vode and of Red Permian Sandstone. The plinth with outflows out of the walls is faced with granites.

³⁴ Faculty of Mining and Geology, Djusina 7, 11000 Belgrade, Yugoslavia. E-mail: <vesnamat@mkpg.rgf.bg.ac.yu>, <nadavask@mkpg.rgf.bg.ac.yu>.

The monument “*Oslobociocima Beograda 1944*” was constructed in 1954. The Main Portal consists of five entrances with short sidewise walls with bas-reliefs. The main portal, the platforms and the steps were made of sandstone blocks from the quarry Osoje near Ljig, and the bas-reliefs of marble from Brach Island (Croatia). Sandstone tiles flag platforms and steps. The portal is composed of 459 visible sandstone blocks of variable size. Randomly location of the different blocks and bushhammering has caused the occurrence of various type of destruction.

Petrography and physical properties of sandstones

Bele Vode sandstone (BV) represents Lower Miocene, shallow-water lacustrine massive to well bedded grayish to yellow-brownish, sometimes pinkish or red colored sediments. They have often cavities, of approximately 0.5 mm in diameter. The coarse- to fine-grained (0.2-1mm) sandstones used as building stone may be described as arkoses, composed of poorly-sorted angular (the gray varieties) or subrounded to rounded (the yellow varieties) grains of quartz (50-60 vol. %), K-feldspar (up to 15 vol.%), plagioclase (up to 5 vol.%), white mica (up to 10 vol.%) and rock fragments (up to 10 vol.%; mainly metamorphic). In the yellow and pinkish to red varieties the contact-pore binding is ferro-calcite, while in the grayish variety, it is quartzitic-argillaceous.

The physical properties of the BV are as follows: bulk density (1880-2420 kg/m³), absolute porosity (7.4-27.9 %) and water absorption (2.43-9.16 %), indicating that the BV are medium heavy and high porous rocks with moderate to high water absorption.

Long-lasting studies under frost, freezing and thawing cycles revealed the BV sandstone as stable rocks, whereas crystallization experiment estimated it as partly unstable. The sandstone durability is influenced by variable porosity and water absorption ability of pore cavities.

The Upper Cretaceous sandstones from the Ljig flysh - quarry Osoje (LO) are subarkose to lithic arenite, brown to gray in color, fine - or fine - to coarse-grained sediments uniform or slightly sorted sometimes with lamination and convolution as well as fissures and cracks parallel to bedding. They are composed of rounded to angular grains of quartz, (55.82 vol.%), feldspar (15.33 vol.%), rock fragments (17.42 vol.%) and white mica (5.02 vol.%). The contact and pore binding is calcitic to argillaceous-quartzitic and slightly ferruginous.

The physical properties of the LO are as follows: medium heavy (2460 kg/m³), medium to high porous (9.85 %) with slight to moderate water absorption (2.25 %).

Red Permian sandstone from the Grza (RPG) is well bedded to massive, with rare lamination and cross bedding. Some parts of the red pile are colorless or grayish colored. It is mainly arkose, rarely quartzitic, and it is composed of subrounded to angular poorly-sorted grains of quartz (50-70 vol. %), feldspar (up to 25 vol.%), mica (up to 2 vol.%) and rock fragments (up to 5 vol.%) – volcanic, chert, quartzite, schist. Contact and pore binding are ferruginous and calcitic or quartzitic.

Physical properties point to the homogeneous quality of the RPG: density (2630-2692 kg/m³), the extreme to moderate porosity (4.9-11.2 %) and the slight to moderate water absorption capacity (1.53 to 3.06 %).

Environmental conditions

Generally, all the buildings in Belgrade are exposed to a medium continental climate with wet and hot summers and long cold winters, often with fogs and snow, and short autumns and springs. The average yearly precipitation, the number of the frost-icy days and the value of the high relative humidity during winter, accelerate the degradation of the sandstones. Simultaneous precipitations (yearly average of 684.5 mm), and temperatures below 0° C (61.5 frost days per year and 17.6 ice days per year), cause freeze/thaw action within materials. Yearly and monthly thermo-oscillations are considerable. Predominant winds are from NE (max. 50-100 km/h). During the summer time, the highest temperatures are accompanied by high insulation. The geographical position of the town, the relief, the climate and the rapid development of industry after 1950 caused high concentrations of air pollution. Yearly average concentration of smoke and SO₂ are relatively low but for individual months (as November, February) daily maximum values in Belgrade are over the recommended limits (50 µm/m³ for smoke and 150 µm/m³ for SO₂). The differences in SO₂ concentration between summer- and wintertime indicate a great input of the pollution, related to the heating systems. The combustion of fossil fuels releases sulfur as SO₂ into the atmosphere. The yearly average pollutant concentrations are relatively low, but the maximum yearly average for

individual years (1991, 1992) was appreciably over the accepted limit in Belgrade. The main effect of SO_2 in the air is the changing of the pH of the rain. These acid rains attack building façades. During the last ten years, the content of SO_2 and smoke show decreasing trend but even these lower contents are damaging for stone.

Salt damages

Several types of degradation are noticed on the studied buildings. The granular disintegration, exfoliation, scaling, blistering and the black crust are the dominant stone pathologies. Formation of thin films and patches of soluble salts follow these damages. The salt efflorescences are found on the surfaces of sandstone elements in the different positions, at the exterior and interior (stone elements that are sheltered from the rain - arcades, porch etc) parts of the façades. It appears mostly at the border between the wet and the dry zones of walls, mostly in the form of white powder-like accumulations.

At the riverbank pillars of the Branko's bridge, the visual evidence of efflorescences are found in arcade's porches where they occur at the upper limit of the ground moisture rising into the wall (Fig. 1). X-ray data show that they are composed of thenardite (Na_2SO_4) and thermonatrite ($\text{Na}_2\text{CO}_3 \cdot \text{H}_2\text{O}$). Water from the soil had leached sodium sulfate/carbonate from the soil and capillary action drew this salt solution into the stone. During the long-lasting moisture, these salts penetrate into the stone surface, collect soot and begin to recrystallize. Losing its mechanical strength, the stone surface disintegrates into powder.

At the less sheltered surfaces of Branko' bridge, the efflorescences occur as white, soft patches on the stone surface (Fig. 2). They are composed of gypsum (CaSO_4) and halite (NaCl), their sources may be different. Gypsum is relatively enriched near the mortar joints and probably it is "leached" from the mortar matrix. Halite can be a result of rainwater evaporation or deicing, but in any case all of these salts accelerate peeling, chipping and later exfoliation of the sandstone blocks (Matovic, 1999).

At the St. Marko church, the observation of the stone panel made of BV sandstone shows minor signs of deterioration and no efflorescence is found on their surfaces (Matovic & Milovanovic, 2001). However, certain parts of the façades made of RPG sandstone are badly deteriorated. The granular disintegration and exfoliation are followed by the forming of the efflorescences and probably sub-efflorescences. The deposits of soluble salts occur on higher parts of façade, but only on the sheltered surface of the arcades porches. X-ray analysis of salt was carried out in samples from the arcade's porches. The data showed various amounts of hydration water in the analyzed phases (thenardite - Na_2SO_4 , mirabilite - $\text{Na}_2\text{SO}_4 \cdot 10\text{H}_2\text{O}$), which are influenced by the relative humidity in the atmosphere. These salts are concentrated from large surfaces into small areas but they produced considerable deteriorations. Their origin is related to the air pollution or to the cement mortar.

Fig. 1. Efflorescence in the upper zone of rising ground moisture (arcade porches of Branko's bridge).

Fig. 2. Efflorescence of gypsum and halite at less sheltered surface.

On the sandstone blocks of the monument “Oslobodiocima Beograda 1944” the following types of degradation were noticed: the exfoliation, the granular disintegration, the flaking and peeling, the forming of the efflorescence and of the black crust. The black crusts and the soluble salts are the results of a significant anthropogenic pollution impact on the sandstone blocks of this monument. The black crusts are composed of gypsum crystals, fly ashes and soot and include also some minerals originating from the sandstone (Fig. 3). The black crusts formed in sheltered areas not exposed to runoff, where there are conditions for the joint between salt solutions (input by capillary transfer) and the fly ashes and soot originating from the polluted air.

The X-ray analysis of sample scratched from the bas-reliefs of the Monument show the presence of thenardite and apthitalite – $(K,Na)_3Na(SO_4)_2$ (Fig. 4). Their source is probably related to the circulating water along mortar joints.

Fig 3. X-ray diffraction patterns of the Monument «Oslobodiocima Beograda 1944» (sample of black crust).

Fig. 4. X-ray diffraction patterns of salt from facades of the same monument.

Conclusions

At the studied sandstone buildings the most important damages are caused by water and frost as well as by the soluble salt migration. The presence of water and moisture favor the solubility and hydrolysis of the corrosion products; having evaporated, the water leaves behind concentrations of salt solutions, which crystallize on the stone surfaces. Phase changes of water-soluble salts are induced by environmental factors, namely variation of temperature and relative humidity. The crystallization of these salts requires water and results in their expansion causing the degradation of stone material. Consequently, the stone surfaces begin to exfoliate.

Analyses show that the main salts/corrosion products are gypsum, thenardite/mirabilite, apthitalite, thermonatrite and halite. The data allowed to group the possible sources of the salt as follows:

- Rain water, as an important source of salt (gypsum, halite, thenardite);
- Soluble parts of the building materials themselves (mortar);
- Ions transported with the rising damp from ground;
- Wet or dry deposition of air polluting substances and their interaction with stone.

References

- Amoroso, G., Fassina, V. (1983) Stone decay and conservation. *Elsevier S.P.*, 445 p.
- Kadijevic, A. (1996) History and architecture of the Zemun bridge King Alexander I Karadjordjevic (in Serbian). *Pinus*, 4, 4-10, Belgrade.
- Matovic, V. (1999) Decomposition of stone in the riverbank pillars of Branko's Bridge (in Serbian). *“Heritage”*, 2 (Eds. Institute for protection of Cultural Monuments of Belgrade), 107-114.
- Matovic, V., Milovanovic, D. (2001) Durability and Decay Type of Sandstone from the Façade of the St. Marko Church in Belgrade (Serbia). In *“Studies in Ancient Structures”*, Istanbul-Turkey, 599-609, CD-version.
- Winkler, E.M. (1994) Stone in architecture, Properties, Durability. *Springer-Verlag*, New York, 310 p.

PETROGRAPHIC NOTES ON THE PRAID SALT BRECCIA (ROMANIA)

Ioan Mârza¹, Vasile Pomârleanu²

Introduction

Salt breccia is a tectonic-diapiric formation that is often mentioned in Romanian literature due to its existence in local salt massifs. The mechanism of salt breccia formation has often been discussed, focussing on source of the clasts. Thus, detailed petrographic aspects that can provide useful information on the lithology and even on the stratigraphic geology of the formations situated immediately above the salt level, were overlooked. Romanian authors mentioned different origins of salt breccia:

- *Tectonic formation* (Popescu – Voiteşti, 1924, 1934, 1943);
- *Sedimentary formation* (Olteanu, 1951; Popescu, 1951);
- *Mixed-origin formation* (Filipescu, 1938).

The term *salt breccia* should only be used with a tectonic meaning, as it is genetically and spatially associated with diapiric structures formed due to the effect of salt tectonics (halotectonics). In other words, salt breccia is a external and cap-rock tectonic-diapiric intrusion body. The development of ideas about the salt diapirism was already presented by Pătruţ *et al.* (1973) and Drăgănescu (1997).

The Praid salt deposit was mentioned and studied by numerous authors such as Poşepny (1867), Popescu – Voiteşti (1943), Maxim (1961), Iorgulescu *et al.* (1962), Paucă (1967), Dragoş (1969) etc. According to Popescu – Voiteşti (1943), the Praid salt diapir pierces the formations of Mediterranean – Sarmatian to Pliocene ages, including volcanic tuffs, andesite agglomerates, etc. In the opinion of this author the age of the Praid salt is Aquitanian. The same author signals the presence of salt breccia in an outcrop in the Corund Valley.

The paper focuses on the study of the metamorphic clast samples collected by one of the authors (I. Mârza) from the inner part of the salt massif (underground). The data are compared with previously published data.

Poşepny (1867) mentioned the existence of a special type of breccia both at the surface of the salt massif as well as inside it. This breccia is formed by angular fragments of “trachytes and crystalline schists”. The author assumed that the crystalline schists originate from the disaggregated trachytic agglomerates that entrapped them.

The rock fragments that form salt breccia in the Praid diapir are located in the external parts as well in the apical part of the massif, inside and outside the salt diapir. The salt is intensely folded, faulted and brecciated during and possibly even after diapirism.

The clasts which form the salt breccia, are represented by various rocks, as sedimentary (marls, clays, sandstones), metamorphic (porphyroids) and magmatic (volcanics) ones.

In the breccia as well as inside the salt massif, elongated and ovoid-shaped lenses as well as almost circular pockets of marls and clay of few meters diameter are found.

Sandstones occur frequently, as angular fragments, sometimes fissured. The white (pure) fibrous salt (recrystallized) infilled the fissures due to tectonic processes.

Fragments of crystalline schists, from few millimeters to over 1 meter in size, occur frequently. The shape of the clasts is angular or slightly rounded, partly possible due to the movements inside the salt. These fragments occur along tectonized areas (microfolded and fractured).

The high frequency of hard rocks (porphyroids, sandstones) of angular aspect is surprising as it supposes their refragmentation in the tectonic-diapir process. Our study focuses on the petrographic description of the metamorphic clasts, called with the generic name of *crystalline schists*, by the previous authors.

1. The petrography of the metamorphic clasts

¹. Babes-Bolyai University, Department of Mineralogy, 1, M. Kogălniceanu Str., RO-3400 Cluj-Napoca, Romania

². Geological Institute of Romania, Caransebeş Str., RO-7000 Bucharest, Romania

The metamorphic clasts were collected in the mining workings (Level 266 m, chambers 6200, 6208) and then analyzed in thin sections. They correspond to porphyroides (metavolcanic rocks – metapyroclastic rocks) having various compositions, as follows: metadolerites, metadacites, metarhyolites, metatuffs. Often they are mylonitized.

Metadolerites are greenish rocks whose structure resembles that of chlorite schists with albite porphyroblasts. According to Mârza (1969) the plagioclase porphyroblasts formed during the tectonic-metamorphic phenomenon. The rock is formed of plagioclase relics caught in albite-quartz micrograiny groundmass, with chlorite and subordinate muscovite. The original structure of the premetamorphic rock of dolerite type is preserved (Pl. I, Fig. 1). Plagioclase relics partially maintain the polysynthetic twinning and are transformed into fine-grained muscovite (“sericite”) and zoisite. The rare magnetite grains represent premetamorphic relics.

The **metadacites** correspond to the most widespread lithic fragments found inside the Praid diapir. The rock is gray, greenish or black in colour. It has schistose structure and porphyric texture (relics of quartz and feldspars) - Pl. I, Fig. 2. The most frequent size of the rock fragments is 10 x 20 cm but there are also blocks ranging over 1 m in length. The disposal of these elongated fragments marks the direction of the tectonic flow of the salt.

Microscopically, the rock reveals the features of a typical porphyroid, with quartzo-feldspathic (0.08 x 0.10 mm – 0.02 x 0.03 mm size of the grains) and chlorite – muscovite (“sericite”) groundmass. Chlorite replaces pseudomorphically, hornblende. Small amounts of zoisite, actinolite, calcite, and iron hydroxides occur also. Apatite (0.03 x 0.04 mm) and zircon microlites (0.02 x 0.08 mm) represent premetamorphic components of the rock.

Plagioclase feldspar is represented by albite, sometimes of *schachbrietalbit* type. It has about 2mm in size and a sub-rounded shape. Some plagioclase crystals preserved the initial twinning but not the original content of An %, due to the metamorphic decalcification of the mineral. Quartz (0.48 x 0.50 mm) exhibits wavy extinction and jagged rims.

The semiquantitative analysis of the mineral components of the rock indicates the following percentages: 37% quartz, 26% feldspar, 12% muscovite, 10% chlorite, 3% opaque minerals (magnetite, pyrite), 11% calcite (secondary origin) and 1% other minerals as apatite, zircon and zoisite. The association of primary minerals corresponds to a dacite (as premetamorphic rock).

Metarhyolites have a gray colour. The structure is schistose, microfolded, and the texture blastoporphyratic. The groundmass is constituted of quartz and feldspar fine grains (0.08 x 0.10 mm) often associated with “sericite”, chlorite, rarely biotite. Orthoclase (Pl. I, Fig.3), plagioclase (polysynthetic twinned albite) and magmatic corroded quartz (relic feature; Pl. I, Fig. 4.) occur as phenocrysts.

The accessory minerals are represented by zircon, monazite, apatite, pyrite and hematite microlites. The presence of hematite and iron hydroxides is responsible for the reddish color of the rock. Rhyolitic volcanic rocks underwent hydrothermal silicification, mirrored by quartz veins of 0.30 – 0.50 mm width.

Mylonitized metarhyolites are characterized by a mylonitized quartzo-feldspathic groundmass (0.15 x 0.20 mm), with high amounts of muscovite (“sericite”) and chlorite.

Rhyolitic metatuffs are micrograiny rocks, having a schistose structure. They are formed mainly from quartz and rarely feldspar, “sericite” (muscovite), chlorite, apatite, zircon and pyrite.

2. The petrography of the Neogene volcanics clasts

Fragments of Neogene volcanics from the Praid salt breccia include andesites with pyroxene and hornblende.

Andesites with pyroxene and hornblende are dark-colored rocks, with a primary vacuolar microtexture. The fabric is subfluidal porphyric-hyalopilitic structure. The groundmass is mainly vitreous and partially microcrystalline (due to the feldspars microliths). The minerals are represented by polysynthetic-twinned plagioclase associated with pyroxene (augite; 0.50 x 0.80 mm size), completely oxidized hornblende and biotite (0.40 x 1.20 mm in size).

The semiquantitative analysis shows the following percentages: glassy-microlithic groundmass (52%), plagioclase (32%), pyroxene (10%), hornblende (3%), biotite (0.5%), opaque minerals (2.5%).

3. The petrography of the sedimentary rock fragments

The sedimentary rocks found in the Praid salt breccia are represented by Neogene sandstones and pelites (clays, marls). The clays and marls are fairly plastic and occur as “pockets”. They can also be laminated in highly-tectonized zones (on the sides of microfolds and in fractured zones).

Quartzo-feldspathic sandstones are the most widespread enclavated fragments inside the Praid diapiric massif. The sandstones have light gray colour, angular shapes and reach from few centimeters to over 1 meter diameter. Their mineralogical composition is as follows: quartz with wavy extinction, feldspar, mica (muscovite, partially oxidized biotite). Apatite, zircon, calcite and glauconite may also occur.

Clays and marls are found in sub-rounded, lens-like pockets and as laminae – in tectonically active zones which mirror the direction of the tectonic flow of the salt. Both clays and marls are basically constituted by clayey minerals (beidellite, montmorillonite), “sericite” (fine-grained muscovite), chlorite and quartz, feldspar, calcite, iron oxi-hydroxides etc.

Conclusions. The Praid salt breccia has a tectonic - diapiric genesis. According to its position in the salt massif, the breccias are located marginal, cap-rock and internal. The internal breccia discussed above is constituted from fragments of metamagmatic (porphyroides) rocks, andesites with pyroxene and amphibole (Neogene pyroclastics) and sedimentary rocks (clays, marls and sandstones). During the process of the salt migration, fragments of enclavated rocks floated in the salt due to the plastic environment. Thus they underwent sometimes syn-cinematic refragmentation that explain their angular shape.

The petrographic analysis presented in the paper shows a perfect resemblance between the fragments of metavolcanic rocks analyzed and the porphyroides of the Tulgheş type epimetamorphics. Fragments originated from epimetamorphic rocks and transported in the Sovata - Praid area of the sedimentation basin (Transylvanian basin) from the eastern border. These fragments of metamorphics, together with Neogene sedimentary rocks located above the salt level were involved in the diapir tectonics. They continuously were spatially redistributed and moved inside the salt deposit due to the mechanic effect of the salt.

Acknowledgments. The authors would like to thank geologist Otilia Sasu for the guidance and assistance she readily provided in the underground of the Praid mine.

References

- Drăgănescu, L. (1997) Origins and genesis of salt massifs (in Romanian) *S.C. Grafica Prahoveană S.A.*, 226 p., Ploieşti.
- Filipescu, M. (1938) Le calcaire de Bădila (Buzău) et quelques considérations sur l'enveloppe du sel. *C. R. Inst. Géol. Rom.*, **XXII** (1933 – 1934), 4–8, Bucureşti.
- Mârza, I. (1969) Evolution of metamorphic units from the south-eastern side of the Muntele Mare (in Romanian). *Edit. Acad. R. S. România*, 166 p., Bucureşti.
- Olteanu, Fl. (1951) Observations on salt breccia within salt massifs from the Mio-Pliocene area between Teleajen River and Bălăneasa stream (in Romanian). *D. S. I. G. R.*, **XXXII**, 12–18, Bucureşti.
- Pătruţ, I., Paraschiv, D., Dicea, O. (1973) On the formation of the diapiric structures in Romania (in Romanian). *Rev. Petrol şi Gaze, M.I.C. – M.M.P.G.*, **XXIV**/9, 533–542, Bucureşti.
- Popescu, Gr. (1951) Observations on the salt breccia and some salt massifs from Paleogene - Miocene area of the Prahova district (in Romanian). *D. S. Inst. Geol. Rom.*, **XXXII** (1943-1944), 3-12, Bucureşti.
- Popescu-Voiteşti, I. (1924) Galets a facettes dans la breche tectonique des massifs de sel de Roumanie. *Rev. Muz. Geol. – Min. Univ. Cluj*, **I**, 1, 96–100, Cluj.
- Popescu-Voiteşti, I. (1934) Geology of salt accumulations (in Romanian). *Rev. Muz. Geol. – Min., Univ. Cluj*, **V**, 1, 1-85, Cluj.
- Popescu-Voiteşti, I. (1943) The salt of the Romanian Carpathian regions (in Romanian). *Fund. Reg. Lit. Artă*, 74 p., Bucureşti.
- Pošepny, Fr. (1867) *II. Studien aus dem Salinengebiet Siebenbürgens. J. d. K. K. Geol. Reich.*, **XVII**, 475–516, Wien.

Fig. 1. Metadolerite clast. N+, 30x.

Fig. 2. Silicified metadacite clast. Quartz veins cross-cutting quartz macrograins having wavy extinction. N+, 45x.

Fig. 3. Metarhyolite clast. Orthoclase crystal in a quartzo-sericitic groundmass. N+, 45x.

Fig. 4. Metarhyolite clast. Magmatic corroded quartz, with wavy extinction. N+, 45x.

CHLORITE-CHLORITOID-GARNET EQUILIBRIA AND GEOTHERMOMETRY IN THE SANANDAJ-SIRJAN METAMORPHIC BELT, SOUTHERN IRAN

Mohssen Moazzen³⁵

Metapelites are important rocks, sensitive to the condition of metamorphism. Thus they can be used to estimate the pressure and temperature of the metamorphism. Despite these characteristics, it is difficult to find low-grade metapelites (greenschist facies) possessing enough phases in equilibrium to have low-variance assemblages. Therefore, the estimation of the pressure and the temperature of the metamorphism in these rocks is difficult. Some investigators have considered this problem and have attempted to calibrate useful geothermometers for low-grade pelitic rocks. The most important geothermometers applicable to low-grade metapelites are chlorite-garnet (for garnet phyllites and garnet schists) and chlorite-chloritoid Fe-Mg exchange geothermometers. Chlorite-chloritoid-garnet and chlorite-chloritoid assemblages were found in metapelites of the Sanandaj-Sirjan metamorphic belt in southern Iran. Other phases in the rocks include muscovite, quartz, magnetite, titanite and zircon. Chloritoid, chlorite, garnet and muscovite were analysed using electron microprobe and SEM. The chemical study of the minerals indicates that chlorites are ripidolite to daphnite. Chloritoids are Fe-rich and garnets are essentially almandine with appreciable amounts of grossular and spessartin. The analysed white micas are muscovites with considerable amounts of paragonite end-member, a celadonite end-member being absent. One calibration of chlorite-chloritoid Fe-Mg exchange geothermometer and three different calibrations of chlorite-garnet Fe-Mg exchange geothermometer were applied to metapelites of the Sanandaj-Sirjan metamorphic rocks. Chlorite-chloritoid thermometry yields temperatures of cca 515-557°C for the rocks. Chlorite-garnet thermometry gives temperatures between 442 and 460°C. The application of these thermometers yields fairly close temperatures for regional metamorphism in the studied area within the chlorite zone, indicating a medium to high greenschist facies. The difference between the results from garnet-chlorite and chlorite-chloritoid thermometers is about 70°C. Most probably, the temperatures obtained using the garnet-chlorite thermometer are more realistic. If this is true, then chloritoid-chlorite thermometer overestimates temperature about 70°C.

³⁵ Department of Geology, Tabriz University, 51664 Tabriz, Iran

GEOCHEMICAL FEATURES OF BERYL FROM VOISLOVA PEGMATITES (SOUTH CARPATHIANS)

Titus Murariu³⁶, Smarada Rădăşanu³⁷, Uwe H. Kasper³⁸, Thorbjorn Schoenbeck³

Introduction

Beryl is one of the most abundant and widespread minerals of beryllium in granite pegmatites.

Ideally, beryl formula can be written as $^{T(2)}\text{Be}_3\text{OAl}_2[^{T(1)}\text{Si}_6\text{O}_{18}]$, but the chemical composition and structure of most beryl samples are, however, more complicated. Whereas Si in $T(1)$ is virtually untouchable, extensive substitutions affect the Be – and Al – polyhedra and vacancies in channels (Černý, 2002). Therefore, from a crystal-chemical reason, beryl formula is briefly represented as:

In granite pegmatites from Carpathian Province of Romania (Mârza, 1980), beryl is found in some pegmatite bodies located in the Getic Subprovince (Teregova field – Semenice Mountains; Voislova and Tâlva field – Poiana Ruscă Mountains; Despina, Pietrele albe, Haneş, Streaia – Lotru Mountains) (Schadler, 1930; Superceanu, 1957; Maieru *et al.*, 1968; Pomârleanu, 1969; Hann, 1987; Murariu, 2001) and Gilău-Muntele Mare Subprovince (Bondureasa Valley – Mârza *et al.*, 1988).

Commonly, beryl occurs as large crystals in the intermediate zones of zoned pegmatite bodies, but was also found as small patches on fractures in albite. In pegmatites from Teregova, beryl (11 kg) is associated with quartz and K-feldspar, with plagioclase, quartz and muscovite; or with columbite, tantalite and montebrasite (Superceanu, 1957). In pegmatites from Bondureasa Valley beryl (7x3 cm) comprises inclusions of tourmaline, quartz and muscovite (Mârza *et al.*, 1988).

This paper reports on the geochemical features of beryl from Voislova pegmatites (Getic Subprovince). Samples of beryl occur as large green crystals. In thin section, the interior of beryl is filled by albite, (Fe,Mn)-phosphates and apatite. So far, only fluid inclusions of Voislova pegmatite beryl were studied by Pomârleanu (1969), but there are no further geochemical data available for beryl.

Analytical methods

The analytical methods used for major compounds of beryl from Voislova pegmatite were XRF and EMP.

XRF analyses were performed with a Philips PW2400 X-ray spectrometer, using the analytical procedure "oxiquant". 72 natural rocks and clays were used to determine the calibration curves of the pertinent elements.

EMP were carried out with a JEOL JXA-8900 instrument using operating current of 20 nA and accelerating voltage of 20 kV. X-ray intensities of the alkalis, the minor (Ti and Mn) and the major elements were counted for 5 s, 40 s, and 60 s, respectively. In order to minimize losses of Na and K the beam diameter was expanded to 10 mm. Components were standardized using natural minerals, glasses of natural rocks and synthetic oxide compounds. The results were corrected using the ZAF procedure (Reed, 1996).

The trace elements are analysed by ICP-MS method (Perkin Elmer/Sciex ELAN 6000 ICP-MS, quadrupole mass spectrometer). Measurements of element concentrations were performed using as internal standards Ru-Re (10ng/ml) to minimize drift effects and two calibration solutions (high purity

³⁶ "Al.I.Cuza" University, Department of Mineralogy and Geochemistry, 20 A, Carol I Blv., RO-6600 Iaşi, Romania.
E-Mail: <tmurariu@geo.uaic.ro>.

³⁷ "Al.I.Cuza" University, Department of Mineralogy and Geochemistry, 20 A, Carol I Blv., RO-6600 Iaşi, Romania.
E-Mail: <smaranda@geo.uaic.ro>.

³⁸ Geologisches Institut, Universität zu Köln, 49a, Zùlpicher Str., D-50674 Köln, Germany.
E-mail: <hu.kasper@uni-koeln.de>.

chemical reagents). A batch of 5 - 7 samples was bracketed by two calibration procedures. Accuracy and precision of determinations were checked with certified reference materials (CRM) (Govindaraju, 1994; Dulski, 2000).

All the analyses were performed at the Geological Institute of Köln University, Germany.

Geochemistry of beryl

Since BeO is one of the major oxides of beryl which cannot be determined by XRF and/or EMP, we recalculated the BeO concentration by stoichiometry, on the base of 3 Be cations per formula unit. The final results of beryl composition are reported in Table 1.

After BeO recalculation, a deficit in major oxides total still persists for all analyses, and we presume that it is caused by H₂O contents, which were not determined and therefore cannot be assessed by stoichiometry. Generally, the H₂O content in beryl pegmatites range between about 1.4 and 2.5 % (Deer *et al.*, 1999; Trueman & Černý, 1982). Rare alkalis (Li, Rb, Cs) do not exceed 0.12 per cent so that they do not affect significantly the sum of oxides.

The structural formula of beryl shows that this mineral accommodates alkali cations in channel and femic cations in octahedral positions (Table 1). The total content of femic cations is minor, and among them Fe²⁺ has the higher level of concentration. In five of the samples Mn prevails over Mg. Just one sample is enriched in Mg relative to Mn.

Table 1.
The composition of beryl from Voislova pegmatites.

Oxides %	EMP					XRF
	Sample 4	Sample 21	Sample 22	Sample 5	Mean	
SiO ₂	62.824	63.918	63.122	62.906	63.193	65.404
TiO ₂	0.000	0.000	0.018	0.000	0.005	0.016
Al ₂ O ₃	18.749	18.316	17.926	17.870	18.215	17.447
FeO	0.180	0.241	0.142	0.160	0.181	0.918
MgO	0.001	0.000	0.000	0.000	0.000	0.290
MnO	0.021	0.014	0.007	0.019	0.015	0.089
CaO	0.179	0.000	0.000	0.019	0.050	0.048
Na ₂ O	2.593	2.078	2.205	2.339	2.304	0.612
K ₂ O	0.000	0.013	0.017	0.025	0.014	0.000
P ₂ O ₅	0.163	0.005	0.033	0.000	0.050	0.233
Cr ₂ O ₃	0.016	0.000	0.000	0.003	0.005	0.005
BeO	13.515	13.550	13.550	13.535	13.537	13.654
Sum	98.241	98.135	97.020	96.876	97.568	98.716
Structural formula based on 18 oxygen atoms						
Si	5.814	5.900	5.891	5.886	5.873	5.979
P	0.013	0.000	0.003	0.000	0.004	0.018
Sum T (1)	5.827	5.900	5.894	5.886	5.877	5.997
Be T(2)	3.005	3.004	3.038	3.042	3.022	2.999
Al	2.045	1.992	1.972	1.971	1.995	1.880
Fe	0.014	0.019	0.011	0.013	0.014	0.070
Mg	0.000	0.000	0.000	0.000	0.000	0.040
Mn	0.002	0.001	0.001	0.002	0.001	0.007
Cr	0.001	0.000	0.000	0.000	0.000	0.000
Sum O	2.062	2.012	1.984	1.985	2.011	1.997
Na	0.465	0.372	0.399	0.424	0.415	0.108
K	0.000	0.002	0.002	0.003	0.002	0.000
Ca	0.018	0.000	0.000	0.002	0.005	0.005
Sum C	0.483	0.373	0.401	0.429	0.422	0.113

The accommodation of femic cations in octahedral sites presumes the accommodation of cations in channels in order to maintain the charge balance: $C_{\square}^{O}Al = C_{(Na,K)}^{O}(Fe,Mg,Mn)$ (1).

The content of alkali cations reaches up to 0.373-0.483 apfu, except for one sample with 0.113-apfu alkalis. For five samples the charge deficit caused by femic cations is compensated at a lesser extent by alkali cations, and this behavior is denoted by the samples position on alkali femic (Fig. 1). For the sample with lower alkali content the substitution (1) is obvious. The sample position along the x axis, which represents $C_{A^{T(2)}}Li = C_{\square^{T(2)}}Be$ substitution (2) indicates that Li is one of the alkali cations in beryl composition and that Na, K, Ca (including Cs and Rb for one of the samples) are mainly involved in compensation of charge balance required by Li accommodation in $T(2)$ sites.

Using ICP-MS analysis, rare alkalis (Li, Rb and Cs) were identified as minor constituents of beryl. The whole content of alkali in beryl is 1.78 wt % (Na = 1.70 %; K = 0.01 %; Li = 0.04 %; Rb = 0.002 %; Cs = 0.02 %). Based on the alkali content, it belongs to a sodium-lithian beryl, for which Na ranges from 0.0 to 2 wt %, Li is under 0.6 wt %, Cs is low (Beus, 1966; Hawthorne & Černý, 1977; Deer *et al.*, 1999).

The relationship between Na/Li and Cs content in beryl is typical of primitive beryl-type pegmatites (Fig. 2; Trueman & Černý, 1982).

The trace element concentrations are generally low, under $0.70 \mu g.g^{-1}$. Higher contents are found for Sr ($1.66 \mu g.g^{-1}$), Ba ($1.18 \mu g.g^{-1}$) and Sc ($1.42 \mu g.g^{-1}$). On the spider diagram (Fig. 3), the trace elements, normalized to continental crust, show that beryl is enriched in Cs and Li and depleted in all other trace elements (Fig. 3).

The concentration level of all 14 rare earth elements is also very low and the total amount attains only $0.2 \mu g.g^{-1}$. However, beryl accommodates LREE ($0.147 \mu g.g^{-1}$) at a higher degree relative to HREE ($0.053 \mu g.g^{-1}$).

The chondrite-normalized pattern of REE of beryl (Fig. 4) displays upwards downwards curved segments, which result in a partly M-shape lanthanides distribution pattern named tetrad effect (e.g. Peppard *et al.*, 1969; Sinha, 1978; Masuda *et al.*, 1987). The tetrad effect on REE patterns is typical of magmatic rocks (and their minerals as well) and is found also in precipitates from fluids. Recent discussions about the tetrad effect deal with highly evolved igneous rocks (e.g. Bau, 1996; Irber, 1999; Irber *et al.*, 1997; Monecke *et al.*, 2002), which typically show an M-shape tetrad effect. Thus, we could infer that beryl pegmatites from Voislova evolved from an igneous system, even if the connection between pegmatites and granites is still unknown in this area.

Fig. 1. (Na+K+Rb+Cs+Ca) – (Fe+Mg+Mn+Cr+V+Sc) relationship in beryl.

1. Beryl from Voislova pegmatites
2. Beryl from Teregova pegmatites
3. Beryl from different types of pegmatites (data from Barton & Young, 2002)
4. Beryl from rare-elements pegmatites, Siberia (Zagorski & Kuznetova, 1990).

Fig. 2. Na/Li – Cs relationship in beryl.

- A - Barren and geochemical primitive beryl-type pegmatites;
 B - Geochemical evolved beryl-columbite and beryl-columbite-phosphate pegmatites; C - Albite-spodumene and complex pegmatites; D - Highly fractionated complex pegmatites.

Fig. 3. Continental crust normalized pattern of trace elements of beryl.

Fig. 4. Chondrite normalized-pattern of REE in beryl.

Conclusions

- analysed beryl from Voislova pegmatite does not fulfill the ideal composition, being enriched in alkalis (mostly in Na and Li) so that it belongs to the sodium-lithian type of beryl;
- although the occurrence of beryl in Voislova pegmatites as large crystals denotes that pegmatites belong to rare-elements pegmatites class, the relationship between Na/Li and Cs content of beryl shows that these pegmatites are not significantly evolved from geochemical point a view;
- the association of beryl with (Fe,Mn) – phosphates and apatite suggests that the main volatile compounds in the crystallization environment were H₂O and P₂O₅;
- beryl is characterized by low concentration of trace elements;
- the chondrite - normalized pattern of REE in beryl suggest an igneous origin of the pegmatites.

Acknowledgments

We are grateful to Dr. Vasile Pomârleanu for providing one of the beryl samples.

References

- Barton, M., Young, S. (2002) Non-pegmatitic deposits of beryllium: Mineralogy, Geology, Phase Equilibria and Origine. In: Beryllium. Mineralogy, Petrology and Geochemistry (P. Ribbe and J. Rosso, Eds.). Mineral. Soc. Amer., *Geochim. Soc.*, **50**, 591-691.
- Bau, M. (1996) Controls on the fractionation of izovalent trace elements in magmatic and aqueous system: evidence from Y/Ho and Zr/Hf and tetrad effect. *Contrib. Mineral. Petrol.*, **123**, 323-333.
- BEUS, A.A. (1966) – *Geochemistry of beryllium and genetic types of its deposits*. Acad. Sci. URSS, Moskva (in Russian, 1960); Engl. Transl. Freeman & Co, 401 p.
- Černý, P. (2002) Mineralogy of beryllium in granite pegmatites. In: Beryllium. Mineralogy, Petrology and Geochemistry (P. Ribbe and J. Rosso, Eds.). Mineral. Soc. Amer., *Geochim. Soc.*, **50**, 405-444.
- Deer, W.A., Howie, R.A., Zussman, J. (1999) Beryl. An introduction to the rock-forming minerals. 2nd Edition. Pearson Education. 116-121.
- Dulski, P. (2000) Reference materials for geochemical studies: New analytical data by ICP-MS and critical discussion of reference values. *Geostandards Newsletter*, **25**, 87-125.
- Govindaraju, K. (1994) Compilation of working values and sample description for 383 Geostandards. *Geostandards Newsletter*, **18**, Sp. Issue, 1-158.
- Hann, H., P. (1987) The pegmatites from South Carpathians (in Romanian). Ed. Acad. Rom., Bucharest, 141 p.
- Irber, W (1999) The lanthanide tetrad effect and its correlation with K/Rb, Eu/Eu*, Sr/Eu, Y/Ho, and Zr/Hf of evolving peraluminous granite suites. *Geochim. Cosmochim. Acta*, **63**, 489–508.

- Irber, W., Forster, H.-J., Hecht, L., Moller, P., Morteani, G. (1997) Experimental, geochemical, mineralogical and O-isotope constraints on the late-magmatic history of the Fichtelgebirge granites (Germany). *Geol. Rdsch.*, **86** (Suppl.), 110-124.
- Maieru, O., Superceanu, C., Apostoloiu, A. (1968) Neue Spodumen und Beryllpegmatite im mittleren süd karpatischen Schiefergebirge (Rumänien). *Geol. Jahrgang*, **17**, 4, Akad. Verlag, 388-397.
- Masuda, A., Kawakami, O., Dohmoto, Y., Takenaka, T. (1987) Lanthanide tetrad effect in nature: Two mutual opposite types W and M. *Geochem. J.*, **23**, 245-253.
- Mârza, I. (1980) Considérations génétique sur les pegmatites du cristallin de Gilău (Monts Apuseni) et la province pegmatitique Carpatique. *An. Inst. Geol. Geofiz.*, **LVII**, 423-432.
- Mârza, I., Pomârleanu-Neagu, E., Pomârleanu, V. (1988) Beryl in some pegmatites in the Bondureasa valley (Apuseni Mountains). *Studia. Univ. „Babeş Bolyai”*, Ser. Geol., **XXXIII/1**, 69-76.
- Monecke, T., Kempe, U., Monecke, J., Sala, M., Wolf, D. (2002) Tetrad effect in rare earth element distribution patterns: A method of quantification with application to rock and mineral samples from granite-related rare metal deposits. *Geochim. Cosmochim. Acta*, **66**, 1185-1196.
- Murariu, T. (2001) The geochemistry of the pegmatites from Romania. *Ed. Acad. Rom.* Bucharest, 356 p. (in Romanian).
- Hawthorne, F.C., Černý, P. (1977) The alkali-metal positions in Cs-Li beryl. *Can. Mineral.*, **15**, 414-421.
- Peppard, D.F., Mason, G.W., Lewey, S. (1969) A tetrad effect in liquid-liquid extraction ordering of lanthanides (III). *J. Inorg. Nucl. Chem.*, **31**, 2271-2272.
- Pomârleanu, V. (1969) Inclusions fluides dans les cristaux de beryl des pegmatites de Roumanie. *Rev. Roum. Géol. Geophys. Géogr.*, Ser. Géologie, **13/2**, 117-121.
- Reed, S.J.B. (1996) *Electron microprobe analysis and scanning electron microscopy in geology*. Cambridge University Press, 215 p.
- Schadler, J. (1930) Ein neues Beryllvorkommen (Teregova, Banat), *Verh. Österr., Geol. Bundesanst.* 10.
- Sinha, S.P. (1978) „Inclined W” and the systematic of the rare earths. *Kemia-Kemi*, **6**, 238-243.
- Superceanu, C. (1957) Rare minerals in granite pegmatites from Banat (I). *Rev. Minelor*, **3**, 140-155.
- Trueman, D.L., Černý, P. (1982) Exploration for rare-elements granitic pegmatites. In : *Mineral. Assoc. Canada, Short Course Hbk* (Černý, P., Ed.), 463-493.
- Zagorski, V.E., Kuzneţova, L.G. (1990) The geochemistry of spodumen-bearing pegmatites and rare-elements bearing metasomatites. *Izd. Nauka*, Novosibirsk, 140 p. (in Russian).

SECONDARY MINERALS FOUND IN OLD MINE GALLERIES FROM ROŞIA MONTANĂ, ROMANIA

Bogdan P. Onac³⁹, Daniel Ş. Vereş, Joe Kearns⁴⁰, Miruna Chirienco⁴¹,
Adrian Minuţ, Radu Breban⁴²

Recent investigation on several secondary minerals formed in old mining galleries in the Cărnic district (Roşia Montană, Romania) enabled us to characterize eight minerals. Out of these eight identified minerals, the discovery of jokokuite is the first reported occurrence in the Carpathians (Szakáll, 2002). A second mineral identified as apjohnite represents a new occurrence in Romania (Udubaşa, 1999). Along with these two rare minerals, some other species of the halotrichite group (pickeringite, halotrichite, apjohnite, and dietrichite), iron sulfates, and K-Al sulfates were identified. Minerals were identified using light microscopy, X-ray diffraction, infrared, and scanning electron microscopy with energy dispersive spectroscopy. The chemical composition of some samples was determined with inductively coupled plasma atomic emission spectrometry. The precipitation of these sulfates is largely controlled by cations substitutions and changes in the temperature, relative humidity, and evaporation along the sampled gallery. The general sequence of sulfate deposition at the investigated site includes the following main steps: (1) oxidation of pyrite by meteoric water seepage through dacites to create ferrous sulfate and sulfuric acid solutions; (2) chemical reaction between dacites and sulfuric acid from which other cations (Al^{3+} , Zn^{2+} , Mn^{2+} , K^{+}) are released into solution; and (3) precipitation of various hydrated sulfates as a result of a combination of dehydration/hydration, oxidation, and neutralization reactions that take place under different microclimatic settings. The specimens are deposited in the Mineralogical Museum of „Babeş-Bolyai” University in Cluj-Napoca, Romania.

Jokokuite - $\text{Mn}^{2+}\text{SO}_4 \cdot 5\text{H}_2\text{O}$ forms pale pink, rosette-like aggregates up to 2-3 cm in length on the walls of an old mining gallery at horizon +958 m, intimately associated with rozenite. The jokokuite crystals have vitreous luster, no cleavage and are easily soluble in water. The average cell parameters obtained on the basis of 29 powder reflections are $a = 6.38(2)\text{\AA}$, $b = 10.70(1)\text{\AA}$, $c = 6.22(2)\text{\AA}$, $\alpha = 97.619(5)^\circ$, $\beta = 110.493(8)^\circ$, $\gamma = 75.88(9)^\circ$. The c cell parameter is smaller than that reported value in the ICDD file 31-836, which may reflect the substitution of Mn^{2+} with Fe^{2+} .

Apjohnite - $\text{Mn}^{2+}\text{Al}_2(\text{SO}_4)_4 \cdot 22\text{H}_2\text{O}$. Found in several samples collected from either floor or walls of old adits. It forms white to yellowish brown or greenish crusts or fibrous and needle-like crystals (up to few centimeters). The unit cell of a representative sample (#1538) as refined by least squares of 48 reflections were found to be $a = 6.266(5)\text{\AA}$, $b = 24.502(2)\text{\AA}$, $c = 21.281(3)\text{\AA}$, and $\beta = 98.692(8)^\circ$. In sample #1541 it appears associated with pickeringite.

Alunogen - $\text{Al}_2(\text{SO}_4)_3 \cdot 17\text{H}_2\text{O}$ appears in association with pickeringite as efflorescences on dietrichite botryoidal aggregates. The prismatic crystals of alunogen are up to 2 mm in length and are extremely thin (<0.5 mm). Up to now, this mineral was mentioned to occur only as efflorescences on metamorphic or igneous rocks (Rădulescu & Dimitrescu, 1966).

Dietrichite - $(\text{Zn}, \text{Fe}^{2+}, \text{Mn}^{2+})\text{Al}_2(\text{SO}_4)_4 \cdot 22\text{H}_2\text{O}$ forms tufted aggregates of acicular crystals and efflorescences along galleries' ceiling. The color is dirty yellow or sometimes greenish. The type locality for this mineral is Baia Sprie (Maramureş, Romania) whereas Roşia Montană represents its second occurrence in Romania.

³⁹ „Babeş-Bolyai” University 1, Kogălniceanu Str., Dept. of Mineralogy, & „Emil Racoviţă” Institute of Speleology, RO-3400 Cluj-Napoca, Romania. E-mail: <bonac@bioge.ubbcluj.ro>.

⁴⁰ The Pennsylvania State University, U.S.A.

⁴¹ „Babeş-Bolyai” University of Cluj-Napoca, 1, Kogălniceanu Str., Dept. of Mineralogy, RO-3400 Cluj-Napoca, Romania. E-mail: <mirona_bv@yahoo.com>.

⁴² S.C. Roşia Montană Gold Corporation S.A., Roşia Montană, Romania.

Halotrichite - $\text{Fe}^{2+}\text{Al}_2(\text{SO}_4)_4 \cdot 22\text{H}_2\text{O}$ was observed as yellowish-brown mammillary aggregates with vitreous luster. It was also found as hair-like efflorescences.

Kalinite - $\text{KAl}(\text{SO}_4)_2 \cdot 11\text{H}_2\text{O}$ is rather abundant in the gallery we investigated and appears as delicate, tiny fibers overlying halotrichite aggregates. Crystals are translucent and if removed from the gallery environment will decompose within minutes into a white milky powder.

Melanterite - $\text{Fe}^{2+}\text{SO}_4 \cdot 7\text{H}_2\text{O}$ forms colorless to translucent, sometimes slightly green fibrous aggregates (up to 4 cm) having vitreous luster. Upon exposure to dry air crystals become white-yellowish and opaque.

Pickeringite - $\text{MgAl}_2(\text{SO}_4)_4 \cdot 22\text{H}_2\text{O}$ was first identified in Romania in Diana Cave, Băile Herculane (Diaconu & Medeşan, 1973). In our investigated occurrence at Roşia Montană the mineral forms shining white to silky thin crystals (3-5 mm in length) covering apjohnite crusts.

Rozenite - $\text{Fe}^{2+}\text{SO}_4 \cdot 5\text{H}_2\text{O}$ is the main component of the rosette-like aggregates found on the ceiling of abandoned adits of the gold deposit at Roşia Montană. The white or colorless fibrous aggregates of rozenite form directly on highly weathered dacites and can reach 3 to 5 cm in length.

References

- Diaconu, G. & Medeşan, A. (1973) Sur la presence du pickeringite dans la grotte Diana (Băile Herculane, Roumanie). *Trav. Inst. Spéol. "Emile Racovitza"*, **XII**, 303-309.
- Rădulescu, D., Dimitrescu, R. (1966) Topographical mineralogy of Romania (in Romanian) *Ed. Acad. Rom.*, Bucharest, 376 p.
- Szakáll, S. (Ed.) (2002) *Minerals of the Carpathians*. Granit, Prague, 480 p.
- Udubaşa, G. (1999) Advances in mineralogy of Romania. *Rom. J. Mineralogy*, **79**: 3-30.

AN UNUSUAL OCCURRENCE OF BERLINITE (AlPO_4) IN THE PHOSPHATE-BEARING SEDIMENTS FROM THE CIOCLOVINA CAVE, ROMANIA

Bogdan P. Onac⁴³, William B. White⁴⁴

This paper reports the first worldwide reported occurrence of berlinite (AlPO_4) formed entirely under sedimentary conditions. Berlinite appears as grayish or colorless fine crystals in vacuoles and along cracks in heavily compacted, phosphate-rich sediments within the Cioclovina Cave, Romania. The X-ray powder diffraction pattern and the hexagonal unit-cell of the Cioclovina berlinite specimen compare well with other published determinations. The lattice parameters are $a = 4.94(4)$, $c = 10.87(1)$ Å, $V = 230.1(3)$ Å³. Microprobe analyses confirmed a nearly ideal formula for berlinite. In situ guano combustion is responsible for the transformation of taranakite and for the dehydration of variscite into berlinite. Berlinite was successfully synthesized from these two minerals at atmospheric pressure and temperatures ranging from 350° to 600°C.

⁴³ „Babeş-Bolyai” University 1, Kogălniceanu Str., Dept. of Mineralogy, & „Emil Racoviţă” Institute of Speleology, RO-3400 Cluj-Napoca, Romania. E-mail: <bonac@bioge.ubbcluj.ro>.

⁴⁴ Materials Research Institute & Department of Geosciences, The Pennsylvania State University, University Park, PA 16802, USA.

THE RELATIONSHIP BETWEEN THE MINERALIZATION OF BRECCIA PIPES AND MINERAL COMPOSITION OF SPELEOTHEMS: EVIDENCES FROM THE CORKSCREW CAVE, ARIZONA (USA)

Bogdan P. Onac⁴⁵, William B. White⁴⁶, Jack W. Hess⁴⁷

Many hundreds of caves hosting breccia pipes are exposed in the massive Mississippian Redwall limestone of the Grand Canyon. The majority of Grand Canyon caves developed during late Mississippian time when the Redwall Formation was emerged. The circular breccia pipes exposed on the cave ceilings are filled with pebbles and cobbles of sandstones, shales and limestones of Paleozoic age (Surprise Canyon Formation), cemented in a carbonatic matrix (Wenrich & Sutphin, 1994). Both clasts and matrix suffered an extensive dolomitization (Wenrich, 1985), being a source of Mg for several minerals deposited in a later stage within the cave. The breccias occurring within this sector of Grand Canyon were mineralized with uraninite, but associated with it are a suite of elements (e.g. As, Mo, V, Ba, Cu, Pb, Fe etc.) that form other ore minerals.

Chemical and mineralogical characterization of the samples was undertaken by X-ray powder diffraction (XRD), X-ray analytical system attached to a scanning electron microscope (SEM-EDAX), electron-microprobe analyses (EMPA), as well as sulfur stable isotope analyses. Morphological observations were undertaken using an optical microscope.

The abundance of gypsum and barite throughout the cave and their isotopically light sulfur isotope composition suggests deposition from hydrothermal sulfuric acid solutions that were also responsible, at least in part, for development of the recent cave passages that dissect older karst breccia bodies. Solution, remobilization and redeposition of some of the ore-related elements have produced a noteworthy assemblage of secondary minerals (i.e., hörnesite, talmessite, carnotite, claudetite, and powellite) that mirrors breccia pipes mineralization. The presence of quartz, calcite, and hörnesite $\pm$ gold in some of the speleothems is considered indicative for a late, low-temperature hydrothermal episode in the cave minerals deposition history. The speleothems in the Corkscrew Cave are diverse in mineralogy, morphology and paragenesis. Mineral deposition varied in both time and space.

Based on the cave morphology and on its assemblage of secondary minerals found in the composition of various types of speleothems, the following events in the cave development and mineral deposition therein were deciphered: (1) Early cave development stage, (2) Formation and mineralization of karst breccia pipes, (3) Reactivation of cave by sulfuric acid speleogenesis, (4) Solution-mobilization-reprecipitation of secondary ore-derived minerals, and (5) Precipitation of minerals under vadose cave conditions.

The mineralogy of speleothems reflects the geochemistry of the groundwater as well as the pH-Eh of the depositional environment. Redox-sensitive elements such as As, Mo, and V are soluble along with U^{6+} in oxidized groundwaters (Drever, 1997); all these elements have been leached from the mineralized breccia pipe, and then transported at short distances by neutral to alkaline, oxidized low-temperature hydrothermal solutions until a redox interface caused their redeposition. Such a redox interface might have been created by either rich organic matter or presence of disseminated pyrite in the limestone bedrock.

Analyzing the relationship between the various minerals forming the coralloids one can understand they can only form under some very specific conditions. These particular settings need be met only in a microenvironment, such as a capillary film at the speleothem surface, close to the breccia pipe body and not anywhere else throughout the main cave passages. Apart for carnotite,

⁴⁵ Babeş-Bolyai" University of Cluj-Napoca, 1, Kogălniceanu Str., Dept. of Mineralogy, RO-3400 Cluj-Napoca, Romania. E-mail: <bonac@bioge.ubbcluj.ro>.

⁴⁶ Department of Geosciences, Pennsylvania State University, USA

⁴⁷ Water Resources Center, Desert Research Institute, Las Vegas, USA

none of the other ore-related minerals described in this paper were cited by Hill & Forti (1997) in their *Cave Minerals of the World* book, nor were they identified in any other cave worldwide ever since.

References

- Drever, J.I. (1997) The geochemistry of natural waters. Surface and ground water environments. Prentice Hall, New Jersey. 436 p. 3rd ed.
- Hill, C.A., Forti, P. (1997) Cave minerals of the world. National Speleological Society, Huntsville, Alabama. 463 p. 2nd ed.
- Wenrich, K.J. (1985) Mineralization of breccia pipes in northern Arizona. *Economic Geology*, **80**: 1722-1735.
- Wenrich, K.J., Sutphin, H.B. (1994) Grand Canyon caves, breccia pipes and mineral deposits. *Geology Today*, 97-104.

CUBIC CoTe₂: SYNTHESIS, STRUCTURE DETERMINATION AND REFLECTED LIGHT INVESTIGATIONS

Franz Pertlik⁴⁸¹

Summary. Single crystals of CoTe₂ were synthesized under hydrothermal conditions at 250°C in concentrated aqueous NaOH solution from the elements cobalt and tellurium. The crystal structure is a representative of structure type “pyrite”, proved by single crystal X-ray work. The reflectance spectra of CoTe₂ were measured with polarized light within the range from 400 to 800 nm, in air and in oil. The comparison of these data with that of the isotypic compounds CoSe₂ (trogtalite) and CoS₂ (cattierite) shows an increase of the reflectance values correlated with the atomic number of the chalcogen atom. The colours change from light cream-brown (CoS₂) to light pink-brown (CoSe₂) to white (CoTe₂) and depend on a shift of the reflectance minimum from ~ 520 nm (CoS₂) to ~ 630 nm (CoTe₂).

Introduction

Investigations within the system Co-Te at temperatures >700°C, pressure one bar, gave no evidence for the existence of the phase CoTe₂ with pyrite type structure (Tengnér, 1938; Haraldsen *et al.*, 1956). On the other hand Bither *et al.* (1968) reported on the synthesis and electrical data of cubic CoTe₂. This compound crystallizes at 65 kilobars and 1200° C. Recent investigations within parts of the system Co-Te-NaOH under hydrothermal conditions now also revealed crystals of CoTe₂ in this cubic modification. For comparison with the compounds CoS₂ and CoSe₂ (cubic modifications) the reflectance values, visible part of the spectrum, were measured. In is worth mentioning, that no reflectance data were reported for pyrite type tellurides until now.

Experimental

Synthesis: Crystals of CoTe₂ were synthesized by hydrothermal treatment of a mixture of each of one gram native cobalt and tellurium in a Teflon lined reactor. The remaining volume of the reactor (total volume ~ 6 cm³) was filled up to 80 % with at room temperature concentrated NaOH solution. After a heating period of one month at 500(10) K single crystals of CoTe₂ up to a diameter of 0.5 mm were maintained. CoTe₂ forms metallic gray octahedra with no pronounced cleavage. The crystal class and the space group of CoTe₂ were proved by single crystal X-ray work, the homogeneity and the stoichiometry by electron microprobe analyses.

X-Ray crystallography: Table 1 presents crystal data, details about intensity measurements, about the structure investigation of CoTe₂ by X-ray work and the final structural parameters. The lattice parameters were calculated from 36 accurate measured 2 θ angles. The collected intensities were corrected for absorption according to crystal size and shape, as well as for Lorentz and polarization effects. The atomic coordinates, given for the isotypic compound FeS₂ (pyrite, Brostigen & Kjekshus, 1969) were used in the first stage of parameter refinement by least squares techniques. Complex scattering functions for neutral atoms were taken from the INTERNATIONAL TABLES (1974).

⁴⁸ Institut für Mineralogie und Kristallographie der Universität Wien. Geozentrum, Althanstraße 14, A-1090 Wien, Austria. E-mail: <franz.pertlik@univie.ac.at>.

Table 1.

Crystal data, details of intensity measurements and structure refinements of CoTe₂
(e.s.d.'s in parentheses)

a = 6.3233(2) Å	Z = 4
V = 252.832 Å ³	Space group: Pa-3; (No. 205)
Measured reflections: 1380 (h, ±k, ±l)	
Unique reflections: 447	
Reflections with Fo > 3 σ Fo: 439	
R value/Rw value; W=[σ Fo] ⁻² : 0.023/0.028	
Variable parameters: 7	
Stoe AED2four circle diffractometer, Mo X-ray tube, graphite monochromator.	
Atomic coordinates and anisotropic displacement parameters.	
2 Co on 4(a) U ₁₁ = 0.0051(2)	8 Te on 8(c) x/a = 0.36809(3)
U ₁₂ = 0.0001(1)	U ₁₁ = 0.0049(1)
	U ₁₂ = 0.0003(1)
Interatomic distances in Å.	
Co – Te = 2.6113(4); 6x	Te – Co = 2.6113(4); 3x
	Te – Te = 2.8916(4); 1x

Reflectance measurements: The CoTe₂ sample was ground with 7 µm SiC abrasive on glass. Polishing was performed with 6 µm diamond paste on nylon polishing cloth and with 1 µm and 0.25 µm diamond pastes on “microcloth” polishing cloth (BUEHLER Ltd.) by hand. The reflectance measurements were performed with linearly polarized light with microscope “Leitz Orthoplan Pol” (using planachromates 20x/0.40, effective N.A. 0.20) and a microscope photometer (photomultiplier S20, type EMI 9558). The diameter of the circular measuring field was 0.05 mm. The reflectance standards used were (W, Ti)C resp. SiC. The reflectance values were measured from 400 to 800 nm in steps of 20 nm in air and in oil immersion (DIN 58884; n = 1.518 at 589 nm) using a Leitz monochromator (Δλ = 7 nm). The reflectance values for CoTe₂ in air and in oil, listed in Table 2, are further represented in graphs (Fig. 1) in comparison with the values for CoSe₂ given by Picot & Johan (1982) resp. CoS₂ given by Demirsoy (1969). The index of refraction n, the absorption coefficient k and the absorption index k have been calculated from the reflectance values in air and in oil according to the formula given by Koenigsberger (1914). The result of these calculations is expressed in three graphs shown in Fig. 2.

Table 2.

Reflectance values for CoTe₂ in the range from 400 to 800 nm together with the values at standard wavelengths (IMA/COM standard) in air (R) and in oil immersion (^{im}R, DIN 58884) in %.

λ(nm)	R	^{im} R	λ(nm)	R	^{im} R	λ(nm)	R	^{im} R
400	51.9	39.8	580	49.7	36.5	760	52.7	40.8
420	51.8	39.7	600	49.5	36.6	780	53.1	40.9
440	51.7	39.5	620	49.4	36.9	800	53.4	50.0
460	51.5	39.3	640	49.5	37.6			
480	51.3	38.8	660	47.7	38.6			
500	51.0	38.2	680	50.4	39.3	470	51.4	39.0
520	50.7	37.6	700	50.8	39.8	546	50.1	36.8
540	50.3	37.0	720	51.5	40.2	589	49.6	36.5
560	50.0	36.7	740	52.2	40.5	650	49.6	38.0

Fig. 1: Spectral reflectance curves of CoTe_2 in air and in oil immersion ($n = 1.518$ at 589 nm). In addition to the reflectance curves for CoSe_2 (Picot & Johan, 1982, with no references to the origin of the sample) as well as the curve for CoS_2 (Demirsoy, 1969) for synthetic crystals are drawn.

Fig. 2: Graph of the index of refraction n , of the absorption coefficient k , and of the absorption index κ ($kappa$) for CoTe_2 calculated from reflectance values in air and in oil

Discussion

The curve of the reflectance values for CoTe_2 shows a minimum at about 630 nm . Therefore, a higher reflectivity at the blue-end of the visible spectrum is responsible for the white colour of CoTe_2 . In the same way, as the minimum of reflectance shifts from CoTe_2 ($\sim 630 \text{ nm}$) to CoS_2 ($\sim 520 \text{ nm}$) the higher reflectivity shifts to the red-end of the visible spectrum. Therefore, the colours of CoSe_2 are light pink-brown and that of CoS_2 light cream-brown (acc. to Picot & Johan, 1982). Marked variations in reflectivity with anion composition are commonly observed in isostructural compounds and minerals of different metals with sulfur, selenium and tellurium. Such variations may be explained by the increased covalent character and electron delocalisation of the metal-chalcogen bond with rising atomic number in the series S-Se-Te (Burns & Vaughan, 1970).

References

- Bither, T.A., Bouchard, R.J., Cloud, W.H., Donohue, P.C., Siemons, W.J. (1968) Transition metal pyrite dichalcogenides. High-pressure synthesis and correlation of properties. – *Inorg. Chem.* **7**, 2208-2220.
- Brostigen, G., Kjekshus, A. (1969) Redetermined crystal structure of FeS_2 (pyrite). – *Acta Chem. Scand.* **23**, 2186-2188.
- Burns, R.G., Vaughan D.J. (1970) Interpretation of the reflectivity behaviour of ore minerals. – *Amer. Mineral.* **55**, 1576-1586.
- Demirsoy, S. (1969) Untersuchungen über den Einfluß der chemischen Zusammensetzung auf die spektralen Reflexionsfunktionen und Mikroindurhärten. – *N. Jb. Miner. Mh.* 1969, 323-333.

- Haraldsen, H., Grønvold, F., Hurlen, T. (1956) Eine röntgenographische und magnetische Untersuchung des Systems Kobalt/Tellur. – *Z. anorg. allgem. Chemie*, **283**, 143-164.
- International Tables for X-ray Crystallography (1974) Vol. IV. Revised and Supplementary Tables. Eds. Ibers, J.A. & Hamilton, W.C. The Kynoch Press – Birmingham, England.
- Koenigsberger, J. (1914) Über Messungen des Reflexionsvermögens und Bestimmung der optischen Konstanten. – *Ann. Physik*, **43**, 1205-1222.
- Picot, P., Johan, Z. (1982) Atlas of the ore minerals. (Transl.: J. Guilloux, Rev.: D. H. Watkinson). Orleans and Amsterdam: B.R.G.M. and Elsevier.
- Tegnér, S. (1938) Über die Phasen CoTe-CoTe_2 und NiTe-NiTe_2 . – *Naturwiss.* **26**, 429.

SOURCE REGIONS OF THE MELT AND FLUID PHASES IN THE PORPHYRY Cu-Au-Mo DEPOSITS AND OTHER VOLCANIC STRUCTURES FROM ALPINE CARPATHIAN CHAIN (ROMANIA)

Ioan Pinte⁴⁹

The aim of the study focuses on several fundamental questions regarding the melt and fluid inclusion studies in the magmatic-hydrothermal environments from the Alpine volcanic area of the Carpathians in Romania. These will be addressed mainly in regard to the primary source regions of saline and sulfide-rich silicate melts and supercritical aqueous fluids, based upon the author's data interpretations (i.g. Pinte, 1993, 1997, 2000, 2002 and unpublished data).

1. Melt and fluid inclusions from the inner part of the porphyry copper deposits were segregated phases evolved during the emplacement of a shallow subvolcanic body, which hosts the main part of the potassic and ore zone. Alternatively, they were transported in a flushing system and injected in the solid country rocks.

2. During the complex evolution of a stratovolcano there is some mafic magmatic input, which periodically supplies sulfur as well as siderophile and calcophile elements. Our petrographic and microthermometric data show evidence that the melt and fluid inclusions originated from deeper mafic magmas and mixed with felsic melts in intermediate magma chamber(s) in several stratovolcanoes.

Finally, it is shown that the silicate melts and supercritical aqueous fluids, segregated from shallow felsic magma chamber(s), remain the most important sources of metals transported by various ligands as chloride - complexes, sulphates etc, mainly in the porphyry copper and epithermal ore deposits. Nevertheless the evolved sulfide-rich melt coming from a deeper mafic magma had enriched substantially the former ore deposits in several late tectono-magmatic events.

References

- Pinte, I. (1993) Microthermometry of the hydrosaline melt inclusions from copper – porphyry ore deposits (Apuseni Mountains, Romania). *Arch. Mineral.* **XLIX**, 165–167. Warsaw, Poland.
- Pinte, I. (1997) The significance of the liquid homogenization temperature in salt melt inclusions. A case study in neogene porphyry copper ore deposits from Metaliferi Mountains (western Romania). *ECROFI XIV*, Nancy, Abstr. vol., 266–267. Nancy, France.
- Pinte, I. (2000) Silicate melt, salt melt, aqueous and CO₂- rich inclusions in felsic and mafic minerals from barren and productive subvolcanic structures from Alpine Carpathian chain (Romania), *Workshop on Melt Inclusions*, Session 6: Magmatic Volatiles, Electronic Program p. 31, 16–18 March 2002, Grenoble, France.
- Pinte, I. (2002) Occurrence and microthermometry of the globular sulfide melt inclusions from extrusive and intrusive volcanic rocks and related ore deposits from Alpine Carpathian Chain (Romania). *Workshop – Short Course on Volcanic Systems, Geochemical and Geophysical Monitoring, Melt Inclusions: Methods, Applications and Problems*, B. De Vivo and R.J. Bodnar (Eds.), Proceedings, Sept. 26–30th, 2002, Seiano di Vico Equense-Napoli, Italy, 177–180.

⁴⁹ Geological Institute of Romania, Cluj-Napoca Branch, P.O. Box 181, RO-3400 Cluj-Napoca 1, Romania, E-mail: <ipinte@email.ro>.

TEXTURAL FEATURES AND MELT INCLUSION TYPES IN ANDESITES AND BASALTIC ANDESITES FROM THE EASTERN CARPATHIANS (CĂLIMANI – GURGHIU – HARGHITA MTS.), ROMANIA

Ioan Pinte⁵⁰, Emilia Mosonyi⁵¹, Zoltan Bardocz²

Oscillatory or discontinuous zoned and sieved plagioclase, resorbed pyroxene and amphibole are important features of the magmatic minerals in volcanic environments. They are related mainly to magma mixing processes and can be successfully used to understand better magma dynamic history in shallow and intermediate magma chambers. Generally, the mafic melt mixes with felsic magmas before or during volcanic eruptions and the final products cover a large spectrum of petrographic types from andesitic to basaltic ones (e.g. Stimac & Pearce, 1992; Hattori, 1993; Kuritani, 2001).

The external zones of the plagioclase phenocrysts show equilibrium either disequilibrium between melt and the magmatic crystals. Thus, the same content of anorthite was observed in the overgrowth zone of all size groups of plagioclase phenocrysts in equilibrium conditions. On the other hand, at disequilibrium, the border of the resorbed plagioclase consisted of several distinct crystallizing centers.

The main types of the melt inclusions found both, in plagioclase and mafic minerals, are silicate melt inclusions (now glass or recrystallized glass) and globular sulphide melt inclusions as zoned and not-zoned primary cavities. Frequently, a supercritical aqueous fluid phase as vapor-rich inclusions (+/- solids) also occurs. The melt and the vapor-rich inclusions were formed as a result of interaction between phenocrysts and sulfur-undersaturated silicate magma during mixing in intermediate magma chambers at ca. 870-1200°C and 1-2 kb, and equilibrium P-T conditions, respectively.

References

- Hattori, K. (1993) High-sulfur magma, a product of fluid discharge from underlying mafic magma: evidence from Mount Pinatubo, Philippines. *Geology*, **21**, 1083 – 1086.
- Kuritani, T. (2001) Replenishment of a mafic magma in zoned felsic magma chamber beneath Rishiri Volcano, Japan. *Bull. Volcanol.*, **62**, 533 – 548.
- Stimac, J. A., Pearce, T. H. (1992) Textural evidence of mafic-felsic magma interaction in dacite lavas, Clear Lake, California. *Amer. Mineral.*, **77**, 795 – 809.

⁵⁰ Geological Institute of Romania, Cluj-Napoca Branch, P.O. Box 181, RO-3400 Cluj-Napoca 1, Romania, E-mail: <ipintea@email.ro>.

⁵¹ „Babeş-Bolyai” University of Cluj-Napoca, 1, Kogălniceanu Str., Dept. of Mineralogy, RO-3400 Cluj-Napoca, Romania. E-mail: <mosonyi@bioge.ubbcluj.ro>.

FLUID INCLUSIONS IN SALT (TRANSYLVANIAN BASIN, ROMANIA)

Vasile Pomârleanu⁵², Ioan Mârza⁵³

The paper presents general morphological aspects of salt crystals, as well as data of fluid inclusions from the Ocna Dej and Slănic Prahova salt deposits.

The salt horizon from Ocna Dej is situated above the Dej volcanic tuff. Chronostratigraphically it belongs to Badenian, to the NN5 zone (Mésároş, Mârza, 1991) or to the NN6 zone (Chira, 2000).

The salt level from the Slănic synclinal (Prahova county) overlaps the Piatra Verde Tuff.

Stratigraphically, both, the volcanic tuff and the salt correspond to the Dej Tuff and to the Badenian salt horizon from Transylvania respectively.

The formation of salt deposits results from successive stages of a specific process taking place at the surface of the salt saturated water in marine basins. Similar processes occur in addition in underground water at various depths as well as at the sediment/water interface.

“Hopper crystals” represent specific crystal growth features, respectively a cubic crystal of salt, in which the faces of the cube have grown more at the edges than in the center, forming as a consequence after several growth steps a pyramidal outline. The genesis of “hopper crystals” is explained by the formation of a fine-grained film of salt at the water surface, due to evaporation and to a continuous increasing solution density. With the procedure of growth of the incipient crystal develops a submerging trend; still the crystals remain at the water surface for a relatively long time, due to the surface tension of the solution. Because one of the crystal faces is in permanent contact with the high concentration and density solution, the crystal growth proceeds vertically.

The salt crystals reach up to 1-2 cm in length (Pl. I, Fig. 1-5). These crystals join each other along small areas, generating crystal aggregates that float at the water surface. If the water surface is hydromechanically disturbed (waves, water currents etc.), the salt aggregates are fragmented and may sink as independent crystalline bodies. The occurrence and high frequency of “hopper crystals” indicate a deposition of the salt under calm hydrodynamic conditions. By contrast, their fragmentation before reaching the mature stage suggests an agitated hydrodynamic environment (Pomârleanu & Neagu, 2003).

Hopper halite crystals were noticed in both deposits, in Ocna Dej, and in Slănic Prahova. The habit of the crystals is outlined by the presence of abundant fluid and solid inclusions (Pl. II; Fig. 1, 2), which underwent some necking down processes (Pl. II; Fig. 3).

A second stage of primary growth of halite crystals takes place at the bottom of a marine basin. The submerged crystals lying at the water/sediment (mud) interface or at various depths within the mud continue to grow. Such a process was observed in NaCl-supersaturated water of the Pata Rât lake, in the neighborhood of Someşeni (Cluj district). At the bottom of this lake – rich in sapropelic mud, numerous “hopper crystals” were identified; some were isolated, others form aggregates. Under these circumstances, a “competitive” crystal growth takes place. As a consequence, layers of salt may form at the bottom of the basin at the final stage of crystallization (Pl. I; Figs. 2, 3).

During the diagenesis, the original salt morphology may be dissolved as a result of the influence of an undersaturated marine water flux, or of meteoric/surface waters. The dissolution starts along the contacts between the crystals and gradually forms cavities within the crystals. Sometimes the dissolution of primary salt evolves in such an advanced manner, that the core of the recrystallized halite (secondary halite; Pl. I; Fig. 4) preserves only very small relics of the original morphology (Pl. II, Fig. 1). The recrystallized, translucent salt contains elongated, monophasic and biphasic fluid inclusions (Pl. I; Fig. 5).

⁵² Geological Institute of Romania, Caransebeş Str., RO-7000 Bucharest, Romania

⁵³ Babes-Bolyai University, Department of Mineralogy, 1, M. Kogălniceanu Str., RO-3400 Cluj-Napoca, Romania.

In these salt crystals fluid inclusions were studied. The measured values of homogenization (*th*) and decrepitation (*td*) temperatures vary between 80 and 220°C. A significant number of inclusions decrepitated before reaching the complete homogenization. Experimental work (Pomârleanu & Neagu, 2003) showed that if a large amount of halite crystals underwent plastic deformation within the salt deposit, they were not anymore suitable for geothermometric investigations. Furthermore, a partial loss of the fluid inclusions content occurred during heating, due to the remarkable plasticity of salt and probably to the presence of cryptofractures. During the study of fluid inclusions some general aspects were noticed: gaseous inclusions were predominant in the halite crystals from the Slănic Prahova deposit, whereas liquid inclusions prevailed in the Ocna Dej deposit (Pl. II; Figs. 5 and 6). The pH of the solutions probably ranges the values of 3 – 6 (acidic to almost neutral), similar with the values mentioned in other salt deposits by Roedder (1984)

The geothermometric data as well as the chemical composition of the fluid-gaseous inclusions in salt deposits (primary salt in particular) may be used as an important tool in evaluating the environment and source of formation of the major masses of halide rocks in paleomarine and lacustrine basins, i.e. exogene vs. hypogene (volcanic).

The present paper is an argument for the importance of further geothermometric studies of the fluid inclusions in salt deposits located in the three main regions with salt deposits in Romania (Transylvania, Maramureş and pre-Carpathian area), taking into account that recent trends in this field of research focus on the *migration of inclusions according to the variation of thermal gradients*.

References

- Chira, C. (2000) The Miocene calcareous nannoplankton and molluscs in Transylvania, Romania (in Romanian). *Ed. Carpatica* 183 p., Cluj-Napoca
- Dubessy, J., Ramboz, C. (1987) The history of organic nitrogen from early diagenesis to amphibolite facies: mineralogical, chemical, mechanical and isotopic implication . *IXth Simposion on Fluid Inclusions. Abstract*, 31-32, Univ. Porto.
- Roedder, E. (1984) The fluid in salt. *Amer. Min.*, **48**, 413-439.

PLATE I.

Plate.1. Hopper halite crystals from Ocna Dej. 1-4: different aspects of the Hopper crystals from the Ocna Dej salt deposit. 5 – Monophasic and biphasic inclusions in halite crystal.

PLATE II.

Plate II. Fluid inclusions in halite crystals. Monophasic (1) and biphasic (2) inclusions in the Ocna Dej salt. The necking down phenomenon (3). Multi-phase inclusions from Slanic Prahova primary salt surrounded by recrystallized one with fluid inclusions containing volatile elements (4, 5, 6).

XRD INVESTIGATION OF THE CLAY FRACTION IN SOME ROMANIAN ZEOLITIC VOLCANIC TUFFS AND DIATOMITES

Dana Pop⁵⁴, Ioan Bedelea⁵⁵, Horea Bedelea⁵⁶ (

Introduction

Between 1998-2000, the most representative occurrences of zeolitic volcanic tuffs and diatomites from Romania were studied in the view of their possible usage for building materials with improved mechanical and insulating properties, in the frame of an EC funded INCO-COPERNICUS project. The clay fraction (< 2 microns) represents a relatively small amount in the whole rock mass; still, its mineralogy may provide clues for understanding the environment of formation, and could be used as a criterion for correlating various horizons and outcrops. On the other hand, the presence of certain clay species even in small amounts may influence the behavior of the raw materials in applications like production of blended cement and concrete, or synthesis of wollastonite or diopside.

Materials and methods

Five samples of zeolitic tuffs, from Slănic (Prahova distr.), Bârsana (Maramureş distr.), Mirşid (Sălaj distr.), Petreştii de Jos, (Cluj distr.), and Pâglişa (Cluj distr.) – the last three representing the same Dej Tuff Horizon, two samples of diatomite from Filia (Covasna distr.), and Adamclisi (Constanţa distr.) and additionally one sample consisting of a green clayey material filling voids in the tuff from Pâglişa were separated into the >2 microns and <2 microns fractions, by using the classical sedimentation method. The < 2 microns fractions were submitted to a standardized physical-chemical procedure established at the Clay minerals Laboratory of the Geological Survey of Denmark (GEUS), Copenhagen. The chemical procedure consisted of Mg²⁺- and K⁺- saturation by using 1M chloride solutions. From each fraction, four distinctive specimens were prepared as oriented films: K⁺-saturated, air dry; K⁺-saturated, heated at 300 °C for one hour; Mg²⁺-saturated, air dry, and Mg²⁺-saturated, solvated with glycerol. The measurements were performed on two types of Phillips goniometers: 1050, and PW 3040 respectively, both using Co-K_α radiation and β-filter.

Results and discussion

In all the studied zeolitic volcanic tuffs, the clay minerals are mainly represented by smectite and/or illite. The samples from Bârsana and Slănic are characterized by the presence of illite, which in the first case is better crystallized. Accordingly, the green colour of these tuff levels is probably due to a Fe-rich illite-type phase. The samples representing the Dej Tuff Horizon (Mirşid, Petreştii de Jos, and Pâglişa) show heterogeneous clay compositions: at Mirşid smectite is the main phase, while at Petreştii de Jos and Pâglişa both illite and smectite are present. In almost all the samples (except for Petreştii de Jos), a prominent peak around 11°50' (2θ) was noticed, which was attributed to clinoptilolite (020). A smectite-type phase is dominant in the green material consisting the infillings in the tuff from Pâglişa.

The clay mineral association in the diatomite samples is clearly distinctive. The diatomite from Filia contains poorly crystallized smectite, illite and kaolinite, while that from Adamclisi is dominated by smectite, illite being present in small amounts.

At this stage of the interpretation, the following can be stated:

⁵⁴ "Babeş-Bolyai" University, Mineralogical Museum, 1, Kogălniceanu Str., RO-3400 Cluj-Napoca, Romania. <danapop@bioge.ubbcluj.ro>.

⁵⁵ Faculty of Geography-History, Dimitrie Cantemir University, 22, Cuza-Vodă St., RO-4300 Tg. Mureş, Romania

⁵⁶ "Babeş-Bolyai" University of Cluj-Napoca, 1, Kogălniceanu Str., Dept. of Geology, RO-3400 Cluj-Napoca, Romania. E-mail: <bedelea@bioge.ubbcluj.ro>.

1. The “classical” procedure of separation of the < 2 microns fraction does not prevent the presence of non-clay minerals, especially zeolites (clinoptilolite). Therefore, the separation of the < 0.2 microns fraction is recommended;
2. The studied zeolitic tuffs are dominated by the presence of smectite and/or illite in the clay fraction;
3. The same tuff level can show distinctive clay mineralogy in different occurrences, probably as a result of the specific local detrital supply or alteration processes;
4. Distinctive mineral phases (smectite-type vs. illite-type) could be responsible for the green color of some zeolitic tuffs;
5. The diatomites show a heterogeneous clay mineral composition, according to the specific conditions of formation.

This work was financed by the INCO-COPERNICUS 15CT96-0712 project.

THE WEBSITE OF THE MINERALOGICAL MUSEUM, BABEŞ-BOLYAI UNIVERSITY, CLUJ-NAPOCA, ROMANIA

Dana Pop⁵⁷, Cosmin Stremţan⁵⁸

Current status of the Romanian Natural Sciences museums websites

Among the 700 records in the on-line national database of Museums and Collections in Romania, set up by the Institute for Cultural Memory (CIMEC) (<http://www.cimec.ro/>, last update February 2003), there are 75 institutions belonging to the Natural Sciences profile. Fourteen of them include mineralogy and/or geology collections, and three represent mineralogical museums proper (Museum of Mineralogy, Baia Mare; Gold Museum, Brad; "Constantin Gruescu" Iron Aesthetic Mineralogy Museum, Ocna de Fier). The list is not exhaustive, for example most of the academic collections, such as the Mineralogical Museum of the Babeş-Bolyai University, are missing.

Based on a general Internet search, it can be stated that currently only a few Romanian Natural Sciences museums have their own website. Among those hosting mineral collections, one can mention: "Grigore Antipa" Natural History Museum from Bucharest, Museum of Oltenia (including the Department of Natural Sciences) from Craiova; "Iron Gates" Region Museum (including the Department of Natural Sciences) from Drobeta Turnu Severin; Prahova County Natural Sciences Museum from Ploieşti, Bruckenthal Museum of Natural History from Sibiu, and Museum of Banat (including the Department of Natural Sciences) from Timişoara. Currently, the only website of a mineralogical museum is that of the Museum of Mineralogy from Baia Mare (<http://www.sintec.ro/muzeu/muzeu1.html>).

It is self evident that there is still much to be done for increasing public knowledge, awareness and interest on this type of museums. A website is definitely an up-to-date, attractive, efficient and not expensive tool.

The website of the Mineralogical Museum, Babeş-Bolyai University in Cluj-Napoca (MMBBU)

The MMBBU is the largest and most complete mineralogical museum in Transylvania, hosting about 16,500 mineral specimens from Romania and abroad, illustrating more than 800 mineral species. Starting with the autumn of 2003, the MMBBU have introduced its own website, in both Romanian and English versions. The website was designed to cover the main fields of activity of the Museum, as well as the presumable interests of various categories of visitors (especially pupils of ages between 10-18, and professionals from Romania and from abroad). The website has the following structure: Homepage, About us, Location, Permanent exhibitions & collections, Temporary exhibitions, What's new (New acquisitions), Exchanges & Donations, Education, Visitor information, Favorite links, Photo gallery. It also provides a quick search engine and a virtual visitor's counter. Details on the content of some of the sections are given below.

The permanent exhibitions and collections (12 in number) are briefly presented as text and images (general views, or individual specimens). On a national basis, some of them have a special value: the only meteorite collection, the second largest gold collection, one of the finest cut gemstone collection, the most complete Romanian gemstones collection.

⁵⁷ "Babeş-Bolyai" University, Mineralogical Museum, 1, Kogălniceanu Str., RO-3400 Cluj-Napoca, Romania.
E-mail: <danapop@bioge.ubbcluj.ro>.

⁵⁸ „Babeş-Bolyai” University of Cluj-Napoca, 1, Kogălniceanu Str., Faculty of Biology and Geology, RO-3400 Cluj-Napoca, Romania.

A relatively new initiative, *i.e.* setting up temporary exhibitions, is hoped to highlight some topics of general interest, such as “Minerals and the human health” (currently on display). The virtual visitors are invited to make their suggestions concerning further topics.

The “What’s new” section provides a list of the acquisitions in the previous year, as a renewed invitation for previous visitors, but also as information for professionals on the museum’s offer of hosting scientifically valuable mineral specimens (type specimens, rare species).

Exchanges and donations are the main procedures providing new samples for the Museum. Some information on the Museum’s policy regarding donations, lists of specimens from Romania and abroad offered for exchanges and categories of minerals that are of interest for the Museum are included in this section.

Education in the Museum is one of the main goals of the MMBBU. According to the target group, several recent projects have been developed. They are included in an electronic form: Quiz-test on Physical properties of minerals (for students), Top-10 of the MMBBU specimens (for high school pupils), and Treasure search (for primary and secondary school pupils).

The Photo gallery consists of 55 spectacular images of mineral specimens from the MMBBU collections, which can be searched by mineral name, and occurrence.

PYROXENITE – MARBLE INTERACTION, A PETROGENETIC GRID AND A WORKING EXAMPLE FROM THE INĂU CRYSTALLINE ISLAND, MARAMUREŞ, ROMANIA

Daniel M. Radu⁵⁹

An exotic occurrence of a pyroxenite xenolith, hosted by dolomitic marbles is located in the Inău crystalline massif, Maramureş (Fig. 1). The interaction pyroxenite-marble at different PT has not yet been considered by the geological literature. Petrogenetic grids for ultramafic rocks were presented by Will *et al.* (1990), Schmaedicke, Okrusch (1997), Schmaedicke (2000). The most appropriate system (CaO-MgO-SiO₂-H₂O-CO₂) was considered by Spear (1993).

The mineralogical assemblages of the pyroxenite xenolith, the host rock and interaction zones are presented in the table 1.

Figure 1. Geological map of Inău and Preluca crystalline massifs with the location of the pyroxenite xenolith.

Table 1.
The mineralogical assemblages of the pyroxenite xenolith, the host rock and interaction zones.

Nucleus (xenolith)	Intermediate zone	External zone	Host rock
Pyroxenite	Reaction skarn	Reaction skarn	
enstatite, forsterite, chromite, pentlandite	clinoenstatite, forsterite, chromite, pentlandite; antigorite, talc, chlorite, anthophyllite, tremolite, dolomite	talc, antigorite, dolomite	dolomite

Considering the mineralogical assemblages presented above, and the ratio MgO/FeO = 5.14 of the pyroxenite, the system CaO-MgO-SiO₂-H₂O-CO₂ (CMSh-CO₂) will be a good approximation of the geological reality. The end-members that are used to calculate the petrogenetic grid are as follows: calcite, dolomite; forsterite, enstatite, diopside; anthophyllite, tremolite; antigorite, talc, chlorite; quartz; H₂O, CO₂. The Thermocalc 3.1 software (Holland & Powell, 2001) using the internally consistent thermodynamic database of Holland & Powell (1998) was used to calculate a petrogenetic grid for the system CMSh-CO₂, with a fluid composition of XH₂O = 0.95 and XCO₂ = 0.05 (Fig. 2).

⁵⁹ S.C. SAMAX Romania S.R.L. 15/1 Moldovei Str., RO-4800 Baia Mare, Romania.
E-mail: <daniel_radu_geo@hotmail.com>.

Fig. 2. Petrogenetic grid for the system CMSH-CO₂. Aluminium silicates are presented only for reference. Two possible paths, (a) and (b) were considered, according to the evolution deduced from aluminosilicates assemblages (micaschists, gneisses, amphibolites) of Preluca massif. The star represent calculated PT for initial pyroxenite-marble interaction.

Often, when the carbonates are in excess, the partial pressure of CO₂ is higher than 0.05. T- CO₂ diagrams were calculated for pressures of 5, 10 and 15 kbars (Fig. 3) that represent a good guide to characterize the fluid composition. Most of the mineral assemblages are stable at temperatures of 50 to 100°C, lower in the presence of a fluid mixture H₂O + CO₂ relatively to a pure water fluid.

The microprobe analyses on enstatite, forsterite, talc, tremolite, anthophyllite, and chlorite were used to extract the maximum PT for the initial pyroxenite mineral assemblage. For XCO₂ = 0.10; 0.15; 0.20, the calculated PT values are: 13.2-708; 11.8-703; 10.9-698 (kbars - °C); these are similar to the maximum PT calculated by Radu (2003) for the Preluca massif, using mineral assemblages from micaschists, gneisses and amphibolites.

Fig. 3. T-CO₂ diagrams for the system CMSH-CO₂. (a) P_{fluid} = 5.0 kbars. (b) P_{fluid} = 10 kbars. Invariant points with the symbols as in figure 2.

Fig. 3. (continued) (c) $P_{\text{fluid}} = 15$ kbars.

We assume that the pyroxenite xenolith represents a fragment from the lower crust/upper mantle, reacting with the dolomitic marble when the uplift episode of the Inău crystalline massif started. The presence of anthophyllite suggests a short metamorphic episode near the univariant curve kyanite-sillimanite at $T > 600^{\circ}\text{C}$. This is confirmed by the rare presence of sillimanite in the mineral assemblages from Preluca micaschists and gneisses.

Acknowledgments

The author is grateful to the Mineralogical Institute, Würzburg, Germany, for microprobe analysis, and to Dr. Nigel Cook, Geological Survey of Norway, for the help and discussions on preliminary works. The suggestions of Prof. Dr. Marin Şeclăman, University of Bucharest, deserve my sincere appreciation.

References

- Holland, T.J.B., Powell, R. (1998) An internally-consistent thermodynamic dataset for phases of petrological interest. *Journal of Metamorphic Geology*, **16**, 309 - 343.
- Radu, D.M. (2003) Thermobarric evolution of the metamorphic rocks from Preluca, Țicău and Codru massifs, NW Transylvania (in Romanian). *Ph.D. Thesis*, Bucharest University, 220 p.
- Schmaedicke, E., Okrusch, M. (1997) Phase relations of calcic amphibole-orthoamphibole-chlorite-talc assemblages, with applications to ultramafic rocks from KTB pilot hole, Bavaria. *Geologische Rundschau, Supplement*, **86**, S212-S221.
- Schmaedicke, E. (2000) Phase relations in peridotitic and pyroxenitic rocks in the model systems CMASH and NCMASH. *Journal of Petrology*, **41**, 1, 69-86.
- Spear, F.S. (1993) Metamorphic phase equilibria and Pressure-Temperature-Time paths. *Mineralogical Society of America, Monograph*, 800 p.
- Will, T. M., Powell, R., Holland, T.J.B. (1990) A calculated petrogenetic grid for ultramafic rocks in the system CaO-FeO-MgO-Al₂O₃-SiO₂-CO₂-H₂O. *Contribution to Mineralogy and Petrology*, **105**, 347-358.

THE GEOCHEMISTRY OF APATITE FROM RĂZOARE PEGMATITES (ROMANIA)

Smaranda Rădăşanu⁶⁰, Titus Murariu¹, Uwe H. Kasper⁶¹, Thorbjorn Schoenbeck²

Introduction

The pegmatites from Răzoare are located in gneisses from Răzoare Formation, one of the four metamorphic formations which were distinguished in the Preluca Mountains by Rusu *et al.* (1981) and Balintoni (1997). The Preluca Mountains represent one of the seven crystalline islands located in NW Transilvania. The metamorphic rocks, which build up these mountains, belong to Baia de Aries group (Rusu *et al.*, 1981; Balintoni, 1997).

Along the massif, the metamorphic grade decreases from SE towards NW and varies from sillimanite to biotite zones of the amphibolite facies. The first metamorphic event (M_1) generated staurolite and kyanite as index minerals, and the second one yielded sillimanite. The second metamorphic event (M_2) acted only upon the Răzoare Formation. The P-T calculations (Radu, 1997) show that the metamorphic peak corresponds to 530⁰-600⁰C temperature and 6-8 kbar pressure. The presence of kyanite in some micaschists indicates that P-T conditions were above the kyanite-sillimanite invariant curve and they fit the initial granitic migmatitic melting (Radu, 1997).

The pegmatite emplacement is connected to M_2 event (Balintoni, 1997; Radu, 1997). Generally, the pegmatites have a granite composition and consist of quartz, feldspars and muscovite as rock-forming minerals and of biotite, tourmaline, garnet, apatite and zircon as accessory minerals. At Răzoare, the pegmatites are enriched in accessory minerals comparative with the pegmatites found in the other three formations.

The apatite is found as large green prismatic crystals (up to about 3 cm in length) or rounded grains associated with feldspars (albite prevailing over K-feldspar), garnet and quartz or, as disseminated crystals among albite, perthite, quartz, tourmaline and garnet, when it can be observed only in thin sections.

This paper deals with the geochemical features of apatite from the Răzoare. Previous researches carried out by Macovei & Pomârleanu (1980) and Mârza *et al.* (1986) using IR and XR spectra stated that fluorine-apatite is the main component of apatite from the Răzoare pegmatites. So far, a comprehensive geochemical study regarding major and trace components of the apatite from the Răzoare pegmatite has not been performed. The available data are qualitative and show that this mineral concentrates same trace elements such as Mn, Fe, Mg, Al, Cu, La, Sm, Eu, Dy (Mârza *et al.*, 1986) or (UO₂)²⁻ (Macovei & Pomârleanu, 1980).

Analytical methods

The major components of seven apatite samples from Răzoare pegmatites were analyzed using XRF and EMP methods. The trace elements were determined for three apatite samples using ICP-MS method.

XRF analyses were performed with a Philips PW2400 X-ray spectrometer, using the analytical procedure "oxiquant". Seventy-two natural rocks and clays were used to determine the calibration curves of the pertinent elements.

EMP were carried out with a JEOL JXA-8900 instrument using operating current of 20 nA and accelerating voltage of 20 kV. X-ray intensities of the alkalis, the minor (Ti and Mn) and the major elements were counted for 5s, 40s, and 60s respectively. In order to minimize losses of Na and K, the beam diameter was expanded to 10 mm. Components were standardized using natural minerals, glasses of natural rocks and synthetic oxide compounds. The results were corrected using the ZAF procedure (Reed, 1996).

⁶⁰"Al.I.Cuza" University, Department of Mineralogy and Geochemistry, 20A, Carol I Blv., RO-6600, Iasi, Romania. E-mail: <smaranda@geo.uaic.ro>, <tmurariu@geo.uaic.ro>.

⁶¹ Geologisches Institut, Universität zu Köln, 49a, Zùlpicher Str., D-50674 Köln, Germany. E-mail: <hu.kasper@uni-koeln.de>.

The trace elements, including all REE, were analysed by ICP-MS method (Perkin Elmer/Sciex ELAN 6000 ICP-MS, quadrupole mass spectrometer). Measurements of element concentrations were performed using as internal standards Ru-Re (10 ng/ml) to minimize drift effects and two calibration solutions (high purity chemical reagents). A batch of 5 - 7 samples was bracketed by two calibration procedures. Accuracy and precision of determinations were checked with certified reference materials (CRM) (Govindaraju, 1994; Dulski, 2000).

All the analyses were performed at the Geological Institute of Köln University, Germany.

Geochemistry of apatite

The apatite can be described by the general formula $A_5(XO_4)_2Z$ (Piccoli & Candel, 2002). The A-site accommodates large cations (e.g. Ca, Sr, Na, Mn^{2+} , Fe^{2+} , Mg, REE, Na, Pb) and comprises two sites that exhibit VII-fold [Ca(2)] and IX-fold [Ca(1)] coordination (Hughes & Rakovan, 2002). The X-site, primarily occupied by P^{5+} (as PO_4^{3-}) exhibits IV-fold coordination and can accommodate other small highly charged cations (e.g. Si^{4+} , S^{6+} , As^{5+} , V^{5+}). The Z site is occupied by the halogens F⁻ and Cl⁻, as well as OH⁻.

The results of EMP and XRF analyses of apatite from the Răzoare pegmatites are reported in Table 1. Because the concentrations of F, Cl and H₂O were not determined, the structural formula was calculated on the basis of 12 oxygen cations. Thus, the deficit of the oxide sum is related to undetermined volatiles and at the same time to the high level of concentration of some elements, as ICP-MS data show (Table 2).

Table 1.

The composition of apatite from the Răzoare pegmatites.

Oxides(%) / Sample	Ap 9	Ap 14	Ap 1	Ap 2	Ap 3	Ap 4	Ap 5
P ₂ O ₅	41.931	41.597	41.292	41.176	41.413	41.01	41.222
SiO ₂	0.022	-	-	-	-	-	-
Al ₂ O ₃	-	-	0.004	-	0.002	-	0.002
TiO ₂	-	-	0.012	-	0.035	-	0.012
FeO	0.151	0.181	0.207	0.147	0.157	0.191	0.175
MgO	-	0.009	0.016	0.03	0.007	-	0.013
MnO	1.085	1.090	1.243	1.158	1.206	1.076	1.171
CaO	55.698	55.397	55.715	55.542	55.177	55.988	55.605
Na ₂ O	0.106	0.116	0.182	0.080	0.095	0.154	0.128
K ₂ O	0.032	-	-	-	0.02	-	0.005
Cr ₂ O ₃	0.052	0.013	0.002	-	0.014	0.005	0.005
Sum	99.077	98.403	98.673	98.133	98.126	98.424	98.339
Structural formula calculated on 12 oxygen atoms base							
P	2.846	2.844	2.824	2.829	2.841	2.815	2.827
Si	0.002	0.000	0.000	0.000	0.000	0.000	0.000
Sum X site	2.847	2.844	2.824	2.829	2.841	2.815	2.827
Fe	0.010	0.012	0.014	0.010	0.011	0.013	0.012
Mg	0.000	0.001	0.002	0.004	0.001	0.000	0.002
Mn	0.074	0.075	0.085	0.080	0.083	0.074	0.080
Ca	4.784	4.793	4.822	4.829	4.790	4.863	4.826
Cr	0.003	0.001	0.000	0.000	0.001	0.000	0.000
Na	0.016	0.018	0.029	0.013	0.015	0.024	0.020
Sum A site	4.887	4.899	4.952	4.935	4.900	4.975	4.940

- concentration below the limit of detection.

From EMP and XRF analyses, the main components of apatite are Ca, P and Mn, followed by Na and Fe. The concentration of MnO and FeO (table 1) fit in with the characteristic range of these oxides in apatite from granite pegmatites associated with granites (0.05-7.9 % MnO; 0.01-2.67 % FeO; Piccoli & Candela, 2002).

Generally, the concentration of Si does not attain the detection limit. As minor components, Al and Ti are found in three samples, with Ti in higher concentration than Al.

The accommodation of Mn, Fe or Mg in Ca sites results from simple exchange, whereas the substitution of Na for Ca requires other elements contribution in order to retain charge balance. The structural formula of the investigated apatite shows that Cr exceeds Al. So far, the single substitution mechanism proposed for the accommodation of Al in apatite is described by the following relationship: NaAlCa_2 (Sha & Chappell, 1999).

Commonly, due to chemical composition and crystal chemistry, the apatite from granite pegmatite is the main host for Y, REE, Th, and U.

Table 2.

**Trace elements of apatite from
Răzoare pegmatites**

$\mu\text{g.g}^{-1}$	Ap.X*	Ap.03	Ap.02
Li	-	49.21	4.97
Rb	12.00	9.24	11.95
Cs	0.17	0.17	0.34
Sr	506	509	219
Y	1950	1565	1564
Zr	5.70	44.64	25.13
Nb	0.10	0.53	1.03
La	166.52	159.00	173.50
Ce	556.11	537.22	562.36
Pr	90.65	84.51	86.53
Nd	432.05	435.01	429.32
Sm	178.87	181.95	172.22
Eu	18.10	20.31	16.28
Gd	277.44	227.73	220.82
Tb	49.52	50.11	48.29
Dy	374.95	310.91	316.59
Ho	68.29	58.30	59.05
Er	177.37	160.72	164.85
Tm	25.11	23.99	24.86
Yb	161.97	158.73	161.17
Lu	20.42	20.50	21.77
Hf	5.11	5.88	8.28
Ta	0.57	0.58	0.71
Pb	90.00	96.12	53.63
Th	1.68	1.94	2.61
U	113.45	104.51	103.31

*Rădăşanu (2002)

The apatite from Răzoare pegmatites is enriched in Sr, Y, REE, Th, U and Pb and depleted in Li, Rb, Cs, Nb, Ta (rare elements) (Table 2). The accommodation of Sr in apatite results from simple substitution for Ca in Ca (2) site (Hughes *et al.*, 1991a); in the investigated samples, strontium attains a high concentration level (table 2).

The REE and Y also occupy Ca site in apatite; LREE prefer the Ca (2) site, whereas HREE and Y prefer the Ca(2) (Hughes *et al.*, 1991b). In order to maintain the charge balance, the substitution of REE and Y for Ca involves coupled substitutions as follows: $\text{REESi}(\text{CaP})_{-1}$, REENaCa_2 (Ronsbo, 1989). In the analyzed samples Si does not attain a significant content in A-site. Therefore, the first substitution is subordinated to the second one that also explains the Na enrichment.

Among the rock-forming and accessory minerals, apatite from Răzoare pegmatites is the main REE and Y concentrator (table 2), being followed by garnet ($49\text{--}287 \mu\text{g.g}^{-1}$ REE; $126\text{--}589 \mu\text{g.g}^{-1}$), feldspars ($0.77\text{--}17.16 \mu\text{g.g}^{-1}$ REE; $0.3\text{--}1.2 \mu\text{g.g}^{-1}$), biotite ($0.87\text{--}5.13 \mu\text{g.g}^{-1}$ REE; $0.1\text{--}2.5 \mu\text{g.g}^{-1}$ Y), tourmaline ($3.63 \mu\text{g.g}^{-1}$ REE; $1.49 \mu\text{g.g}^{-1}$ Y) and muscovite ($0.60\text{--}7.15 \mu\text{g.g}^{-1}$ REE; $0.1\text{--}6.7 \mu\text{g.g}^{-1}$ Y) (Rădăşanu, 2002).

Despite the interest for the uranium in apatite, the basic crystal chemistry of its structural substitution is still unknown. The analyzed apatite is enriched in U comparatively with Th; the maximum concentration level of these trace elements is also found in apatite (Table 2). Low concentrations are detected for Zr, Hf, Nb and Ta (Table 2). The ratios Zr/Hf (1.11–7.59) and Nb/Ta (0.22–1.44) are typical of apatite from granite pegmatites (1.0–27.2 Zr/Hf, 0.2–2.2 Nb/Ta; Zhang & Liu, 2001), which derived from a magmatic system. Thus, these ratios for apatite suggest that pegmatites from Răzoare

crystallized from a melt.

Based on the Y/Ho, Zr/Hf ratios and the shape of chondrite-normalized patterns of REE in rocks and minerals, the nature of crystallization environment can be identified (Bau, 1996, 1997; Irber, 1999; Gramaccioli & Pezzotta, 2000; Monecke *et al.*, 2002). In a volatile enriched environment, the trace elements behavior is not only controlled by their charge and radius. Thus, elements with similar properties, such as Y–Ho, Zr–Hf, Nb–Ta and Sr–Eu, do no longer show a coherent behavior and their ratios deviate from that of typical chondrites (28 Y/Ho; 39 Zr/Hf; 17.57 Nb/Ta; 139 Sr/Eu). At the same time, chondrite-normalized patterns of REE are irregular, showing four upwards-downwards segments referred to as lanthanides tetrad effects. Anomalous behavior of Y and REE and of Zr and Hf, hosted by different minerals is due to the chemical complexation

with a wide variety of ligands such as non-bridging oxygen, F, B, P, Cl etc., so that the non-chondrite ratios denote a high activity of complexing factors in the depositing fluids (Bau, 1996; Gramaccioli & Pezzotta, 2000).

Our results regarding Y/Ho, Zr/Hf, Nb/Ta and Sr/Eu ratios show certain contrasts. The first ratio (26.48-28.55 Y/Ho) does not significantly deviate from chondrite ratio, but the values of the next ratios (1.11-7.59 Zr/Hf; 0.23-1.45 Nb/Ta; 13.47-27.94 Sr/Eu) turn out to be much lower than those of chondrites proving that the crystallization of apatite proceeded from an enriched volatile melt.

The concentration of incompatible elements in apatite depends on complex compounds type. Bau & Dulski (1995) suggest that the complexation with fluorine is the major cause for Y/Ho > 28 values, whereas the complexation with bicarbonate is assumed to generate values < 28. In the investigated samples the second trend was found, suggesting that crystallization environment was not enriched in F and the fractionation of Y from HREE was limited.

The chondrite-normalized pattern of REE shows that the apatite is enriched in HREE (Fig. 1) with a narrow range of $(Ce/Yb)_{CN}$ values (0.88-0.90). Generally, the REE fractionation pattern of pegmatite apatite displays HREE enrichment, with $(Ce/Yb)_{CN}$ varying from 0.2 to 0.9 (Fig. 1; Belousova *et al.*, 2002). The shape of the curves, characterized by convex segments, infers the presence of the tetrad effect. The calculated values of the tetrad effect ($TE_{1,3}$) based on Irber's model vary between 1.12-1.15. The fractionation trend of the REE and the presence of the tetrad effect found for our samples are typical of apatite from granite pegmatites derived from a residual melt of magmatic systems.

Fig.1. Chondrite-normalized pattern of REE for apatite from Răzoare pegmatites (2-4) and for pegmatite apatite in general (1; Belousova *et al.*, 2002).

Conclusions

- The deviation of the Răzoare pegmatite apatite from the apatite ideal composition is a result of the accommodation of Mn, Na and Fe in Ca-sites.
- The concentrations of Mn, Na and Fe vary in a narrow range and are typical of apatites from granite pegmatites associated with magmatic granite.
- Among the both accessory and rock-forming minerals, the apatite represents the main host of Y, REE, U, and Th.
- The values of Zr/Hf, Nb/Ta and Sr/Eu ratios, the REE fractionation pattern and the presence of tetrad effect denote that crystallization of apatite proceeded from volatile-enriched melt. The value of Y/Ho ratio found for investigated samples suggests that the fluorine was not the main volatile component in the pegmatite melt. Based on previous studies (Macovei & Pomârleanu, 1980; Mârza *et al.*, 1986), the fluorine concentration level was suitable to give rise to fluorine-apatite.
- The composition of apatite from Răzoare pegmatites suggests the crystallization of these rocks from a granitic melt, which could result from partial melting of the metamorphic host-rocks. The anatectic origin of the pegmatites was formulated by Mârza (1985) and it is supported by P-T data obtained by Radu (1997) for the metamorphic host-rocks.

References

- Balintoni, I. (1996) The geotectonics of the metamorphic terrains from Romania (in Romanian). *Ed. Carpatica*, Cluj-Napoca, 176 p.
- Bau, M. (1996) Controls on the fractionation of isovalent trace elements in magmatic and aqueous system: evidence from Y/Ho and Zr/Hf and tetrad effect. *Contrib. Mineral. Petrol.*, **123**, 323-333.
- Bau, M. (1997) The tetrad effect in highly evolved felsic igneous rocks – a reply to the comment by Y.Pan. *Contrib. Mineral. Petrol.*, **128**, 409-412.
- Bau, M., Dulski, P. (1995) Comparative study of yttrium and rare-earth elements behaviours in fluorine-rich hydrothermal fluids. *Contrib. Mineral. Petrol.*, **119**, 213-223.
- Belousova, E.A., Griffin, W.L., O'reilly, S.Y., Fisher, N.I. (2002) Apatite as an indicator mineral for mineral exploration: trace-element compositions and their relationship to host rock type. *J. Geochem. Explor.*, **76**, 45-69.
- Dulski, P. (2000) Reference materials for geochemical studies: New analytical data by ICP-MS and critical discussion of reference values. *Geostandards Newsletter*, **25**, 87-125.
- Govindaraju, K. (1994) Compilation of working values and sample description for 383 Geostandards. *Geostandards Newsletter*, **18**, Special Issue, 1-158.
- Gramaccioli, C.M., Pezzotta, F. (2000) – Geochemistry of yttrium with respect to rare-earth elements in pegmatites. *Mem Soc. Italiana di Sci. Nat. e Museo Civico di Storia Nat. di Milano*, **XXX**, 111-115.
- Hughes, J.M., Cameron, M., And Crowley, K.D. (1991a) Ordering of divalent cations in the apatite structure: crystal structure refinements of natural Mn- and Sr - bearing apatite. *Amer. Mineral.*, **76**, 1857–1862.
- Hughes, J.M., Cameron, M., And Mariano, A.N. (1991b) Rare-earth-element ordering and structural variations in natural rare-earth-bearing apatites. *Amer. Mineral.*, **76**, 1165–1173.
- Hughes, J.M., Rakovan, J. (2002) The crystal structure of apatite, $\text{Ca}_5(\text{PO}_4)_3(\text{F}, \text{OH}, \text{Cl})$. In Phosphates. Geochemical, Geobiological and Material Importance (M.J. Kohn, J. Rakovan & J.M. Hughes Eds.) *Mineral. Soc. Amer., Geochim. Soc.*, **48**, 1-11.
- Irber, W (1999) The lanthanide tetrad effect and its correlation with K/Rb, Eu/Eu*, Sr/Eu, Y/Ho, and Zr/Hf of evolving peraluminous granite suites. *Geochim. Cosmochim. Acta*, **63**, 489–508.
- Macovei, V., Pomârleanu, V. (1980) The study of IR spectra of the apatites from Romania. *Stud. Cerc. Geol. Geofiz. Geograf., ser. Geol.*, **25**, 51-54 (In Romanian).
- Mârza, I. (1985) The genesis of magmatic ore deposits (in Romanian). Vol. II: Ortomagmatic and pegmatitic metallogeny (in Romanian). *Ed. Dacia*, 331 p.
- Mârza, I., Znamirovski, V., Moşiu, A. (1986) The pegmatite apatite from Răzoare (Maramureş District) – The investigations based on X-Ray diffraction, neutron activation and microprobe analyses (in Romanian). *Stud. Cerc. Geol. Geofiz. Geograf., Ser. Geol.*, **31**, 13-17.
- Monecke, T. Kempe, U., Monecke, J., Sala, M., Wolf, D (2002) Tetrad effect in rare earth element distribution patterns: A method of quantification with application to rock and mineral samples from granite-related rare metal deposits. *Geochim. Cosmochim. Acta*, **66**, 1185–1196.
- Piccoli, P.M., Candela, P.A. (2002) Apatite in igneous systems. In vol.: Phosphates. Geochemical, Geobiological and Material Importance (M.J. Kohn, J.Rakovan & J.M. Hughes Eds.) *Mineral. Soc. Amer., Geochim. Soc.*, **48**, 255-287.
- Radu, D.M. (1997) Geothermometry of metapelitic rocks from the western part of Preluca crystalline complex, Maramureş. *Rom. J. Mineral.*, **78**, 21-30.
- Rădăşanu, S. (2002) The geochemistry and geothermometry of the pegmatites from Preluca Mountains (in Romanian). *Ph.D. Thesis*, “Al.I.Cuza” University from Iaşi, 172 p.
- Reed, S.J.B. (1996) Electron microprobe analysis and scanning electron microscopy in geology. *Cambridge University Press*, 215 p.
- Ronsbo, J.G. (1989) Coupled substitutions involving REEs and Na and Si in apatite in alkaline rocks from the Ilimaussaq intrusion, South Greenland, and the petrological implications. *Amer. Miner.*, **74**, 896-901.
- Rusu, A., Balintoni, I., Bombiţă, G., Popescu, G. (1981) Geological map of Romania (Preluca File), sc.1:50.000. *Geol. Inst. of Romania*, Bucharest.
- Sha, L-H., Chappell, B. (1999) Apatite chemical composition, determined by electron microprobe and laser-ablation inductively coupled plasma mass spectrometry, as a probe into granite petrogenesis. *Geochim. Cosmochim. Acta*, **63**, 3861-3881.
- Zhang, H., Liu, C.Q. (2001) Sr/Eu ratio in apatite as a recorder of fluid exsolution from pegmatite-forming melts. *11th Annual V.M. Goldschmidt Conference*, Abstr. vol., p. 3314.

THE ONTOGENY OF SPELEOTHEMS

Charles Self⁶²

Speleothems are secondary mineral deposits whose growth in caves can be studied by mineralogical techniques. One of these techniques is the ontogeny of minerals, which is the study of individual crystals and their aggregates as physical bodies rather than as mineral species. Ontogeny of minerals as a scientific subject has been developed in Russia but is poorly understood in the West. In this lecture, I will introduce the basic principles of this subject and explain a hierarchy scheme whereby mineral bodies can be studied as crystal individuals, aggregates of individuals, associations of aggregates (termed koras), and as sequences of koras (ensembles). Selenite needles are crystal individuals, most other speleothems are aggregates, while the association (kora) of calcite stalactites and stalagmites is known even to members of the public. Most cavers understand that crystallization in caves is a cyclic process, the product of any one cycle being termed (in ontogeny) an ensemble.

Individuals, aggregates, koras and ensembles are classed as "minor mineral bodies" because they can be studied by mineralogical rather than by petrographic techniques.

Ontogeny of minerals is not simply a new classification system for mineral bodies, it is a method by which past crystallization environments can be interpreted. The structure and texture of minor mineral bodies can be directly related to environmental factors at the time of their development. Speleothems are ideal subjects for this type of study, since there are few common mineral species in caves, yet there is a great variety in the forms that these minerals can take.

⁶² Bristol Speleological Society, Bristol, UK.

SPECTROSCOPIC INVESTIGATION OF SOME OBSIDIAN ARCHAEOLOGICAL ARTIFACTS

Viorica Simon⁶³, Corina Ionescu⁶⁴, Liviu Dărăban¹

Introduction

Obsidian is a naturally occurring volcanic glass, of high interest both to geologists, archaeologists and physicists (Earle & Ericson, 1977; Hughes, 1988). Geologists and physicists are interested in the chemical and respectively the physical properties of obsidian, while archaeologists are primarily interested in its cultural features.

Because of its workability and capacity to form sharp edges during the splitting, obsidian was preferred by most of the pre-metal ancient cultures for obtaining tools and weapons, at the same time being the object of continuous commercial exchanges. Elaborate networks sometimes carried away the obsidian raw material as well as the artifacts into the areas where it was not naturally available and where it attained a position of high value.

The ability to identify the geological sources of the archaeological obsidian artifacts provides a key to the reconstruction of extinct systems of contact, trade and population movements. Different patterns of obsidian distribution in space and time may also offer information on the social structures of the prehistoric peoples.

A number of different methods have been developed to "fingerprint" or characterize obsidian in correlation with the geological sources. For example, obsidian artifacts from the Mediterranean islands have been recently investigated (Villeneuve *et al.*, 2002) using electron paramagnetic resonance of Fe³⁺ ions, thus revealing that in the Aegean area the main natural source of obsidian was Melos Island. In Anatolia the main obsidian sources were the Agicol and Ciftlik locations.

It is well-known that the main Central European obsidian sources were the Viničky (Zemplén Mts., East-Slovakia) and Eperjes (Tokaj Mts., North-Hungary) occurrences (Thorpe *et al.*, 1984), related to Neogene volcanics. The obsidian from Slovakia, known as Carpathian 1-type, is grey-black or black in colour and slightly translucent. The Hungarian obsidian, of Carpathian 2-type has black colour and is almost opaque.

The present study focusses on the iron content (obtained by electron paramagnetic resonance and X-ray fluorescence) of archaeological obsidian artifacts discovered in the sites from the north of Romania as well as some obsidian artifacts of Slovakian origin. The resulted data have been compared, with the aim to identify the provenance sources of Romanian obsidian artifacts.

Samples and methods

The investigated samples have been provided by the National History Museum of Transylvania (Cluj-Napoca, Romania). They consist in fragments of Neolithic obsidian artifacts (tools and weapons) originated from the archaeological sites of North-Romania (Piatra Curmeni and Oraşul Nou, both in Oaş area) and from the eastern Slovakian sites (Zemplén Mts.).

Electron paramagnetic resonance (EPR) measurements have been performed at room temperature, in X band (9.4 GHz) by means of a standard JEOL-JES-3B spectrometer with 100 kHz field modulation.

X-ray fluorescence spectra have been recorded with a SPARK instrument equipped with LiF analyser. The Roentgen tube of the instrument is provided with copper anti-cathode.

⁶³ "Babeş-Bolyai" University, Faculty of Physics, 1, Kogălniceanu Str., RO-3400 Cluj-Napoca, Romania. E-mail: <viosimon@phys.ubbcluj.ro>.

⁶⁴ "Babeş-Bolyai" University of Cluj-Napoca, 1, Kogălniceanu Str., Dept. of Mineralogy, RO-3400 Cluj-Napoca, Romania. E-mail: <corina@bioge.ubbcluj.ro>.

Results and discussion

The complex EPR spectra recorded from the obsidian patterns are mainly due to iron in different states and site location. Sometimes to these spectra can also contribute other paramagnetic ions occurring in obsidian. For all the samples investigated in this study, the EPR spectra (Fig. 1) consist in two resonance lines, a weak narrow line with $g_{\text{eff}} = 4.3$ arising from Fe^{3+} isolated ions, at a magnetic field of 1600 G, and a broad line having the line-width around 2000 G, characterized by $g_{\text{eff}} = 2.0$, which may be considered as an envelope over the signals arising from all the resonant centres contained in the sample⁶⁵. The amplification degrees used to record the EPR spectra presented in Fig. 1 are different for the samples (a), (b) and (c) and they are in the relation 10:7:2. As may be noticed from the resonance spectra, in correlation with the intensity of the large lines with $g_{\text{eff}} = 2$, the obsidian sample found in Piatra Curmeni (Fig. 1a) contains few paramagnetic centres compared with those in the other samples. One can also notice that the number of the isolated Fe^{3+} ions related to the number of all paramagnetic species, in each sample, decreases in the same order, from the Piatra Curmeni obsidian followed by the Slovakian obsidian and finally by the Oraşul Nou one.

Fig. 1. EPR spectra of obsidian samples from Piatra Curmeni -Romania (a), Slovakia (b) and Oraşul Nou - Romania (c).

⁶⁵ The possibility to discriminate the Fe^{3+} ions among the other paramagnetic species indicates that obsidian can be also subject of magnetic investigations.

Villeneuve *et al.* (2002) consider that condensed clusters of Fe^{3+} ions give rise to a resonance line at $g_{\text{eff}} = 2.0$. In addition to the signals with $g_{\text{eff}} = 2.0$, 2.2 and 4.3 they also detected broad resonance lines that move towards weaker fields when temperature decreased (from 300 to 5 K), and in contrast with the first ones, whose position and width do not depend on temperature, their width can either increase or decrease. They attributed these signals to microcrystals of iron oxides (hematite, magnetite, pseudobrookite) or silicates, present as inclusions in the amorphous groundmass of obsidian. The contribution of every signal to EPR spectrum is different for each source, according to the different thermodynamic conditions that occurred during obsidian formation. Thus, it is possible to discriminate *a priori* one geological source from another using EPR spectra.

X-ray fluorescence analysis brings also important information on obsidian patterns (Kowalski *et al.*, 1972, Biro *et al.*, 1986). The X-ray fluorescence spectra for three of the investigated samples are presented in Fig. 2. The obsidian sample found in the Piatra Curmeni site (Fig. 2a) has a low iron content if compared with Slovakia (Fig. 2b) and Oraşul Nou (Fig. 2c) samples.

Fig. 2. X-Ray fluorescence spectra of obsidian artifacts from Piatra Curmeni - Romania (a), Slovakia (b) and Oraşul Nou - Romania(c).

Biro *et al.* (1986) already evidenced slight differences both in the main components as well as in the trace elements between Slovakian and Hungarian obsidian occurrences. In agreement with these data, our results suggest the idea that the Neolithic obsidian samples from both sites (Piatra Curmeni and Oraşul Nou) originate from the Carpathian group (Slovakian-Hungarian occurrences). The Piatra Curmeni obsidian artifacts reveal closer resemblance with the Slovakian obsidian (Carpathian 1-type), while the Oraşul Nou sample is slightly different, suggesting the provenance from the Hungarian occurrences (Carpathian 2-type).

Conclusions

The iron content evidenced by X-ray fluorescence in the Slovakian obsidian is higher than that of the obsidian sample found in Piatra Curmeni but less than for the obsidian sample found in Oraşul Nou.

In the same relation is the concentration of all the other paramagnetic elements, which contribute to the large EPR line.

The EPR and X-Ray fluorescence results suggest an idea, that the investigated patterns are originate from the Carpathian obsidian group: the sample found in Piatra Curmeni from Carpathian 1-type (Slovakian) while the sample found in Oraşul Nou was probably brought from Carpathian 2- type (Hungarian) obsidian sources.

They denote trade relations along the rivers from this region, as the obsidian artifacts were found at relatively long distances from the sources.

References

- Earle, T.K., Ericson, J.E. (1977) Exchange Systems in Archaeological Perspective, In Earle, T.K. & and Ericson, J.E. (Eds.) Exchange Systems in Prehistory. *Academic Press*, New York, 3-12.
- Hughes, R.E. (1988) The Coso Volcanic Field Reexamined: Implications for obsidian and hydration dating Research, *Geoarchaeology*, **3**, 253.
- Villeneuve, G., Duttine, M., Poupeau, G., Rossi, A., Scorzelli, R.B. (2002) Electron Paramagnetic Resonance of Fe^{3+} ion in obsidians from Mediterranean islands. Application to provenance studies, *IV-th Conference on Natural Glasses*, Lyon -France, August 2002.
- Thorpe, O.W., Warren, S.E., Nandris, J.G. (1984) The Distribution and Provenance of Archaeological Obsidian in Central and Eastern Europe, *J. Archaeol. Science*, **11**, 183.
- Kowalski, B.R., Schatzki, T.F., Stross, F.H. (1972) Classification of archaeological artifacts by applying pattern recognition techniques to trace element data, *Anal. Chem.*, **44**, 2176.
- K.T. Biro, I. Posgai, A. Vadar (1986) Prehistoric Flint Mining and Lithic Raw Material Identification in the Carpathian Basin, *Acta Archaeol. Acad. Scient. Hungaricae*, **38**, 17.

PILLOW-LAVAS OF THE POVLEN MT. (WESTERN SERBIA)

Danica Sreckovic-Batocanin⁶⁶, Biljana Nikolin², Nada Vaskovic⁶⁷

Introduction

In the western ridge of the Povlen Mountain one of the largest outcrops of basic volcanic rocks (covering about 4 km²), occurring as pillow lavas, has been observed (Fig. 1). In the lower parts of this unit, epiclastic breccias composed of serpentinite, diabases of ophitic, intersertal or quench texture as well as glassy lava clasts, prevail. Pillow lavas of different size (from 10 x 15 cm up to 20 x 25 cm) and heaving a weakly developed amygdaloidal structure, progressively become more abundant upwards. They are dark-green in colour, highly hydrothermally altered and cataclastically deformed. In the marginal parts of this unit, tuffaceous sandstone with well developed, clearly visible granulometric gradation overlain by Triassic limestones was noticed.

Fig.1. Pillow-lavas of the Povlen Mt.

These lavas were identified as diabases according to Mojsilovic *et al.* (1966; 1975). According to their geochemical similarities with a nearby situated ophiolite complex, they were considered as the highest part of the oceanic crust (Sreckovic-Batocanin, 2001). They were tectonically separated from the lower parts of the ophiolitic complex, that are exposed some 8 km further south.

⁶⁶ Faculty of Mining and Geology, Djusina 7, 11 000 Belgrade, Serbia. E-mail: <danicasreckovic@yahoo.com>.

⁶⁷ Faculty of Mining and Geology, Djusina 7, 11 000 Belgrade, Serbia. E-mail: <nadavask@mkpg.rgf.bg.ac.yu>.

Analytical techniques

The samples were examined in thin sections for the identification of the weathering processes and the intensity of the phenocrysts accumulation.

The major and trace elements of all samples were determined by XRF techniques (PHILIPS PW 2400) in the Mineralogical Institute in Köln (Germany). Samples were treated with Li-tetra borates before to be analysed. The estimation of the relative precision is based on international standards.

The mineral composition was determined on a JOEL 740 SEM microprobe at the Aristotle University of Thessaloniki (Greece), having the following operating parameters: 10 s counting time, 20 kV accelerating voltage, 15 nA beam current and ca. 2 μm impact diameter. Natural silicates and oxides were used as standards.

Geological setting

The investigated rocks occur in the so-called »Mesozoic zone« of Petkovic (1930-1931). According to Mojsilovic *et al.* (1966, 1975), within this zone, except some large blocks, different sedimentary, igneous and metamorphic rocks were considered members of the »Diabase-chert formation«, until Dimitrijevic & Dimitrijevic (1974) explained it as an ophiolitic melange. This area is a part of the western subzone of Vardar Zone Composite Terrane (Karamata & Krstic, 1996) and it is composed of a dismembered ophiolite complex, Triassic, Liassic and Cretaceous carbonate rocks, and, less abundant, of sandstone blocks, embedded in a clayey-silty matrix. It represents the relic of the marginal basin, existing at least from the Late Triassic to Late Cretaceous and located between the Kopaonik block and the Drina-Ivanjica unit.

Mineral and chemical composition

As these lavas have a similar mineral composition (plagioclase with An_{22} in average, augite, ore minerals as accessories and chlorite, calcite, quartz, epidote-zoisite, prehnite, actinolite as secondary minerals), they were separated mainly according to their texture and mineralogy, in:

1. Ophitic metabasaltic andesites (with a glomeroporphyric texture)
2. Ophitic quartz-metabasaltic andesites (>10% quartz; ophitic texture)
3. Porphyritic metabasaltic andesites.

These rocks are highly altered, and their geochemical values (Tables 1, 2) clearly indicate basaltic andesites with OFB signature (Fig. 2). The low content in refractory elements is probably the result of their concentration in the lower members of the ophiolite complex.

Fig. 2. Plot of the Povlen Mt. pillow lavas on the Zr-Ti/100–Sr/2 diagram (Pearce & Cann, 1973). Abbreviations: OFB – ocean floor basalts; IAB – island arc basalts; CAB – calcalkaline basalts.

Table 1.

Representative major element analyses of the Povlen Mt. pillow lavas.

Wt%	DB-6	DB-8	DB-13
SiO ₂	54.90	55.45	55.71
TiO ₂	1.18	1.15	1.44
Al ₂ O ₃	14.23	13.78	13.71
Fe ₂ O ₃	2.08	1.09	3.83
FeO	7.50	9.00	7.00
MnO	0.14	0.14	0.13
MgO	5.32	5.95	4.56
CaO	6.64	5.93	5.00
Na ₂ O	4.95	4.47	4.56
K ₂ O	0.08	0.24	0.05
P ₂ O ₅	0.12	0.11	0.14
SO ₃	0.03	0.04	0.03
LOI	3.48	3.64	3.40
Σ	100.73	100.08	99.65
CIPW norm			
Q	2.84	2.62	10.13
Or	0.47	1.42	0.30
Ab	41.89	37.82	38.59
An	16.37	16.83	16.79
Di	12.94	9.85	5.79
Hy	16.89	25.76	15.74
Ol	0.00	0.00	0.00
Mt	3.22	1.12	5.74
Il	2.24	2.18	2.73
Ap	0.28	0.25	0.32
%An	28.00	31.00	30.00

Table 2.

Representative trace element analyses of the Povlen Mt. pillow lavas

Ppm	DB-6	DB-8	DB-13
Sc	38	37	36
V	273	261	319
Cr	70	87	15
Co	38	38	33
Ni	42	52	21
Cu	44	39	22
Zn	69	78	74
Ga	8	8	11
As	10	2	2
Rb	6	0	5
Sr	85	121	108
Y	35	32	38
Zr	116	110	127
Nb	4	4	5
Mo	12	12	12
Cs	7	16	3
Ba	30	17	16
Ce	12	17	3
Nd	18	19	10
W	6	6	0
Pb	4	9	6
Th	8	5	5
U	5	5	4

Conclusions

Pillow-lavas of the Povlen Mt. represent the uppermost member of the ophiolite complex whose lower part is exposed some 8 kms southward. They were cooled rapidly at low temperature

conditions, giving rise to skeletal crystals of oxide minerals, fan-shaped epidote, plagioclase with quenched texture, as well as to the presence of volcanic glass. The low pressure conditions (acc. to Mahood & Baker, 1986) are reflected in the pyroxene chemistry - low Al content in M1 site (0.003-0.028) and the presence of vacuoles (up to 5 mm in diameter) subsequently filled by calcite, quartz, prehnite and chlorite. After their consolidation, these rocks were metamorphosed at the lower part within the prehnite-actinolite facies (temperature around 250-300°C and pressure up to 2 kbars), in the presence of fluids whose composition changed from CO₂-poor (prehnite formation), to CO₂-rich (calcite formation). These metabasaltic andesites have a subalkaline character, with low K content (0.05-0.24%) and high Y/Nb ratio.

The amygdaloidal structure and the absence of cherts should indicate their genesis in a shallow-water environment. The presence of breccias, according to Nicolas *et al.* (1989) suggest the pillow lavas formation along a paleo-transform fault, where the breccias represented relic screens over the basaltic floor.

Acknowledgements

The authors are indebted to Dr. Almut Katzemich from the Mineralogical Institute in Köln (Germany) who performed XRF-analyses and to professor George Christofides from Aristotle University (Thessaloniki, Greece) who performed microprobe analyses. Finally, we would like to acknowledge the thoughtful review by Academician Stevan Karamata.

References

- Dimitrijevic, M.D., Dimitrijevic, M.N. (1974) On genesis of "Diabase-chert formation". *Geol. Glasnik*, **7**, 333-350, Titograd.
- Karamata, S., Krstic, B. (1996) Terranes of Serbia and neighbouring areas. In *Terranes of Serbia* (Eds. Knezevic, V. & Krstic, B.), 25-40, Belgrade.
- Mahood, G.A., Baker, D.R. (1986) Experimental constraints on depths of reaction of mildly alkalic basalts and associated felsic rocks. *Contrib. Min. Petrol.*, **93**, 251-264.
- Mojsilovic S., Filipovic I., Baklajic Dj., Djokovic I., Navala M. (1966) Explanatory text for the Basic Geol. Map of SFRJ, sheet Valjevo, L-34-136, 1:100.000, SGZ, Beograd (In Serbian).
- Mojsilovic S., Filipovic I., Avramovic V., Pejovic D., Tomic R., Baklajic D., Djokovic I., Navala M. (1975) Basic geological Map of SFRJ, sheet Valjevo L-34-136, 1:100.000, Izdanje SGZ, Beograd.
- Nicolas, A., Ceuleneer, G., Boudier, F., Misseri, M. (1988) A structural mapping in the Oman ophiolites: mantle diapirism along an ocean ridge. *Tectonophysics*, **151**, 27-56.
- Petkovic K. (1930-1931) Geological Map of the Yugoslavian Monarchy, 1: 1.000.000. Geol. Inst. Univ. Beograd & Geol. Inst. Zagreb (Ed. Libraire Francois Bach) Beograd.
- Pearce, J.A., Cann, J.R. (1973) Tectonic setting of basic volcanic rocks determined using trace elements analyses. *Earth Planet Sci. Lett.*, **19**, 290-300.
- Sreckovic-Batocanin, D. (2001) Petrology of the Tejci ophiolite complex (Western Serbia) – *Ph.D. Thesis* (unpublished), 167 p., Belgrade.

SOME GEOCHEMICAL REMARKS ON BASALTS FROM BĂIȚA BIHOR (BIHOR MTS., ROMANIA). MAJOR ELEMENTS

Dan Stumbea⁶⁸

Geographically, Băița Bihor is located in the southwestern area of Bihor Mts., a subdivision of Northern Apuseni Mountains. The geological setting is controlled by a very complicated tectonic situation, in which Bihor, Codru and Arieșeni structural units are involved; the lithology of these units consists of limestones (the Bihor unit), dolomites and limestones (the Codru unit), clays and sandstones (the Arieșeni unit). This structure is penetrated, during four magmatic stages (Berbeleac, 1988), by banatitic intrusives (Upper Cretaceous-Lower Paleogene); the largest of it being a batholith made up of granites, granodiorites and pegmatites. The batholith belongs to the third magmatic stage and exhumed in the Valea Seacă area.

The present paper deals with some basic dykes of the fourth magmatic stage, named previously "basalts" by Stoici (1983) and showing no relationship with the felsic batholith. Based on the chemical data published by the quoted author, the study focuses on a more detailed geochemical characterization of the basic dykes. The chemical data concern the major elements and were performed using the wet chemical technique.

The plot of 65 samples supposed to be basalts, shows a wide range of samples distribution in TAS diagram (SiO_2 vs. $\text{Na}_2\text{O}+\text{K}_2\text{O}$). Thus, the samples plot in the fields of basalts, trachybasalts, tephrites, basaltic andesites, basaltic trachyandesites, trachyandesites and finally andesites. De La Roche-Leterier diagram, based on the standard formula parameters of Si, Na, K, Fe, Ti (R1) versus Ca, Mg, Al (R2) leads to similar results.

In order to find out whether this conclusion is confirmed, a statistical processing of all 65 analyses has been performed considering their major elements. This approach revealed two maximums in the distribution of SiO_2 contents: at about 48.7 wt% and at 52.7 wt% (Fig. 1). Based on this fact and knowing that SiO_2 has generally a strong bimodal character in magmatic rocks (it means two maximums - at about 52.2 wt% and at 73.0 wt% respectively) we assumed that in the case of samples from Băița Bihor there are two populations of data: a basaltic population (with $\text{SiO}_2 < 50.7$ wt%) and an andesitic one (with $\text{SiO}_2 > 50.7$ wt%).

Our attempt to establish similar limits by means of a statistical approach performed on data concerning other major elements has generally failed; except for MgO , which revealed three populations of data, the distribution of all major elements shows both symmetrical and asymmetrical gaussian patterns.

The replot of basalt data in both TAS and De La Roche-Leterier diagrams confirms that the limit of 50.7wt% SiO_2 statistically identified by our research is quite realistic. Our further study focuses only on basalt samples (28 analysis).

The CIPW norm sustains the assumption that the dykes from Băița Bihor belong to both basalt and tephrite categories. Thus, the normative mineralogical composition of tephrites shows a lower content of normative quartz, plagioclases and pyroxenes, as well as a higher one concerning normative orthoclase, feldspathoids and olivine as compared with basalts. In terms of normative olivine, its amount of about 14% in tephrites (in contrast with only 5% in basalts) suggests that at least some of the tephrites belong to the basanite category.

The distribution of data in both $\text{TiO}_2\text{-K}_2\text{O-P}_2\text{O}_5$ (Pearce *et al.*, 1975) and $\text{FeO-(Na}_2\text{O+K}_2\text{O)-MgO}$ diagrams (Irvine & Baragar, 1971) suggests that the basalts of the Băița Bihorului area belong to a calc-alkaline series and confirms the earlier opinion of Berbeleac (1988) who associates the banatites of Băița Bihorului to a magmatic arc tectonic setting.

⁶⁸ University „Al .I. Cuza”, 20A Carol I Bvd., 6600, Iași, Romania. E-mail: <dstumbea@geo.uaic.ro>.

Fig. 1. The frequency vs. SiO₂ diagram shows maximus at 48.7wt% (middle of 47.7-49.7wt% range) and at 52.7wt% (middle of 51.7-53.7wt% range). The limit between the two populations of data has been theoretically drawn at 50.7wt% (middle of 49.7-51.7wt% range).

References

- *** (1981) Basaltic volcanism study project. 1981. Basaltic Volcanism on the Terrestrial Planets. *Pergamon Press, Inc., New York*, 1286 p.
- Berbeleac, I. (1988) Ore deposits and global tectonics (in Romanian). *Ed. Tehn. Bucureşti*, 327 p.
- Stoici, S.D. (1983) The metallogenetic district of Băiţa Bihorului (in Romanian). *Ed. Acad., Bucureşti*, 190 p.
- Pearce, T.H., Gorman, B.E., Birkett, T.C. (1975) The TiO₂-K₂O-P₂O₅ diagram: A method of discriminating between oceanic and non-oceanic basalts. *Earth and Planetary Science Letters*, **24**, 419-426.
- Irvine, T.N., Baragar, W.R.A. (1971) A guide to the chemical classification of the common volcanic rocks. *Can. J. Earth Sci.*, **8**, 523-548.

THE METAMORPHIC BASEMENT OF THE ROMANIAN CARPATHIANS: A DISCUSSION IN THE LIGHT OF RADIOGENIC K-Ar AND $^{40}\text{Ar}/^{39}\text{Ar}$ DATING

Carol Strutinski⁶⁹, Adrian Puşte⁷⁰, Rodica Stan⁷¹

Over 600 K-Ar and $^{40}\text{Ar}/^{39}\text{Ar}$ ages published during the last four decades on rocks of the metamorphic basement from the Romanian Carpathians, the Transylvanian Basin and the eastern part of the Pannonian Basin are statistically treated and discussed. The radiometric ages have been assigned to four groups, designated as: pre-Hercynian (older than 375 Ma), Hercynian (375-251 Ma), „Kimmerian“ (250-121 Ma) and eo-Alpine (younger than 120 Ma). They have been further sorted by the type of mineral fraction on which they were performed and plotted by distinct symbols on a map having the principal tectonic lines as a background. For the sake of convenience, different individual data coming from the same area/unit and having the same or similar characteristics (age and processed mineral) have been put together and represented on the map by only one symbol.

The principal features to be noticed on the distribution map are as follows:

1. East Carpathians. The age distribution shows a by and large SW-NE polarity, with the oldest, mainly Hercynian, ages in the NE, close to the boundary with the sedimentary units of the Outer Dacides, and the youngest Eo-Alpine ages in the SW, in the Rodna massif.

2. South Carpathians. In the Lower Danubian Units an unusual large number of pre-Hercynian ages points to the preservation of a Cadomian or Caledonian province through Hercynian and Alpine times, while in the Upper Danubian, Getic and Supragetic units the Hercynian and „Kimmerian“ ages prevail. Eo-Alpine ages are rather rare and are usually confined to the innermost Danubian zones and to some shear planes.

3. Northern Apuseni Mountains (i.e. Inner Dacides). The Bihor Autochthonous shows an poorly-defined polarity in the distribution of ages, as compared to the East Carpathians. However, it may be regarded as the mirror image of the latter, as here the oldest (Hercynian) ages are situated to the west and south-west, while the younger (Eo-Alpine) ages occur in the east and north-east. The few radiometric ages obtained on rocks of the Biharia and Baia de Arieş units fall almost all in the eo-Alpine group. The situation is somewhat complicated in the Codru unit, located between the Bihor Autochthonous in the NW and the Biharia units in the SE, where a few analyses of amphibolitic rocks yielded Hercynian or even older ages.

4. The Transylvanian Depression. In the north-western part of the Transylvanian Depression (including the Preluca, Țicău and Codru “islands”) only three out of 31 samples indicate Upper Jurassic – Lower Cretaceous ages, the remaining 28 yielding Eo-Alpine ones. Three isolated age data obtained on drill cores located in the eastern and southern parts of the Transylvanian Depression are consistently older („Kimmerian“ or Hercynian), and correlate well with the corresponding ages from the adjacent inner parts of the East and South Carpathians.

5. Eastern part of the Pannonian Basin. Most ages are „Kimmerian“. However, in boreholes located between Oradea and Satu Mare some metamorphic basement rocks yielded eo-Alpine ages, that were lately reported also from the Hungarian side of the area (Arkai *et al.*, 1998).

⁶⁹ Postfach 10 28 20, 66028 Saarbrücken/Germany, E-mail: <carl_stru@yahoo.de>.

⁷⁰ “Babeş-Bolyai” University, Department of Mineralogy, 1, Kogălniceanu Str., RO-3400 Cluj-Napoca, Romania. E-mail: <adipuste@bioge.ubbcluj.ro>.

⁷¹ Geological Institute of Romania, POB 181, 3400 Cluj-Napoca/Romania.

The existing data strongly support the idea that within the Romanian tract of the Carpathians the most pervasive Alpine rejuvenation of basement rocks occurred in a belt of about 140-150 km width. Within this belt, the crystalline rocks were intensely reworked, undergoing a metamorphic event of Barrovian (medium PT) type before the final Austrian exhumation, evidenced by most of the Alpine ages. The eo-Alpine metamorphic belt is cropping out today on the flanks of the Mureş Zone, i.e. in the Rodna massif to the NE, and in the Northern Apuseni to the west. Away from it, the ages get progressively older and outline a broad Hercynian metamorphic province, having its thermal axis apparently displaced by only some tens of kilometers from the Alpine one. In the most external part of the South Carpathians a pre-Hercynian province could be outlined, obviously representing the north-western extension of the Moesian Plate. Within the Romanian Carpathians the radiometric K-Ar and $^{40}\text{Ar}/^{39}\text{Ar}$ ages, as well as newer fission-track data, do not sustain re-heating processes above 300°C and corresponding regional metamorphic events during meso- and neo-Alpine times (Dallmeyer *et al.*, 1998, 1999; Bojar *et al.*, 1998).

References

- Arkai, P., Berczi-Makk, A., Hajdu, D. (1998) Alpine prograde and retrograde metamorphisms in an overthrust part of the basement, Great Plain, Pannonian Basin, Eastern Hungary. *Acta Geol. Hungarica*, **41/2**: 179-210.
- Bojar, A.V., Neubauer, F., Fritz, H. (1998) Cretaceous to Cenozoic thermal evolution of the southwestern South Carpathians: evidence from fission-track thermochronology. *Tectonophysics*, **297**: 229-249.
- Dallmeyer, R.D., Neubauer, F., Fritz, H., Mocanu, V. (1998) Variscan vs. Alpine tectonothermal evolution of the Southern Carpathian orogen: constraints from $^{40}\text{Ar}/^{39}\text{Ar}$ ages. *Tectonophysics*, **290**: 111-135.
- Dallmeyer, R.D., Pană, D.I., Neubauer, F., Erdmer, P. (1999) Tectonothermal evolution of the Apuseni Mountains, Romania: Resolution of Variscan versus Alpine events with $^{40}\text{Ar}/^{39}\text{Ar}$ ages. *The Journal of Geology*, **107**: 329-352.

MINERAL CHEMISTRY OF THE PRIMARY MAGMATIC MINERAL ASSEMBLAGE OF THE „DEJ TUFF” (ROMANIA)

Alexandru Szakács⁷²

Introduction

The Middle Miocene rhyolitic Dej Tuff Complex (the „Dej Tuff”) consisting of volcanoclastic sandstones and conglomerates, medium to fine-grained tuffs, and alternating thin layers of tuffs, tuffites and tuffaceous marls, records the inception of the Neogene volcanic activity in Romania. It crops out along the north-western and northern margin of the Transylvanian Basin, as well as along its southeastern margin („Persani Tuff”). It was also found in many boreholes throughout the whole basin.

Former mineralogical investigations on the „Dej Tuff” focused mainly on its characteristic post-depositional diagenetic mineral assemblage. Its primary magmatic mineral assemblage is known only from classical petrographic studies. The aims of this study are 1) to identify the primary magmatic rock-forming mineral phases, either essential or accessory ones, 2) to distinguish them from accidental admixed phases, and 3) to investigate in detail, using modern analytical techniques, their mineral chemistry.

Mineral chemistry

Mineral chemistry of the major rock-forming minerals – feldspar, biotite, amphibole, pyroxene – as well as accessory phases – allanite, Fe-Ti oxides, etc. – have been systematically investigated through a large number of microprobe analyses performed with a JEOL-733 Superprobe at the University College (University of London).

Plagioclase crystalloclasts show a broad spectrum of An content, ranging from 9.6% to 70.7% with most of them containing 15-40% An. The An% distribution histogram (Fig. 1a) indicates two frequency maxima at 20-25% An and 30-40% An, respectively. The bimodal distribution of An% in plagioclase population may suggest magma mixing processes. An% values over 50% have mostly been found in plagioclase phenocrysts of dacitic to andesitic lithic clasts. Both normal and recurrent zoning of plagioclase have been recorded with An variation in individual crystalloclasts of 1.4-19.4%. Consistent sequential variation of An content of plagioclase has been pointed out at Cepari, where upward-decreasing An contents across three sequences are recorded.

K-feldspar is sporadically present in the „Dej Tuff”. Orthoclase crystalloclasts are mostly an admixed component in the volcanoclastics, but primary magmatic sanidine could be an accessory phase as it was recorded in a few outcrops (Fig. 1b).

According to the analytical data, **biotite** belongs to a transitional field between magnesian and ferrous biotites. High Fe^{3+} with respect Fe^{2+} , as well as lack of the correlation between Mg and Fe^{2+} , suggests various stages of alteration. TiO_2 content varies between 3.62-6.27%, most values being around <4%. These relatively high TiO_2 contents are not typical for rocks originating from crustal granitoidic magmas. Biotite chemistry strongly suggests that its composition has been influenced by the presence of cogenetic amphibole, pyroxene - and possibly olivine - in the melt (Fig. 2). According to the biotite systematics of Abdel-Rahman (1994), the „Dej Tuff” biotites belong to both calc-alkaline and alkaline types, the main trend being consistent with the calc-alkaline affinity.

⁷² Romanian Academy, Institute of Geodynamics "Sabba S. Stefanescu", 19-21, Jean-Louis Calderon Str., RO-70201, Bucharest, Romania. E-mail: szakacs@geodin.ro

Fig. 1. Mineral chemistry of feldspar: a - An% distribution histogram; b - Plagioclase and K-feldspar compositions in the An-Ab-Or diagram. Symbols represent samples originating from different localities.

Fig. 2. MgO-FeO-Al₂O₃ diagram for biotites (Speer, 1984). I: biotites associated with muscovite and topaz, II: biotite with no other minerals associated, III: biotite associated with amphibole, pyroxene and olivine. Symbols represent samples originating from different localities.

Amphiboles are present only in few outcrops of the “Dej Tuff” as an accessory component. They belong to the group of calcic amphiboles (hornblende), most crystalloclasts being magnesio-hornblende and magnesio-hastingsite. Few edenite grains have also been identified. Whether the calcic amphiboles are crystallized from the “Dej Tuff” magma or not, has been tested in several ways. Amphibole-biotite equilibrium test shows a relatively good negative correlation of the $\text{Fe}^{2+}/(\text{Fe}^{2+} + \text{Mg})$ ratio, as well as a positive correlation of Mn content, in coexisting biotite and hornblende for 3 out of 4 samples, whereas Ti shows no correlation trend. The amphibole-melt equilibrium has been tested by comparing amphibole composition with the composition of the volcanic glass. Amphibole/glass ratio shows values of 0.56 for Na and 0.34 for K which - provided $\text{Na}_2\text{O} > 3\%$ - is indicative for crystal-melt equilibrium. In the trivariate $\text{SiO}_2\text{-FeO+Fe}_2\text{O}_3\text{-Na}_2\text{O+K}_2\text{O}$ diagram of Wones & Gilbert (1982) the amphibole crystalloclasts of the “Dej Tuff” plot in the rhyolite-dacite field, whereas in the $\text{SiO}_2\text{-FeO-Fe}_2\text{O}_3$ diagram they plot in the andesite-basalt field. These data are consistent with the magmatic origin of the calcic amphiboles. Moreover, they suggest the provenance of the amphibole from a less differentiated part of a magma body.

Pyroxene is also a minor or accidental mineral phase in the “Dej Tuff”, encountered as two clinopyroxene species: augite and pigeonite (Fig. 4). Augite appears at well-determined levels in the volcanoclastic sequence at some occurrences (e.g. Cepari), while pigeonite occurs randomly at other localities (Bobâlna, Aluniş). Unlike pigeonite, an externally admixed component, augite - which participates with up to 4-5% within a specific level at Cepari, where biotite is lacking - shows chemical features, such as TiO_2 content, compatible with the chemistry of other mafic minerals present in the “Dej Tuff”, hence its crystallization from the same parental magma is likely. The primary magmatic clinopyroxene in the “Dej Tuff” at Cepari has been determined as a magnesium-rich augite.

Fig. 3. Plot of Dej-tuff amphiboles in the $\text{SiO}_2\text{-FeO-Fe}_2\text{O}_3$ and $\text{SiO}_2\text{-FeO+Fe}_2\text{O}_3\text{-Na}_2\text{O+K}_2\text{O}$ diagrams of Wones & Gilbert (1982).

Fig. 4. Plot of pyroxene compositions. The symbols represent samples originating from different localities.

Fe-Ti oxides are ubiquitous accessory minerals in the “Dej Tuff”. Their mineral chemistry allows the recognition of three groups: (1) a homogeneous group, formed by ilmenite, (2) a more heterogeneous group, with spinel-structure oxides, represented by titano-magnetite, and (3) accidentally occurring minerals, as two grains of rutile. The main part of the analyzed samples contain both ilmenite and titano-magnetite. Systematic compositional change of the Fe-Ti oxides has been pointed out at Cepari across compositional sequences, similar to the An% content variation of plagioclase. Fe-Ti oxide geothermometry yielded results consistent with these trends (Szakács, 2000b).

Allanite is also an ubiquitous accessory phase showing relatively homogeneous composition. Ce_2O_3 ranges between 11.53-13.23%, La_2O_3 between 7.08-8.37%, and Nd_2O_3 between 2.28-3.35%. Relatively low MnO (< 0.8%) and high TiO_2 (up to 2.4%) are characteristic for the “Dej Tuff” allanites, in concordance with the chemistry of other primary magmatic minerals. Compositional differences have been pointed out between allanite grains occurring at different levels in the lithological columns.

Fig.5 Variation in allanite composition according to the occurrence/position (1-6). Symbols represent samples originating from different localities.

Zircon and **apatite** are frequent accessory phases in the “Dej Tuff” but they have not been subjected to chemical investigation.

Monazite and **xenotime** are for the first time identified in the “Dej Tuff”. They represent sporadically occurring accessory minerals. Monazite appears as submillimetric subhedral grains, in places associated with apatite, which forms epitaxial overgrowths. Xenotime was encountered in only two grains, as epitaxial overgrowths on euhedral zircon crystals. The chemical compositions of six monazite and one xenotime grains are given in Table 1 and 2.

Table 1.

Microprobe analyses of monazite grains from the „Dej Tuff”(oxides in wt%).

Sample	Al ₂ O ₃	SiO ₂	P ₂ O ₅	CaO	TiO ₂	SrO	ZrO ₂	La ₂ O ₃	Ce ₂ O ₃	Nd ₂ O ₃	Sm ₂ O ₃	Total*
TD139-3A-mz1	0	0	32.70	0.47	0	0	0	15.52	26.63	12.38	1.52	89.22
TD166-6B-mz1	0.15	0.63	28.20	0.45	0.15	0.23	0.68	20.50	28.40	7.69	0.45	87.53
TD166-6B-mz2	0.12	1.00	30.13	0.63	0.1	0.37	0.63	20.57	29.52	7.83	0.60	91.61
TD166-8C-mz1	0	0	32.58	6.81	0.28	2.23	0	13.15	23.03	7.71	0.97	86.76
TD166-8C-mz2	0.11	0	33.12	1.53	0	0	0	14.85	25.95	10.3	1.07	86.93
TD166-8C-mz3	0.08	0	32.72	1.02	0.13	0.13	0	15.87	28.97	10.8	1.31	91.03

Table 2.

Microprobe analyses of a grain of xenotime from the „Dej Tuff” ”(oxides in wt%).

Sample	CaO	Y ₂ O ₃	Dy ₂ O ₃	Er ₂ O ₃	Yb ₂ O ₃	P ₂ O ₅	Total*
TD-166-5A-xn1	1.17	52.5	3.08	2.9	4.12	29.43	93.24

* Low Total % is due to the fact that Th has not been analyzed

Inclusion-relationships between accessory minerals allowed the reconstruction of the early crystallization history of the “Dej Tuff” magma: monazite – apatite – zircon – Fe-Ti oxides. Allanite has not been observed to have any inclusion-relationships with other accessory mineral(s).

Conclusions

The primary magmatic mineral components of the “Dej Tuff” include the main rock-forming minerals quartz, plagioclase and biotite, the minor components K-feldspar, amphibole and pyroxene, the ubiquitous accessory phases as Fe-Ti oxides, zircon, apatite, allanite, and the rare accessory minerals monazite and xenotime. The mineral chemistry of the mafic components strongly suggests that biotite, amphibole (magnesio-hornblende and magnesio-hastingsite) and clinopyroxene (Mg-rich augite) have crystallized from the same parental magma having relatively high Ti contents. The accessory mineral association includes REE-bearing minerals such as allanite, monazite and xenotime pointing to the REE-rich chemistry of the “Dej Tuff” magma. The primary mineral assemblage of the “Dej Tuff” might be a result of pre-eruptive differentiation in shallow magma chambers. Whole-rock major and trace element chemistry as well as Sr and Nd isotopic ratios (Szakács, 2000a) fully support such a hypothesis.

Acknowledgements

This work benefited by the logistic support for field-work and sampling provided by the Geological Institute of Romania. Microprobe analyses were performed at the University College (University of London) supported by the Royal Society. The professional and technical support provided by Hilary Downes and Andrew Beard during analytical work are highly appreciated. Corina Ionescu is thanked for improvement of the manuscript.

References

- Abdel-Rahman, A.F.M. (1994) Peraluminous magmas. *Journal of Petrology*, **35**, 2, p. 525-541.
- Szakács, A. (2000a) Petrological and tephrological study of the Lower Badenian volcanic tuffs from the north-western Transylvanian Basin. (in Romanian), PhD thesis, University of Bucharest.
- Szakács, A. (2000b) Inferring eruption sequences and magma chamber conditions from volcanogenic sediments: The Miocene Dej Tuff Complex in the Transylvanian Basin, Romania. IAVCEI General Assembly 2000, Bali, Indonesia, Abstracts, p. 131
- Speer, A. (1984) Micas in igneous rocks. in S.W. Bailey (ed). Micas, *Reviews in Mineralogy*, **13**, p. 299-356.
- Wones, D.R., Gilbert, M.C. (1982) Amphiboles in the igneous environment. in D.R. Veblen & P.H. Ribbe (eds): Amphiboles: petrology and experimental phase relations. *Reviews in Mineralogy*, vol. 9B, p. 355-390.

VOLBORTHITE, A HYDROUS COPPER-VANADATE FROM PHOSPHATE-BEARING ARGILLITES IN DÉDESTAPOLCSÁNY, UPPONY MTS. (N-HUNGARY)

Sandor Szakáll⁷³, I. Sajó⁷⁴

Secondary phosphates rarely have been found in the Palaeozoic argillites of the Uppony Mts. They occur in the fissure-fillings of brecciated or clastic argillite, mainly in near-surface positions in the surroundings of Dédestapolcsány and Nekézseny (Elsholtz *et al.*, 1974). The geochemical investigations of these argillites revealed some uranium enrichment, which shows good correlation with phosphorous. During the new mineralogical investigation of the Dédestapolcsány phosphate-rich area, a hydrous copper-vanadate, volborthite ($\text{Cu}_3\text{V}_2\text{O}_7(\text{OH})_2 \cdot 2\text{H}_2\text{O}$), was found for the first time in the Carpathians.

Volborthite found in the fissures shows yellow-brown, yellow-green, olive-green, and dark green colours. The common appearance includes: powdery scattering, micro-crusts, porous masses, globular and stalagmite-like aggregates (up to 1–3 mm). According to SEM studies these aggregates consist of 10–30 μm long oriented grown needles, which built up lamellae. The largest lamellae are up to 0.3–0.6 mm.

According to XRD patterns and the appearance of volborthite, we can distinguish at least two varieties, which crystallized differently. The yellowish varieties are the least crystallized, while the greenish varieties are the most crystallized phases. The *d* values show excellent agreement with JCPDS 26-1119 file, however, intensities – with exception of 100 reflexion – are smaller. Unit cell data are: *a* = 10.604 Å; *b* = 5.879 Å; *c* = 7.202 Å; β = 94.81°.

Three chemical analyses carried out with CAMECA CAMEBAX SX50 electron microprobe operated at 25 kV and 0.1 mA. Average of the three analytical results is as follows (in weight %): CuO: 40.97; V₂O₅: 25.14; P₂O₅: 0.64; Al₂O₃: 1.14; Cl: 0.41.

Accompanying minerals associated with volborthite are secondary, i.e. goethite, hematite, jarosite, Al phosphates (e.g. kingite, wavellite, evansite, vashegyite) and other phosphates (e.g. beraunite, probably turquoise). The occurrences of volborthite have been found mainly in the oxidation zone of sedimentary uranium deposits (Guillemin, 1956; Leonardsen & Petersen, 1974). The Dédestapolcsány volborthite is definitely connected with uranium-enrichment in the Uppony Mts. It is a near-surface precipitation, together with other secondary minerals, but secondary U-minerals have not been identified yet.

Volborthite sample from Dédestapolcsány is preserved in the mineral collection of the Herman Ottó Museum (Miskolc, Hungary), under catalogue number 24532.

References

- Elsholtz, L., Selmecziné, A.P., Selmeczi B. (1974) *Földt. Közl.*, **104**, 328–335.
Guillemin, C. (1956): *Bull. Soc. franc. Minéral. Cristallogr.*, **79**, 219–275.
Leonardsen, E.S., Petersen, O. V. (1974) *Amer. Mineral.*, **59**, 372–373.

⁷³ Department of Mineralogy and Petrology, University of Miskolc, Miskolc, Hungary. E-mail: askszs@gold.uni-miskolc.hu.

⁷⁴ Chemical Research Centre of the Hungarian Academy of Sciences, Budapest, Hungary.

TETRAHEDRITE – TENNANTITE FROM THE ROŞIA MONTANĂ GOLD-SILVER EPITHERMAL ORE DEPOSIT, APUSENI MOUNTAINS (ROMANIA)

Călin G. Tămaş⁷⁵, Dan Costin¹

The Roşia Montană area belongs to the Transilvanides geotectonic unit (Săndulescu, 1984, Balintoni, 1997). This area is located in the northern part of the so-called Golden Quadrangle from South Apuseni Mountains. The mining activity was always very important at Roşia Montană. Recently, the mining archaeological results have proved that the exploitation of gold and silver started before the Roman times (Cauuet *et al.*, 2003).

The Roşia Montană ore deposit was assigned to the low-sulfidation type by Mârza *et al.* (1997) and Tămaş (2002). Fluid inclusions study (Tămaş & Bailly, 1998, 1999; Tămaş, 2002) confirmed the low salinity (1-7.8 wt.% NaCl equiv.) and the low homogenization temperatures (198-340°C) of the mineralizing fluids from the Roşia Montană ore deposit.

The hydrothermal alterations are well developed in the Roşia Montană ore deposit (Tămaş, 2002). The most striking one is potassium metasomatism (with the formation of adularia). "Sericite" (muscovite) alteration is also widespread in the whole ore deposit. Intermediate and advanced argillic alteration is also occurring as overprint. High-grade ore zones are marked by intense silicification. The carbonate-type alteration is present in very restricted areas.

According to Hedenquist's (2003) reinterpretation of the epithermal systems classification, the Roşia Montană ore deposit may be assigned to an intermediate sulfidation type.

At Roşia Montană the ore is hosted mainly by the breccia-pipe bodies, and at a lesser extent by the dacitic breccias occurring near the volcanic vent, or in the Neogene and Cretaceous sediments. The ore bodies are represented by phreatomagmatic and phreatic breccia structures, veins, stockworks, and impregnations. Placers and paleoplacers with gold content were also reported.

The Roşia Montană ore deposit is centered on the Cetate phreatomagmatic breccia-pipe body. This breccia structure was examined by Tămaş (2002) who interpreted the mechanism of formation, as phreatomagmatic. The Cetate breccia was exploited during centuries at the surface as well as in the underground. Within the Cetate breccia, different genetic types of mineralization were pointed out (Tămaş, 2002): impregnations, quartz-adularia veinlets, quartz (amethyst) veinlets and geodes, quartz-pyrite veinlets, veins rich in Mn-gangue minerals, phreatic breccias, clast-supported breccias. Reflected light microscopy, electron microscopy and microprobe investigations were carried out (Tămaş, 2002) in order to clarify the mineralogy of ore minerals in the Roşia Montană ore deposit. The present contribution focuses on minerals of the tetrahedrite-tennantite group, identified in different -mineralizations within the Cetate breccia, as well as in two breccia bodies from the Cărnăc Hill (Cărnăc I, and Cărnăc II breccias).

The chemical composition of tetrahedrite grains was determined by microprobe analyses (CAMECA SX-50, from BRGM, Orleans, France; accelerating voltage 20kV, beam current of 20nA, minimum detection limits ca. 0.1 wt. % for all elements). These data show large variations of the major elements of the tetrahedrite being emphasized, as follows: 24.04 - 42.99 wt.% Cu, 0 - 15.99 % wt. Ag, 0.33 - 2.42 wt.% Fe, 5.33 - 7.36 wt.% Zn, 0.08 - 29.47 wt.% Sb, 0.97 - 19.94 wt.% As, 22.37 - 28.1 wt.% S. Small quantities of other elements were detected: 0 - 6 wt.% Pb, 0.18 - 0.38 wt.% Mn, 0 - 1.33 wt.% Cd. The weight percentages were converted to the number of atoms based on 29 atoms per formula unit (acc. to Johnson *et al.*, 1986). The number of atoms per formula unit (*p.f.u.*) for each element in the structural formula of tetrahedrite is variable: 7.06 - 10.26 Cu, 0 - 2.71 Ag, 0.10 - 0.72 Fe, 1.25 - 1.82 Zn, 0.01 - 4.03 Sb, 0.22 - 4.04 As, 12.67 - 13.21 S. The cell dimensions were calculated using the compositions, in atoms per formula unit (Johnson *et al.*, 1987). The values of parameter "a" range between 10.22(3) Å and 10.51(3) Å.

⁷⁵ „Babeş-Bolyai” University of Cluj-Napoca, 1, Kogălniceanu Str., Dept. of Mineralogy, RO-3400 Cluj-Napoca, Romania. E-mail: <ctamas@bioge.ubbcluj.ro>, <dcostin@bioge.ubbcluj.ro>.

The number of S atoms is very close to 13, which represents the ideal number of S for tetrahedrite. The inverse correlation among Sb and As indicates the presence of a continuous solid solution tetrahedrite – tennantite. Meanwhile, the plots of Fe vs. Zn display substitutions close to ideal 2 atoms (Fe, Zn) stoichiometry. Tetrahedrite from Roşia Montană is generally rich in Sb. Fe content is generally very low comparing to Zn content. The analysed tetrahedrites contain 10 atoms of Cu per formula unit or less. Cu-Ag substitution, up to approximately 2 atoms *pfu* determines the smaller values than 10. This substitution is confirmed by the negative correlation between these two elements. Most data indicate a distinct positive correlation between Sb and Ag contents. The plot of Ag vs. As contents of the same data, shows that Ag and As have little tolerance for each other (acc. to Johnson *et al.*, 1986). Data plots indicate a relationship between the unit cell dimension and the number of Ag atoms, i.e. the cell size expands with increasing of Ag content. The values of (As+Sb) vs. S and (Ag+Cu) vs. S with 4 vs. 13 and 10 vs. 13 respectively, are constant, indicating an almost ideal stoichiometry.

The chemical composition of tetrahedrite-tennantite from Roşia Montană epithermal ore deposit is strongly dependent on the styles of mineralization. For instance, we may separate at least three types of tetrahedrite:

1. type 1 – intermediate Ag-tetrahedrite (1.40-1.92 atoms p.f.u.), related to quartz-adularia veinlets within Cetate breccia;
2. type 2 – tetrahedrite with heterogeneous Sb-As participation, with As ranging from 0.22-4.04 atoms p.f.u., and without Ag, related to quartz-pyrite veins hosted by Cetate breccia;
3. type 3 – Ag rich tetrahedrite (1.72 – 2.71 atoms p.f.u.), from Cărnic I breccia body.

References

- Cauuet, B., Ancel, B., Rico, Ch., Tămaş, C. (2003) Ancient mining networks. French Archaeological Expeditions 1999-2001 (in Romanian). In: Damian, P. (Ed.) *Alburnus Maior I*, Ed. CIMEC, 530 p., Bucureşti, 471-530.
- Balintoni, I. (1997) The geotectonic features of the metamorphic terrains from Romania (in Romanian). Ed. *Carpatica*, Cluj-Napoca, 176 p.
- Hedenquist, J.W. (2003) Epithermal ore deposits – styles, characteristics and exploration. *Short Course, University Babeş-Boyai Cluj-Napoca, 23 April 2003*.
- Johnson, N.E., Craig, J.R., Rimstidt, J.D. (1986) Compositional trends in tetrahedrite. *Canadian Mineralogist*, **24**, 385-397.
- Johnson, N.E., Craig, J.R., Rimstidt, J.D. (1987) Effect of substitutions on the cell dimension of tetrahedrite. *Canadian Mineralogist*, **25**, 237-244.
- Mârza, I., Tămaş, C., Ghergari, L. (1997) Low sulfidation epithermal gold deposits from Roşia Montană, Metaliferi Mountains, Romania. *St. Cerc. Geol., Geofiz., Geogr., Ser. Geol.*, **42**, 3-12.
- Săndulescu, M. (1984) The geotectonic of Romania (in Romanian). Ed. *Tehn.*, Bucureşti, 336 p.
- Tămaş, C.G. (2002) Breccia pipe structures related to some hydrothermal ore deposits in Romania, (unpubl. Ph. D thesis), 336 p.
- Tămaş, C.G., Bailly, L. (1998) Fluid inclusion study for Roşia Montană ore deposit. *Rom. Journ. Mineral Deposits*, **78**, Suppl. 1, 97-98.
- Tămaş, C.G., Bailly, L. (1999) Roşia Montană low-sulfidation ore deposit – evidence from fluid inclusion study. *Studia Univ. Babeş-Bolyai, Geologia*, **XLIV**, 1, 49-56.

THE RELATIONSHIPS BETWEEN BRECCIA STRUCTURES AND ORE VEINS. A CASE STUDY FROM THE CETATE HILL, ROŞIA MONTANĂ ORE DEPOSIT (ROMANIA)

Călin G. Tămaş⁷⁶, Lucreţia Ghergari¹, Corina Ionescu¹, Béatrice Cauuet⁷⁷

Introduction

The Roşia Montană epithermal ore deposit (Fig. 1) is well known due to its gold content and the Cetate phreatomagmatic breccia pipe structure (Tămaş, 2002). This breccia body that occurs in the Cetate Hill is not the single breccia structure within this massif, other breccia structures were identified additionally at the surface and in the underground.

The genesis of the Cetate phreatomagmatic breccia pipe was followed by subsequent phreatic processes, finally resulting in several phreatic breccias and veins, superimposed on the previous maar-diatreme couples (Tămaş, 2002).

Two ancient mining fields were discovered in the Cetate Hill - Zeus and Găuri - and subsequently used for archaeological-mining investigations (Cauuet *et al.*, 2003). Within these ancient mining networks (Fig. 1) two new breccia structure occurrences were discovered allowing to point out various aspects regarding the relation between breccias and the veins.

Fig. 1 – The geological map of the Roşia Montană area: 1 – Cretaceous flysch; 2 – Neogene sedimentary rocks; 3 – Cetate-type dacite; 4 – Phreato-magmatic breccia pipes (the Cetate breccia); 5 – Fluidization channel of the Cetate breccia pipe (“Glamm” Formation); 6 – Base-surge deposits; 7 – Andesite pyroclastic deposits (Rotunda type); 8 – Faults; 9 – Lakes.

Zeus mining field

⁷⁶ „Babeş-Bolyai” University, Department of Mineralogy, 1, Kogălniceanu Str., RO-3400 Cluj-Napoca, Romania. E-mail: <ctamas@bioge.ubbcluj.ro>, <ghergari@bioge.ubbcluj.ro>, <corinai@bioge.ubbcluj.ro>.

⁷⁷ Université Toulouse Le Mirail, 5, Allée Antonio Machado, 31520 Toulouse, France.

The ore body in the Zeus mining field is represented by a breccia dyke structure (Tămaş, 2002), completely hosted by the Cetate dacite. At the surface, the breccia body has 20 m length and a variable width, ranging from 0.5 up to 2 m. The known vertical extension of the Zeus breccia body is thought to reach up to 20 m. The breccia grades into unbrecciated dacite. The breccia is monolithologic (dacitic) and clast-supported. The matrix is missing, only a thin rim of quartz cement being present around the angular dacite fragments. The potassic alteration (formation of adularia) is the most representative one. Sericite alteration is also present, but no silicification was noticed so far.

Two mineralization stages can be distinguished for the Zeus breccia dyke: a pre-tectonic stage and a post-tectonic one. The first one is represented by small crystal of pyrite scattered inside the dacite clasts. This mineralization is a pre-brecciation event. The post-tectonic (post-brecciation) stage generated two types of mineral association: a) thin quartz rims around the dacite clasts (only of minor importance due to the very small width of the rims) and b) vein structures (four vein structures are so far noticed) that cross-cut longitudinally the breccia dyke structure.

The metal input is different in different parts of the breccia dyke. For example, the ore grades within the breccia body, where there is no vein overprinting, are about 1g/t Au and 3 g/t Ag⁷⁸. The ore grades along the veins including a narrow breccia zone rise to more than 4.2 g/t Au and 42 g/t Ag. For the whole breccia body, including the overprinting veins, the grades reach an average value of about 2.2 g/t Au and 10 g/t Ag.

Găuri mining field

The Găuri mining field is located in the south-western part of the Cetate massif. The modern mining level (+855 level) cross cuts a breccia body, which also occurs on the surface, on the northern slope of the Găuri area.

The Găuri breccia body has a pipe-like shape, with 20 m diameter. The pipe is slightly tilted towards the north. It was also partially opened by an ancient fire-setting exploitation zone (stopping). The Găuri breccia pipe is visible in the ceiling and on the walls in the upper level of the exploitation zone.

The breccia body is a matrix-supported structure with the matrix exceeding the clasts participation. In the underground (both in the remnants of the ancient stopings and in the modern mining network) as well as at the surface, various rocks were identified: metamorphic rocks (garnet micaschists, gneisses), sedimentary rocks (Cretaceous flysch, as sandstones and shales) and magmatic rocks (dacites). The metamorphics and the Cretaceous flysch fragments are rounded, while the dacite clasts are in general angular. The contacts of the breccia body with the host rock (the Cetate dacite) are sharp, although within the host rock a network of sheeted fissures develops. At the surface, on the northern slope of the Găuri area, the eastern rim of the breccia body is marked by a well-defined banded quartz vein (about 2.5 cm width). The fragments are included in a finely comminuted rock-flour type matrix. Both the breccia body and the host rock (dacite) are adularized and sericitized. Furthermore, intense silicification of only the dacite host rock and the breccia contacts was noticed.

The ancient stopping from the Găuri mining field focused on a vein-like structure, in fact a narrow stockwork zone containing "chinga"⁷⁹ and quartz veinlets. It has a variable width, from few centimeters up to 1 m. An intense silicification is to be noticed. The main ore minerals are pyrite, Ag-minerals and electrum. The mean ore grades of the ore in this stockwork structure are 2.2 g/t Au and about 30 g/t Ag.

This structure penetrates only few meters inside the main Găuri breccia body, being stopped due to the abundance of clay-sized material within the breccia matrix (very fine comminuted rock-flour matrix). In this case, contrary to the case of the Zeus mining field, the breccia body did not allow the development of the subsequent vein-like structure. The ore grades gradually decrease along the crosscutting of the vein through the breccia body, from 0.65 g/t Au and 23 g/tAg close to the breccia – dacite contact up to 0.3 g/t Au and 5 g/t Ag few meters away.

Conclusions

⁷⁸ Made by Roşia Montana Gold Corporation, Analabs Gura Roşia.

⁷⁹ A local name used for a mixture of very fine grained minerals, mainly clay minerals and silica, forming a fluid mud which was "injected" under some pressure, inside the breccias or dacite.

The different peculiarities of the Zeus and Găuri breccia structures, as well as their relationship with the later hydrothermal activity, are summarized in Table 1. Important differences, besides several common features, were pointed out for these breccia structures.

Table 1.

Comparative presentation of the breccias from the Zeus and Găuri ancient mining fields, Cetate Hill, Roşia Montană

Peculiarities	Zeus mining field	Găuri mining field
Brecciation mechanism	Tectonic	Phreato-magmatic
Morphology	Tabular – breccia dyke	Columnar – breccia pipe
Clast vs. matrix participation	Clasts prevail; clast-supported breccia;- - open-space breccias	Matrix prevail; matrix-supported breccias
Clast lithology	Monolithologic (dacite)	Polyolithologic (dacite, metamorphics, Cretaceous flysch)
Matrix composition	Quartz and adularia cementing coarse dacite fragments	Fine grained fragments of different rocks (rock flour) host the clasts
Breccia – host rock contact	Gradual	Sharp
Alteration	Adularia and sericite alteration; silicification is missing	Adularia and sericite alteration; very intense silicification of the contacts and host rock
Favorizing factors	Optimal: subsequent hydrothermal cement enveloped the rock fragments and several vein structures crosscut the breccia body rising the ore grades	Minimal: the development of subsequent veins/stockwork was stopped by the breccia body; the ore grades decrease within the breccia body
Mineralization	Common sulfides, Ag-minerals, gold	Common sulfides, Ag-minerals, gold

References

- Cauuet, B., Ancel, B., Rico, Ch., Tămaş, C. (2003) Ancient mining networks. The French archaeological missions 1999-2001. In Damian, P. (Ed.) - *Alburnus Maior I, Monographic series*, 531 p., Bucharest; p. 467-526.
- Tămaş, C.G. (2002) Breccia-pipe structures associated to some hydrothermal ore deposits from Romania (in Romanian). *Ph.D. Thesis*, Univ. Babeş-Bolyai, Cluj-Napoca, Romania, 336 p.

MINERAL INCLUSIONS IN RUBY FROM ANDILAMENA, MADAGASCAR

Panjawan Thanasuthipitak⁸⁰, Theerapongs Thanasuthipitak¹, Kobkarn Phetphu¹

Gem – quality rubies from the Andilamena area in eastern Madagascar have recently entered the gemstone market in commercial quantities. These rubies typically occur as short-prismatic or plate-like crystals with saturated purplish red to red colour. Triangular growth pattern on their basal surfaces is common. Mineral inclusions in 31 out of 245 samples were studied using Raman spectroscopy and scanning electron microscopy. The most common mineral inclusions are rutile, apatite and zircon. Monazite is sometimes identified. Staurolite, spinel, mica(?) and pyrite(?) are occasionally found.

Rutile occurs as both primary and secondary crystals. Primary rutile forms isometrical to elongated crystal of orange to reddish brown or gray to black colour with metallic luster standing out in high relief. Secondary short rutile needles form dense clouds often concentrating in the center of some ruby crystals. These represent exsolved rutile needles and are oriented in 3 directions, parallel to second-order hexagonal prism. After heat treatment, the secondary rutile needles were completely dissolved.

Apatite forms colorless euhedral to subhedral short prismatic crystals with rounded edges. Colourless zircon occurs as both, isolated rounded grains, with or without halos, as well as tiny clusters. Colourless to light yellow crystals of monazite are not as common as rutile and apatite.

Other minerals, namely staurolite, spinel (hercynite), mica(?) and pyrite(?) are also noticed.

According to the mineral inclusions and despite the relatively high iron content (ranging from 0.269 to 0.421 wt% FeO) of the Andilamena ruby, which indicates a genetic relation to basalts, the presence of certain inclusions such as staurolite suggests a metamorphic affiliation.

⁸⁰ Chiang Mai University, Dept. of Geological Sciences, 50200 Chiang Mai, Thailand. E-mail: tpitak@hotmail.com.

PETROLOGY OF GARNET-AMPHIBOLITE WITH WHITE MICA FROM VRANJSKA BANJA SERIES (SERBIAN-MACEDONIAN MASSIF, SE SERBIA)

Nada Vaskovic⁸¹, Vesna Matovic¹, Danica Sreckovic-Batocanin¹

Introduction

The Upper Complex of the Serbian-Macedonian Massif (SMM) named also Vlasina Complex – (VC) is mostly composed of greenschist with subordinate medium to high grade metamorphics (Dimitrijevic, 1959; Petrovic, 1965). This Ripheo-Cambrian complex was firstly intruded by Ordovician granitoids, later by Paleogene Surdulica granitoid and underwent the extensive Oligocene-Miocene volcanic activity. According to the spatial position, lithology and metamorphic features, the VC was divided into six series as follows: Vlasina (Vaskovic & Tasic, 2000), Vranjska Banja, Bozica, Lisina, Jaresnik and Stajevac (Babovic *et al.*, 1977).

According to Karamata & Krstic (1996), the Vlasina Complex represents a composite unit termed Ranovac-Vlasina-Osogovo terrane which is a part of the Carpatho-Balkanides Composite Terrane (Fig. 1).

Fig. 1. Terranes of East Serbia (acc. to Karamata & Krstić, 1996). CBCT - Carpatho-Balkanides Composite Terrane; VCMT - Vrska Cuka-Miroc terrane, SPPT - Stara Planina - Porec terrane, KT - Kučaj Terrane, HT - Homolje terrane, RVOT - Ranovac-Vlasina-Osogovo terrane. SMCT - Serbo-Macedonian Composite Terrane, VZCT - Vardar Zone Composite Terrane, VBS - Vranjska Banja Series.

⁸¹ Faculty of Mining and Geology, Džusina 7, 11 000 Belgrade, Serbia. E-mails: <nadavask@mkpg.rgf.bg.ac.yu>, <vesnamat@mkpg.rgf.bg.ac.yu>, <danicasreckovic@yahoo.com>

This study focuses on the medium to high grade metamorphic unit named Vranjska Banja Series (VBS). It comprises the southwestern part of the Vlasina Complex and is composed of a variety of gneisses, micaschists and amphibolites with subordinate occurrences of stromatitic migmatites, quartzites and marbles (Babovic *et al.*, 1977, Vaskovic & Tasic, 2000).

The purpose of this paper is to present the first petrologic data for the garnet-amphibolites with white mica found during the geological mapping of this area (Vaskovic & Tasic, 1996).

Geological framework

The VBS contains a group of polymetamorphic rocks affected by medium and high grade metamorphic events and localized lower grade overprint (Vaskovic, 1998).

The VBS constitutes the southwest part of the VC and extends east of Juzna Morava River (Fig. 1). According to Babovic *et al.* (1977), the VBS represents the deepest part of the VC, progressively metamorphosed under the influence of the Caledonian granitoid intrusions (Bozica and Jaresnik). To the west VBS is bounded by Juzna Morava River and to the south by the greenschists of the Stajevac Series.

Gneisses and micaschists constitute a major proportion of the rocks and yield important petrological constraints (Vaskovic & Milovanovic, 1996; Vaskovic, 1998) regarding the P-T evolution of SMM as a very important geological unit in the structural framework of the Balkan Peninsula.

During its long geological evolution the VBS has been affected by three phases of deformation (D1, D2, D3) and metamorphism (M1, M2, M3) which resulted in a very composite fabric (Babovic *et al.*, 1977; Vaskovic & Tasic, 2000; Vaskovic, 1998). According to Vaskovic (1998), M1 is characterized by a low-grade mineral assemblage (350-450°C, 3.5 kbar) defining S₁; as a result of prograde P-T evolution M2 was recorded by a continuous growth of garnet, staurolite and kyanite under conditions of 520-630°C and 5-7 kbar; M3 is characterized by transformation of garnet to chlorite and staurolite and kyanite to sericite, as well as by formation of stilpnomelane in micaschists in the northeastern part of the VBS. The predominant planar fabric represents a regional schistosity developed during D2 deformational episode. During and after D2 an extensional crenulations cleavage locally developed.

The southwestern part of this series shows a monoclinical structure dipping to the southwest. In the northern part this structure softly curves to the west-southwest (Babovic *et al.*, 1977).

Petrology

Amphibolites occur as thin elongated lenses, and rarely as dykes up to 20 m thick, parallel to the main schistosity within gneisses of the VBS. According to their mineral assemblage, three varieties can be distinguished: common amphibolite, garnet-amphibolite and garnet-amphibolite with white mica. The first two varieties occur mostly at the northern and western part of the VBS while the third appears only at its southeastern part close to the Viti Bor village (Fig. 1) as a lens of about 8 m thick and 50 m long, within two-mica gneisses and garnet-chlorite gneisses. It is schistose, dark green in colour, medium- to coarse-grained with large porphyroblasts of garnet (up to 1 cm) and very thin mica-rich accumulations (0.2 to 0.5 cm thick and 1-2.5 cm long).

The amphibolite from the Viti Bor village is composed of hornblende, plagioclase, garnet, white mica, chlorite and small amounts of quartz. As accessory minerals, leucoxene, ilmenite, titanite and apatite are found.

Hornblende (45.25 % vol. rock) occurs as prismatic or anhedral porphyroblasts up to 0.7-3 mm long or makes up to 0.3 mm long matrix grains. It forms irregular elongated lenses arranged as linear sets parallel to main schistosity or bands up to 1.5 mm thick in association with garnet, plagioclase, epidote, rarely quartz, and ilmenite/leucoxene or titanite. Some larger porphyroblasts are poikilitic, quartz and ilmenite being the most common inclusions. According to Leake *et al.* (1997), it is a magnesio-hornblende, usually homogeneous in composition. However, some grains are characterized by a little higher value of Na^A and Na^{M4}.

Garnet (13.24 % vol. rock) is euhedral to subhedral, about 0.5 to 4 mm, rarely up to 1 cm, in size and often includes quartz grains of less than 0.1 mm. Chemical zoning is not remarkable in the grains. Almandine varies from 63-65 mol %, grossular from 19-22 mol. %, pyrope from 2-5 mol. % and spessartine from 5-6 mol. %.

Plagioclase (9.25 % vol. rock) occurs in subhedral to anhedral porphyroblasts up to 1 mm long or forms elongated aggregates with quartz and/or amphibole. It includes epidote-zoisite, quartz, amphibole and ilmenite. The chemical composition falls in the range of oligoclase (21 to 27% an).

White mica (5.50 % vol. rock) is 0.1 to 1.25 mm long. It forms aggregates or bands up to 1 mm thick between amphiboles. The Si content ranges from 6.3 to 6.6, Mg from 0.29 to 0.46 and Fe from 0.22 to 0.32. All analysed flakes plot close to join $Si+0.5Al=9$ indicating that the white mica belongs to the muscovite-celadonite series termed phengite and phengitic muscovite (Graeser & Niggli, 1967). Both white micas show a minor internal variation in composition from core to rim.

Chlorite occurs as primary and retrograde phase. The first one is associated with white mica and amphiboles, while the second one is a product of garnet and amphibole alteration. The primary flakes range from 0.15 to 1.5 mm in length and sometimes form irregular lenses. Its orientation is parallel to the main schistosity. According to the classification of Hey (1954), the chlorites are repidolites ($X_{Fe}=0.46$) with quite homogeneous Si (5.21) and Al (2.78-2.79) contents.

Epidote is anhedral or tabular to stubby prismatic, mostly elongated grains ranging from 0.1 to 1 mm in length. The tiny grains appear as inclusions in plagioclase. It is the main interstitial phase settled between amphiboles and white mica. Very rarely it forms accumulations. Microprobe analyses were made on grains associated with white mica and grains associated with amphiboles. The first one has higher content of Al and lower content of Fe and Ca than the second one. The $Fe^{3+}/Al+Fe^{3+}$ value of epidotes vary from 0.145 (grain associated with white mica) to 0.224 (grain associated with amphibole). According to classification of Holdaway (1972) they correspond to Al-epidote. The decrease of pistacite content suggest condition of moderate $f(O_2)$ during metamorphism (Liou, 1973).

Titaniferous minerals (ilmenite, leucosene, titanite) and apatite occur as interstitial phases or inclusions in amphiboles.

P-T condition of metamorphism

Ca-amphibole (hornblende) compositions are sensitive to changes in metamorphic condition, especially to changes in temperature (e.g. Laird & Albee, 1981; Spear, 1981; Holland & Blundy, 1994).

The mineral assemblage of the amphibolite (Mg-hornblende + garnet + plagioclase + white mica + epidote) enables the application of the garnet-hornblende - GH (Graham & Powell, 1984) and garnet-plagioclase thermometer - GP (Blundy & Holland, 1990), as well as the garnet-hornblende-plagioclase-quartz (GHPQ) barometer (Kohn & Spear, 1990).

Calculations of temperatures using GH and GP are not consistent. Temperatures estimated by Graham & Powell method range between 420 and 485°C. The GP calibration of Blundy & Holland (1990) yields temperatures of 530-585°C. These temperatures are consistent with the 520-630°C formation temperatures estimated for the micaschists (Vaskovic, 1998) and with 520-550°C for the garnet-chlorite-white mica and garnet-chlorite gneisses (Vaskovic & Milovanovic, 1996).

Pressure estimated for four samples in conjunction with temperatures estimated from the GP varies from 5.2 to 6.1 kbar. Thus, it is in accordance with pressures of 5.1 –7.4 kbar calculated for micaschists of the VBS (Vaskovic, 1998).

The experimental work of Velde (1965, 1967) and Massone & Schreyer (1987) shows that phengite with Si content ≥ 3.3 p.f.u. crystallizes under rather high P/T conditions. Si content of 6.3-6.6 p.f.u. in white mica suggests pressure of 4-5 kbar in the temperature range of 350 to 450°C according to Massone & Schreyer (1987). That is not in agreement with P-T conditions estimated using garnet-plagioclase thermometer and garnet-hornblende-plagioclase-quartz barometer.

Conclusion

Among amphibolitic rocks within the Vlasina Complex (Upper part of the Serbian-Macedonian Massif), occurring as lenses, rarely dykes or small bodies, three varieties have been recognized according to the mineral composition: common amphibolite, garnet amphibolite and garnet-amphibolite with white mica (Vaskovic, 1998). No data are available about their chemistry, but relics of primary fabric in some occurrences revealed their igneous origin.

The garnet amphibolites with white mica studied here belong to the Vranjska Banja Series being associated with gneisses and micaschists metamorphosed in the temperature range of 520-

630°C under the pressure of 5.1 –7.4 kbar (Vaskovic, 1998). For them the peak of metamorphism occurred at temperature range of 530 to 585°C under pressure of 5.2 to 6.1 kbar.

This study has shown that the metamorphism, which affected the rocks of the Vranjska Banja Series, had similar intensity in the whole area.

References

- Babovic, M., Roglic, C., Avramovic, V., Maric, S. (1977) Explanatory book for the sheet Trgoviste with Radomir. Basic Geological maps 1:100 000, Federal Geological Institute, 58 p., Belgrade (in Serbian with English summary).
- Blundy, D., Holland, J.B. (1990) Calcic amphibole equilibria and new amphibole-plagioclase geothermometer. *Contrib. Min. Petrol.*, **104**, 208-224.
- Dimitrijevic, D.M. (1959) Basic characteristics of the column of the Serbian-macedonian Mass. *First Symposium of Serbian Geological Society* (oral report), Belgrade.
- Graeser, S., Niggli, E. (1967) Zur Verbreitung der Phengite in den Schweizer Alpen: ein Betrag zur Zoneographiie der Alpinen Metamorfoze. In: *Etage tectoniques. Colloque de Neuchatel*. Institut de Geologie de l'Universite' de Neuchatel, 33-42.
- Graham, C. M., Powell, R. (1984) A garnet - hornblende geothermometer. Calibration, testing, and application to the Pelona Schists, southern California. *J. M. G.*, **2**, 13-31.
- Hey, M.H. (1954) A new review of chlorites. *Mineral. Mag.*, **30**, 277-292.
- Holdaway, M.J. (1972) Thermal stability of Al-Fe epidote as a function of fO₂ and Fe content. *Contrib. Min. Petrol.*, **37**, 307-340.
- Holland, T., Blundy, J. (1994) Non-ideal interactions in calcic amphiboles and their bearing on amphibole-plagioclase thermometry. *Contrib. Min. Petrol.*, **116**:433-447.
- Karamata, S., Krstić, B. (1996) Terranes of Serbia and neighbouring areas. In *Terranes of Serbia* (Eds. Knezevic V. & Krstic B.), 20-40, Belgrade.
- Kohn, M.J., Spear, F.S. (1990) Two new geobarometers for garnet amphibolite, with applications to southern Vermont. *Am. Mineral.*, **75**, 89-96.
- Laird, J., Albee, A.L. (1981) Pressure-temperature and time indicators in mafic schists: their application to reconstructing the polymetamorphic history of Vermont. *Am. J. Sci.*, **281**, 127-175.
- Leake, B.E., Wooley, A.R., Arps, C.E.S., Birch, W.D., Gilbert, M.C., Grice, J.D., Hawthorne, F.C., Kato, A., Kisch, H.J., Krivovichev, V.G., Linthout, K., Laird, J., Mandarino, J.A., Maresch, W.V., Nickel, E.H., Rock, N.M.S., Schumacher, J.C., Smith, D.C., Stephenson, N.C.N., Ungaretti, L., Whittaker, E.J.W., Youzhi, G. (1997) Nomenclature of amphiboles: report of the Subcommittee on Amphiboles of the International Mineralogical Association, Commission on New Minerals and Mineral Names. *Canad. Mineral.* **35**, 219-246.
- Liou, J.G. (1973) Syntesis and stability relations of epidote, Ca₂Al₂FeSi₃O₁₂(OH). *J. Petrol.*, **14**, 381-413.
- Massone, H.J., Schreyer, W. (1987) Phengite barometry based on the limiting assemblage with K-Feldspar, phlogopite and Quartz. *Contrib. Min. Petrol.*, **96**, 312-324.
- Petrovic, B. (1965) Fabric of the Vlasina Crystalline Complex in the area of the Crna Trava. Ph. D. Thesis, Faculty of Mining and Geology, Belgrade University (unpublished), 250 p. (in Serbian)
- Spear, F.S. (1981) Amphibole-plagioclase equilibria: an empirical model for the relation albite + tremolite + edenite + quartz. *Contrib. Min. Petrol.*, **77**, 355-364.
- Vaskovic, N. (1998) P-T condition of the mica schists from the Vranjska Banja Series. *13th Yugoslavian Geological Congress*, **3**, 41-59, Herceg Novi (in Serbian, English summary).
- Vaskovic, N., Milovanovic, D. (1996) Garnet-white mica-chlorite gneisses of the Viti Bor (Vranjska Banja Series, Serbia). In *Terranes of Serbia* (Eds. Knezevic & Krstic), 183-190. Belgrade.
- Vaskovic, N., Tasic, Z. (1996) Geological map of the Surdulica granodiorite massif 1: 50 000. Federal Ministry of Economy, Belgrade.
- Vaskovic, N., Tasic, Z. (2000) Geology of the Surdulica Granodiorite masiff and its surroundings. Ed. Federal Ministry of Economy, 128 p. (in Serbian), Belgrade.
- Velde, B. (1965) Phengite micas: synthesis, stability and natural occurrences. *Am. J. Sci.*, **263**, 886-913.
- Velde, B. (1967) Si⁺⁴ content of natural phengites. *Contrib. Min. Petrol.*, **14**, 250-258.

MINERALOGY OF PLIOCENE TO PLEISTOCENE PELITIC SEDIMENTS OF THE GREAT HUNGARIAN PLAIN

Istvan Viczián⁸²

This paper deals with the mineralogical composition of pelitic sediments of the Great Hungarian Plain. Published data as well as unpublished data on analyses made in the laboratories of the Geological Institute of Hungary based on almost 150 samples are presented. Determinations were made mainly by X-ray diffraction, the data were systematically corrected by comparison with the results of thermal analysis and partly chemical analysis. All the data were revised and recalculated in a uniform system in order to obtain comparable results.

In the bulk composition the dominant clay minerals are *smectite*, *illite/smectite*, *illite* and *chlorite*. In the <2 µm fraction the same minerals occur, however, expanded phases are more dominant. Triple mixed-layer *illite/smectite/chlorite* and *kaolinite* of various degree of disorder may appear. Clay minerals are essentially detrital, derived from various areas of the surrounding Carpathians and Alps (Viczián, 2002). Sub-basins may differ in degree of disorder and quantitative proportions of clay minerals and quantitative relations of other phases like *calcite*, *dolomite*, *quartz* and *feldspars* depending on relatively permanent source areas and transport directions. Smaller variations in the transport directions as shown by the micromineralogical composition are normally not reflected in the clay mineral record. Climatic variations during the Pleistocene seem to have no significant effect. In the South Tisza Basin and Maros Alluvial Fan *well crystallised detrital phases* prevail, while in the Körös Basin more mature sedimentary material of *lesser crystallinity*, higher *kaolinite* and *very low carbonate contents* can be found.

The clay, carbonate, feldspar and iron minerals deposited may have been modified by flow systems of ground water (Varsányi *et al.*, 1997, Varsányi & Ó. Kovács, 1997). In the upper flow regime comprising Pleistocene and Pliocene horizons of the South Tisza Basin and Maros Alluvial Fan dissolution of *carbonate minerals* and *albite* and ion exchange on *clay minerals* may occur. In the stagnant ground waters filling the Pleistocene and Pliocene beds of the Körös Basin neof ormation of *pure smectite* and *kaolinite* from dissolution of *albite* and dissolution of *carbonates* may be inferred from hydrogeochemical and mineralogical data.

Amorphous iron hydroxides underwent crystallisation and reduction producing, in a downward sequence, amorphous "*limonite*", *goethite* and *siderite*.

No diagenetic K-fixation or illitisation occur in this level. However, some kind of *palaeopedological illitisation* may have occurred in those continental sediments. The first main step of *burial diagenetic illitisation* (Tanács & Viczián, 1995) as well as of kerogen diagenesis starts in the lower groundwater regime, which corresponds to the Upper Pannonian formation group, i. e. below the formations discussed in the present paper.

References

- Tanács, J., Viczián, I. (1995) Mixed-layer illite/smectites and clay sedimentation in the Neogene of the Pannonian Basin, Hungary. *Geol. Carpath., Ser. Clays*, **4**, 1, 3-22.
- Varsányi, I., Matray, J.-M., Ó. Kovács, L. (1997) Geochemistry of formation waters in the Pannonian Basin (southern Hungary). *Chem. Geol.* **140**, 89-106.
- Varsányi, I., Ó. Kovács, L. (1997) Chemical evolution of groundwater in the River Danube deposits in the southern part of the Pannonian Basin (Hungary). *Appl. Geochem.* **12**, 625-636.
- Viczián, I. (2002) Clay mineralogy of Quaternary sediments covering mountainous and hilly areas of Hungary. *Acta Geol. Hung.* **45**, 3, 265-286.

⁸² Geological Institute of Hungary, Budapest, E-mail: viczian@mafi.hu

THE GENESIS OF CALCITE RAFTS FROM CAVES – A POINT OF VIEW

Iosif Viehmann⁸³¹, Lucreția Ghergari⁸⁴²

The present work is a conclusion of our previous observations and experiments concerning the genesis of calcite rafts (Viehmann, 1992; Viehmann *et al.*, 1997). The main raft deposit studied was discovered in 1987 in the Hoanca Apei Cave (Bihor Mountains, Romania). Our results were compared with the ones obtained by Black (1953) in Carlsbad Caverns (New Mexico, USA), Pomar *et al.* (1975) in Spain, and by Forti & Chiesi (1995) in Italy.

The main resemblance consists in the growth manner of the rhombohedral calcite crystals: these are partially developed, growing only on the lower part of the raft, at the contact with water. This crystallogenic mechanism conditions raft floating. The same conclusion was drawn by Pomar *et al.* in 1975.

The second cause of calcite rafts floating on the surface of undisturbed cave waters is the *per ascensum* elevation of the water in the pool. The filling of the cave pool may be repeated several times, with the same calcite rafts floating at each filling.

Another cause of raft floating is the network of skeletal crystals retaining small air bubbles. When calcite rafts become heavier, they sink in the pool and thicken the local sediment.

References

- Black, D.M. (1953) Aragonite Rafts in Carlsbad Caverns, New Mexico. *Science*, 117, 84-85
- Forti, P., Chiesi, M. (1995) A proposito di una particolare forma di calcite flotante. *Atti Mem. Comiss. Grote "E. Boegan" Trieste*, 32, 43-53,.
- Pomar, L., Gines, A., Moya, G., Ramon, G. (1975) Nota previa la petrologia y mineralogia de la calcite flotante. *Endins, publ. d'Espeleo Balear*, Mallorca, 2, 1-6.
- Viehmann, I. (1992) Experimental methods in studying the cave rafts. *Theoretical and Applied Karstology*, 5, 213-215.
- Viehmann, I., Ghergari, L., Onac, B.P. (1997) Cristallographical observations on calcite rafts from three Romanian caves. *Proceedings of the 12th International Congress of Speleology, La Chaux de Fonds*, Switzerland, 1, 227-416.

⁸³ "Emil Racovitza" Speleological Institute, Clinicilor 5, Cluj-Napoca, Romania
"Babeş-Bolyai" University, Dept. of Mineralogy, Kogălniceanu 1, Cluj-Napoca, Romania

MINE-WASTE MANAGEMENT

Şerban Vlad⁸⁵, Dan Costin¹

Mineral extraction and processing, especially metal mining, exposes fresh rock surfaces and produces crushed and milled waste. Such material, exposed to weathering, poses a potential danger to the environment. Sulfide-bearing mineralization, if oxidized, has the potential to produce: a) sulfuric acid, known as acid rock drainage (ARD), and b) increased heavy-metal mobility. Significant contamination of surface and groundwater may result from these two processes. Anyway, not all sulfide-containing waste materials present a threat to the environment, even though they may contain geochemically important heavy-metal concentrations.

Through detailed mineralogical and geochemical investigation, it is possible to evaluate and quantify the environmental threat from mine materials and materials to be mined. Based on the mineralogic and thermodynamic assessment of the geochemical processes that are likely to occur in unmined material or that are currently taking place within mined materials, geologically and economically reasonable waste management decisions can be made.

⁸⁵ "Babeş-Bolyai" University, Dept. of Mineralogy, 1, Kogălniceanu Str., RO-3400 Cluj-Napoca, Romania. Email: <vlad@bioge.ubbcluj.ro>, <dcostin@bioge.ubbcluj.ro>.

LOW MAGNITUDES OF MATRIX STRESS RELATED TO AQUATHERMAL EFFECT AND MONTMORILLONITE DEHYDRATION IN THE CĂLDĂRUŞANI-TULNICI GEODYNAMIC POLYGON, ROMANIA

Dorel Zugrăvescu⁸⁶, Gabriela Polonic¹, Victor Negoită¹

On the basis of various physical measurements carried out in very deep wells, a method for determining the subsurface stress magnitudes was elaborated by the Geodynamic Institute of the Romanian Academy. Using such a methodology a stress study was undertaken within the Căldăruşani-Tulnici Geodynamic Polygon (depth interval 5-7 km) whose results were discussed on the occasion of the International Geophysical Conference (April 2000, Bucharest-Romania). The significance of the stress magnitude variations within the polygon areal was also presented on the occasion of the International Geological Correlation Program IGCP-430 Workshop (June 2000, Covasna-Romania). The conclusions of this work focus on the Oligocene deposits units from the western part of the polygon, in which the grain-to-grain stress (*matrix stress*) decreases drastically.

As a cause of registered low stress values, an other essential information was specified in the above mentioned work, stating that "the thick pre-Sarmatian detritic sedimentary sequence (mainly shales with local evaporites) was rapidly and deeply buried to more than 8 km, giving rise, in sealing circumstances, to very large overpressures within the filling fluids of the rock pores.

All these overpressured geological formations, located along the western limit of the geodynamic polygon have been studied in our institute. We found, as a rule, geopressures as much as 30-50 MPa above the normal hydrostatic pressures. The geological formations are also characterized by shale undercompaction and pore space waters less saline compared with their surroundings.

Such macrovolumes of rocks seem to be entirely isolated without any connection to surface outcrops. In a continuous sedimentary deposition, the sealing moment occurred significantly earlier, at a shallower depth, when the temperature was lower than today.

As a rule, at the time of continuous sedimentation and burial of the geological formations, the temperature rises, leading to the increase of the volume of a given content of water. If the water is located in an open system it may expand freely and a certain quantity of water will be expelled. But, if the same system is closed as might be expected in a large, impermeable shale environment, the water may not expand to the same degree, and consequently, the pressure will increase by an aquathermal pressure effect.

The behaviour of such sealed rock macrovolumes in connection with temperature and pressure variations is the first topic of our work, as the pressure increasing rate in aquathermal conditions is higher than that in non-aquathermal conditions. The rate of this aquathermal pressuring effect is a function of the geothermal gradient and the capacity of the shale to retain the generated pressure.

That is why, we elaborated a proper methodology and adequate graphical solutions/representations relating temperature, water density and its specific volume.

Temperature maps at 5 km and 6 km depth have been drawn, geothermal gradient and pressure gradient calculations on the basis of various borehole measurements have been carried out and finally low values of grain to grain/matrix stress were determined for most of the boreholes cutting the 5-7 km depth interval of the geodynamic polygon.

⁸⁶ Geodynamic Institute of the Romanian Academy, 19-21, J.-L. Calderon Str., RO-70201 Bucharest-37, Romania. E-mail: <dorezugr@geodin.ro>.

The last part of our work deals with the montmorillonite dehydration showing that the montmorillonite alteration to illite starts in the depth interval 1.8–2.3 km and continues at an increasing rate to a depth interval of 3–4.5 km, due to the rock thermal conductivities and geothermal gradients variability on the entire territory of the geodynamic polygon.

This alteration process involves two main physico-chemical changes within rock water characteristics. The first change consists in desorbing the adsorbed water and its transfer into particle spaces as free water. The second step is represented by water expansion, which follows the rules presented in the first part of this paper.

In order to assess the stages of montmorillonite dehydration process, electric logs, sonic logs, temperature logs and radioactive logs (gamma ray, neutron and gamma-gamma) recorded during the customary borehole operations have been used.

Appendix:

LIST OF THE PAPERS PUBLISHED BY UNIV. PROF. DR. LUCREȚIA GHERGARI

UNIVERSITY COURSES

1. Arghir, G., **Ghergari, Lucreția** (1991) Crystallography and Mineralogy (in Romanian, in original: Cristalografie și mineralogie), Cluj-Napoca, 480 p.

GUIDE-BOOKS FOR PRACTICAL STUDIES

1. Arghir, G., **Ghergari, Lucreția** (1983) Crystallography - Mineralogy – Guidebook for practical studies (in Romanian, in original: Cristalografie - Mineralogie - Indrumător de lucrări practice), 371 p., *Technical University Cluj-Napoca*.
2. Arghir, G., **Ghergari, Lucreția** (1989) Crystallography - Mineralogy – Guidebook for practical studies (in Romanian, in original: Cristalografie - Mineralogie - Indrumător de lucrări practice), 371 p., 2nd edition, *Technical University Cluj-Napoca*.
3. Mureșan, I., **Ghergari, Lucreția**, Bedeleian, I. (1986) Mineral identification (in Romanian, in original: Determinator de minerale), vol. I, 396 p., vol. II: 233 p., *Technical University Cluj-Napoca*.

OTHER PAPERS

1. **Mirza, Lucreția** (1973) The use of membranes as sensors in acide-basic conductometric titration against a very conductive background (in Romanian, in original: Utilizarea membranelor ca senzori în titrarea conductometrică acido-bazică pe fond foarte conductibil; *Ph.D. Thesis abstract, Babeş-Bolyai University*, 49 p., Cluj-Napoca).
2. **Ghergari, Lucreția** (1994) The geologic and mineralogic teaching at Cluj University between 1919 - 1994 (in Romanian, in original: Invățământul geologic-mineralogic la Universitatea din Cluj în perioada 1919 - 1994, *Babeş-Bolyai University*, 41 p., Cluj-Napoca).
3. **Ghergari, Lucreția** – collab. at the vol. Lazarovici, Gh., Maxim, Z (1995) Gura Baciului – an archaeological monography (in Romanian, in original: Gura Baciului – monografie arheologică), *Ed. Muz. Nat. Ist. Transilv.*, 436 p., Cluj-Napoca, 1995.

SCIENTIFIC PAPERS

1. Stoicovici, E., **Ghergari, Lucreția**, Moțiu, A. (1957) Contribution to the knowledge and utilization of tourmaline (in Romanian, in original: Contribuții la cunoașterea și valorificarea turmalinei). *Bulet. Univ. "V. Babeș" și "Bolyai"*. Ser. Șt. Naturii, **I**, 1-2, 315-324.
2. Stoicovici, E., **Ghergari, Lucreția**, Mârza, I. (1959) The study of calcium, magnesium, iron and manganese carbonates from the Apuseni Mountains. I. The geological study of the Runc region with special regard to the metamorphic carbonates (in Romanian, in original: Studiul carbonaților de calciu, magneziu, fier și mangan din Munții Apuseni. I. Studiul geologic al regiunii Runc, cu privire specială asupra carbonaților metamorfici). *Studia Univ. Babeş-Bolyai*, Ser. Geol.-Geogr., **II**, 1, 107-131.
3. Maxim, Al., Lucca, V., Marincea, V., Clichici, O., Șuraru, N., Dușa, A., Florei, N., Șuraru, M., Ionescu, G., Moțiu, A., Ghiurcă, V., Nicorici, E., Băluță, Cr., **Ghergari, Lucreția**, (1960) Cluj limestones on the territory of Cluj (Building materials). I. The Baci Limestone (in Romanian, in original: Calcarele grosiere de pe teritoriul orașului Cluj (Materiale de construcție). I. Calcarele de la Baci (Cluj); *Studia Univ. Babeş-Bolyai*, Ser. Geol.-Geogr., **II**, 1, 7-62.
4. **Ghergari, Lucreția**, Mârza, I. (1961) The study of bentonite from Palazu Mare (Dobrogea) (in Romanian, in original: Studiul bentonitului de la Palazu Mare (Dobrogea); *Studia Univ. Babeş-Bolyai*, Ser. Geol.-Geogr., **II**, 1, 83-93.
5. Stoicovici, E., **Ghergari, Lucreția**, Mârza, I. (1962) The study of diagenetic phosphorite from Romania (in Romanian, in original: Studiul unor iviri de fosforit diagenetic din R.P.R.). *Stud. Cercet. Geol. Geogr.*, **VII**, 3-4, 577-595, București.
6. Mârza, I., **Ghergari, Lucreția**, (1963) Bentonite from the Chioarului Valley (Maramureș region) – Preliminary note [in Romanian, in original: Bentonitul de la Valea Chioarului (reg. Maramureș) - Notă preliminară]. *Rev. Minelor*, **1**, 41-43, București.

7. Mârza, I., **Ghergari, Lucreția**, (1963) Some observations regarding silicolites from the Senonian chalk (Southern Dobrogea) [in Romanian, in original: Observații privind silicolitele din creta senoniană din Dobrogea de sud]. *Studia Univ. Babeş-Bolyai*, Ser. Geol.-Geogr., 2, 7-14.
8. **Ghergari, Lucreția**, Mârza, I., Ionescu, G. (1964) Contributions to the geological and mineralogical study of bentonite from Ocna Mureş [in Romanian, in original: Contribuții la studiul geologic și mineralogic al bentonitului de la Ocna Mureş]. *Studia Univ. Babeş-Bolyai*, Ser. Geol.-Geogr., 1, 23-37.
9. Mârza, I., **Ghergari, Lucreția**, Ionescu, G. (1965) Orthoamphibolites from the Baia de Arieş Series, Orăști – Belioara region (Arieş Basin). [in Romanian, in original: Ortoamfibolitele din seria de Baia de Arieş, regiunea Orăști – Belioara, Bazinul Arieşului]. *Studia Univ. Babeş-Bolyai*, Ser. Geol.-Geogr., 1, 43-50.
10. Mârza, I., **Ghergari, Lucreția**, Mînzăraru, L. (1966) Considérations sur la gnse et la composition des bentonites de Rzoare (Roumanie); *Bull. Serv. Carte Gol. Als. Lorr.*, 19, 3-4, 213-220, Strasbourg.
11. Treiber, I., **Ghergari, Lucreția**, Mârza, I., Ionescu, G. (1967) Microtectonic research in the Jidovina Dacitic Massif (the Arieş Valley) [in Romanian, in original: Cercetri microtectonice n masivul dacitic Jidovina, V. Arieşului]. *Studia Univ. Babeş-Bolyai*, Ser. Geol.-Geogr., 2, 45-55.
12. Mârza, I., **Ghergari, Lucreția** (1967) A geological and mineralogical study of bentonite from the Valea Chioarului (Maramureş region) [in Romanian, in original: Studiul geologic și mineralogic al bentonitului de la Valea Chioarului, reg. Maramureş]. *Anal. Univ. Bucureşti*, Seria Şt. Naturii, Ser. Geol.-Geogr., XVI, 1, 31-45.
13. Şuteu, Al., **Mrza, Lucreția** (1970) L'tude roentgnostructurale des trihtropolyacides; *Revue Roum. de Chimie*, 15, 10, 1563-1572.
14. Liteanu, C., **Ghergariu-Mrza, Lucreția** (1970) Titration conductometric with resistance indicator. II - Titration of the acid chlorhydric in solutions concentrated of sodium chloride with resistance membrane PVC + plasticizer + Na-montmorillonite; *Revue Roum. de Chimie*, 15, 12, 1871-1882.
15. Liteanu, C., **Ghergariu-Mrza, Lucreția** (1971) The conductometric titration with indicating resistance. III – On the mechanism of the PVC+Allassion CS (cationite R-H)+diostylphthalate (plasticizer) membrane as indicating resistance [in Romanian, in original: Titarea conductometric cu rezisten indicatoare. III - Despre mecanismul de funcţionare a rezistenei indicatoare membran PVC + alasion CS (cationit R-H) + dioctilftalat (plastifiant)]. *2nd Nat. Conf. for Analytical Chemistry, Braşov*, I, 265-270, 22-26 septembrie 1971.
16. Auslander, D., Lenart, I., Rus, I., **Mrza, Lucreția** (1971) L'action des ultrasons sur le processus de cristallisation; *Seventh International Congress on Acoustics*, 24 M 2, 113-116, Budapest.
17. Şuteu, Al., **Mrza, Lucreția** (1972) Roentgenographic study of caesium phosphomolybdovanadate (in Romanian, in original: Studiul roentgenografic al fosfomolibdovanadatului de cesiu. *Lucrri şt., Ser. Matem.-Fiz.-Chim.*, 55-57, Oradea.
18. Liteanu, C., **Ghergariu-Mrza, Lucreția** (1972) The conductometric titration with indicating resistance - III. Mechanism of the PVC + Allassion CS (Cationite R-H) + dioctylphthalate (plasticizer) membrane as indicating resistance; *Talanta*, 19, 980-984, Great Britain.
19. Mrza, I., Sntimbreanu, A., Moiu, A., Plfi, S., **Mrza, Lucreția**, Curea, C. (1973) Pegmatites with molybdenite from Vina (the Apuseni Mountains) [in Romanian, in original: Pegmatitele cu molibdenit de la Vina (Munii Apuseni)]. *Stud. Cercet. Geol. Geofiz. Geogr.*, Ser. Geol., 18, 2, 317-327, Bucureşti.
20. Mrza, I., **Mrza, Lucreția**, Poszet, T. (1974) Hisingerite, a new mineral identified in the epimetamorphic copper ore from Fagul Cetţii (Blan) [in Romanian, in original: Hisingeritul, un nou mineral identificat n zcmntul cuprifr epimetamorfozat de la Fagul Cetţii (Blan)]; *Studia Univ. Babeş-Bolyai*, Ser. Geol.-Mineral., XIX, 1, 9-14.
21. Mrza, I., **Mrza, Lucreția** (1974) On the presence of hisingerite in pyrometasomatic iron ore from Maşca (Iara Valley, Apuseni Mountains) [in Romanian, in original: Asupra prezenei hisingeritului n zcmntul pirometasomatic de fier de la Maşca (Valea Iara, Munii Apuseni)]; *Stud. Cercet. Geol. Geofiz. Geogr.*, Ser. Geol., 19, 69-75, Bucureşti.
22. **Ghergari, Lucreția**, Bereczky, A., Bereczky, E. (1978) Morphodynamic research on barite from the basemetal ore from Nistru (Maramureş region) (in Rumanian, in original: Cercetri morfodinamice asupra baritului din zcmntul polimetalic de la Nistru (jud. Maramureş); *Studia Univ. Babeş-Bolyai*, Ser. Geol.-Geogr., XXIII, 1, 3-15.
23. Stoicovici, E., Mureşan, I., **Ghergari, Lucreția**, Bedeleian, I., Voiculescu, L., Blu, C., Gbos, L. (1978) L'tude minralogique, ptrographique et ptrochimique de la rhyolite kaolinise de Parva, dpartement de Bistria-Nsud; *Studia Univ. Babeş-Bolyai*, Ser. Geol.-Geogr., XXIII, 2, 3-18.
24. **Ghergari, Lucreția**, Liteanu, C. (1978) Conductometric titration with indicating resistance - IV. The use of membranes with some organic and inorganic acids as active components, for HCl titration in a highly conducting medium; *Talanta*, 25, 9-14, Great Britain.

25. Bedelea, I., **Ghergari, Lucreția**, Mârza, I., Moțiu, A., Mureșan, I., Țirlea, I. (1979) The catalogue of the meteorites collection from the Mineralogical Museum of the University of Cluj-Napoca; *Studia Univ. Babeş-Bolyai*, Ser. Geol.-Geogr., **XXIV**, 2, 3-22.
26. Mészáros, N., **Ghergari, Lucreția** (1979) Lithological and stratigraphical research of the Tertiary deposits from Rohia area [in Romanian, in original: Cercetări litologice și stratigrafice asupra depozitelor terțiare din regiunea Rohia (Târgu Lăpuș)]. *Studia Univ. Babeş-Bolyai*, Ser. Geol.-Geogr., **XXIV**, 2, 39-47.
27. Mârza, I., Bedelea, I., **Ghergari, Lucreția**, Mureșan, I., Voiculescu, L. (1981) Petrometallogenetic observations on the basemetal deposit from Ruschița; the extending area of the principal deposit [in Romanian, in original: Observații petrometalogenetice asupra zăcămintului polimetalic de la Ruschița, zona de extindere a zăcămintului principal]. *Studia Univ. Babeş-Bolyai*, Ser. Geol.-Geogr., **XXVI**, 1, 3-30.
28. Mureșan, I., **Ghergari, Lucreția**, Mârza, I., Bedelea, I. (1981) Contribution to the mineralogical characterization of the raw materials and of basic refractory products [in Romanian, in original: Contribuții la caracterizarea mineralogică a materiilor prime și a produselor refractare bazeice]. *Mem. Secțiilor Științifice Academ.*, Ser. 4, IV, 2, 243-258.
29. **Ghergari, Lucreția**, Mârza, I., Pomârjanschi, G., Hudrea, I. (1983) Argilisation hydrothermales associées aux roches volcaniques du complexe ophiolitique de la région Buru-Cheile Turzii, Monts Apuseni (Roumanie); *Studia Univ. Babeş-Bolyai*, Ser. Geol.-Geogr., **XXVIII**, 1, 2-13.
30. Micu-Semeniuc, R., Hila, E., Doboș-Roman, G., **Ghergari, Lucreția** (1983) Investigation of some complex metal thiocyanates with chlatration properties; *Revue Roum. Chimie*, **28**, 5, 471-485, 1983.
31. Mârza, I., **Ghergari, Lucreția**, Frățilă, V., Sudrigean, Tr., Mureșan, I., Bedelea, I., Pop, E., Bedo, M. (1983) Utilization test of bentonitic clays from Borzeti area – Cheile Turzii, with the view to obtain cement with high grade of whiteness [in Romanian, in original: Incercări de valorificare a argilelor bentonitice din zona Borzești-Cheile Turzii (jud. Cluj), pentru obținerea cimentului cu grad de alb superior]. *Simp. Valorificarea substanțelor nemetalifere*, 3, 65-73.
32. **Ghergari, Lucreția**, Petrescu, I., Simuț, D. (1985) Paleoclimatic and paleogeographic considerations of the Sarmatian formations from Aștileu (Oradea), based on the clay minerals study (a preliminary note) [in Romanian, in original: Aprecieri paleoclimatice și paleogeografice asupra sarmațianului de la Aștileu (Oradea), pe baza studiului mineralelor argiloase (notă preliminară)]. *Crisia*, **XV**, 467-471, Oradea.
33. Mârza, I., **Ghergari, Lucreția**, Bedelea, I. (1985) Alterations of Neogene volcanic tuffs from Transylvania (Romania); *VIII-th Congress of the Regional Committee on Mediterranean Neogene Stratigraphy Symposium on European Late Cenozoic Mineral Resources*, 360-362, Budapest, 15-22 sept., 1985.
34. Mureșan, I., **Ghergari, Lucreția**, Bedelea, I. (1986) Contributions to the petrographic characterization of the volcanic tuffs from the Borod Basin (in Romanian, in original: Contribuții la caracterizarea petrografică a tufurilor vulcanice din bazinul Borod). *Studia Univ. Babeş-Bolyai*, Ser. Geol.-Geogr., **XXXI**, 1, 67-77.
35. **Ghergari, Lucreția**, Nicolescu, Șt., Cociuba, M. (1986) Manjiroite and todorokite from Ocna de Fier, Romania. *Studia Univ. Babeş-Bolyai*, Ser. Geol.-Geogr., **XXXI**, 2, 3-12.
36. Cozubaș, M., **Ghergari, Lucreția**, Mârza, I. (1986) Hydrothermal transformation (argillic stage) of the volcanics from the Cărnic and Cetate massifs from Roșia Montană (Alba district). [in Romanian, in original: Transformarea hidrotermală (faza argilitică) a vulcanitelor din masivele Cărnic și Cetate de la Roșia Montană (jud. Alba)]. *Crisia*, **XVI**, 539-551, Oradea.
37. **Ghergari, Lucreția**, Mészáros, N., Zotoiu, B. (1987) Recherches minéralogiques et pétrographiques sur les formations de la série marine inférieure de Leghia (Département de Cluj). In vol. The Eocene from the Transylvanian Basin, 255-260, Cluj-Napoca.
38. Mureșan, I., Bodea, I., Bedelea, I., **Ghergari, Lucreția**, Mârza, I. (1987) Caractérisation minéralogico-pétrographique de quelques andésites de la région de Baia Mare et de leurs coulées recristallisées; *Studia Univ. Babeş-Bolyai*, Ser. Geol.-Geogr., **XXXII**, 1, 35-44.
39. Mureșan, I., **Ghergari, Lucreția**, Mârza, I., Bedelea, I. (1987) Le complexe bariolé inférieur paléocène de la région Iara-Huedin-Hodișu, département de Cluj, contributions minéralogiques-pétrographiques; *Studia Univ. Babeş-Bolyai*, Ser. Geol.-Geogr., **XXXII**, 1, 21-33.
40. **Ghergari, Lucreția**, Nicolescu, Șt. (1987) Tibiscumit - a new mineral of the smectites group; *Studia Univ. "Babeş-Bolyai"*, Ser. Geol.-Geogr., **XXXII**, 2, 29-40.
41. Constantiniuc, Valeria, Constantiniuc, E., Mârza, I., **Ghergari, Lucreția**, Bojin, D. (1987) La présence des tellures dans les gisements polymétalliques néogènes de Rodna Veche (Monts de Rodna); *Rev. Roum. Géol. Géophys. Géogr.*, Ser. Géol., **31**, 15-22.
42. **Ghergari, Lucreția**, Bucur, I.I. (1988) L'étude minéralogico-pétrographique de certains dépôts argileux liasiques du compartiment médian de la zone de Reșița - Moldova Nouă (Carpathes Méridionales); *Studia Univ. Babeş-Bolyai*, Ser. Geol.-Geogr., **XXXIII**, 1, 29-38.

43. Stoicovici, E., **Ghergari, Lucreția**, Bedelea, I., Popa, M., A., Treiber, I., Imreh, I., Hoțiu, I., Băluță, C., Gábos, Z., Fuchs, H., Voiculescu, L., D. (1988) Etude minéralogique des sables quartzeux-kaoliniteux de l'extension de la Carrière Cornești - Aghireș; *Studia Univ. Babeş-Bolyai*, Ser. Geol.-Geogr., **XXXIII**, 1, 57-68.
44. Mârza, I., **Ghergari, Lucreția**, Mirea, O. (1988) On the presence of celadonite and analcite-bearing cinerites (Dej tuff) in the salt deposit from Ocna Dej (Romania); *Studia Univ. Babeş-Bolyai*, Ser. Geol.-Geogr., **XXXIII**, 2, 77-83.
45. **Ghergari, Lucreția**, Mârza, I., Mészáros, N. (1989) Lithologic study on the Merian Stratotype (Mera, Cluj district); In vol. The Oligocene from the Transylvanian Basin, Romania, 55-66, Cluj-Napoca.
46. Mureșan, I., **Ghergari, Lucreția**, Bedelea, I., Hoțiu, I. (1989) Contributions à l'étude des sables quartzeux-kaoliniteux de Sânpaul Ouest et de Dealul Varului, département de Cluj; Geological Formations of Transylvania, Romania. In vol.: The Oligocene from the Transylvanian Basin, Romania, 387- 424, Cluj-Napoca.
47. **Ghergari, Lucreția**, Mureșan, I., Ivan, I., Bengescu, M. (1989) Jarosite and römertite within the Oligocene quartz and kaolin-bearing sands in the Șard-Mihăiești area (Cluj district). In vol.: The Oligocene from the Transylvanian Basin, Romania, 451-458, Cluj-Napoca.
48. **Ghergari, Lucreția**, Petrescu, I., Tudoran, V. (1989) Mineralogical-petrographical studies on the Oligocene rocks of the Curtuiș, Buzuș and Ileanda beds with the view to reconstituting the paleoenvironment. In vol.: The Oligocene from the Transylvanian Basin, Romania, 459-468, Cluj-Napoca.
49. **Ghergari, Lucreția**, Mârza, I., Ivan, I., Nicolescu, S. (1989) Zeolite-bearing microlites within the banatitic granites on Valea Drăganului (the Apuseni Mountains); *Studia Univ. Babeş-Bolyai*, Ser. Geol.-Geogr., **XXXIV**, 1, 31-45.
50. **Ghergari, Lucreția**, Dan, T. (1989) The presence of metalaumontite in the Băiuț ore deposit (Maramureș County), *Studia Univ. Babeş-Bolyai*, Ser. Geol.-Geogr., **XXXIV**, 1, 47-52.
51. Mârza, I., **Ghergari, Lucreția**, László, K. (1989) Liesegang structures (hydrothermal-metasomatic) in the pebbles of Badenian conglomerates from polymetallic ore deposit at Băia-Coranda (Hondol, the Metalliferous Mts.). *Studia Univ. Babeş-Bolyai*, Ser. Geol.-Geogr., **XXXIV**, 1, 87-96.
52. Mârza, I., Mészáros, N., **Ghergari, Lucreția**, Bedelea, I., Șuraru, N. (1989) Excursion guidebook. Symp. "Volcanic tuffs in Transylvania" (in Romanian, in original: Ghidul excursiei, Simpozionul Tufurile vulcanice din Transilvania), Univ. Babeş-Bolyai Cluj-Napoca, 18-33, oct. 1989.
53. Mârza, I., **Ghergari, Lucreția**, Nicolescu, S. (1990) Mushketovization in Romanian ore deposits; *Studia Univ. Babeş-Bolyai*, Ser. Geol.-Geogr., **XXXV**, 1, 57-67.
54. **Ghergari, Lucreția**, Hosu, Al., Ivan, I. (1990) Mineralogical composition of the lower Badenian sediments al Lăpuș de Sus (Hunedoara county) - paleoenvironment indicator; *Studia Univ. Babeş-Bolyai*, Ser. Geol.-Geogr., **XXXV**, 2.
55. **Ghergari, Lucreția**, Oniciu, L., Mureșan, L., Pîntea, A., Topan, V., A., Ghițoiu, D. (1991) Effect of additives on the morphology of lead electrodeposits; *J. Electroanal. Chem.*, **313**, 303-311.
56. **Ghergari, Lucreția**, Nicolescu, S., Torok, I. (1991) Mineralogy and petrography of the Balta-Sărată volcanic tuff (Caraș-Severin county). In vol. The volcanic tuffs from the Transylvanian Basin, Romania, 325 -330, Cluj-Napoca.
57. **Ghergari, Lucreția**, Mészáros, N., Mârza, I., Chira, C., Filipescu, S., Ivan, I. (1991) Contributions to the petrographic and chronostratigraphic knowledge of the tuffs in the Cojocna area. In vol. The Volcanic Tuffs from the Transylvanian Basin, Romania, 207-216, Cluj-Napoca.
58. **Ghergari, Lucreția**, Bedelea, I., Codrea, V., Hosu, Al., Bereckmeri, L. (1991) Additional data on the lithological nature of the "volcanic tuffs" (Liassic) at Șuncuiș (Pădurea Craiului Mountains). In vol.: The volcanic tuffs from the Transylvanian Basin, Romania, 357-364, Cluj-Napoca.
59. Mârza, I., **Ghergari, Lucreția**, Mészáros, N., Nagy, A., Băciu, C., Ivan, I., Chezan, S. (1991) L'étude du tuf de Dej dans son extrémité occidentale d'affleurement (Cluj-Napoca-Mera-lara-Borzești) - Bassin de Transylvanie. In vol.: The volcanic tuffs from the Transylvanian Basin, Romania, 183 - 190, Cluj-Napoca.
60. **Ghergari, Lucreția**, Mârza, I., Bedelea, I. (1991) Phenomenes d'alteration supergene associées aux tufs volcaniques du Bassin de Transylvanie. In vol.: Geological Formations of Transylvania, Romania, 3. The Volcanic Tuffs from the Transylvanian Basin, Romania, 293 - 302, Cluj-Napoca.
61. **Ghergari, Lucreția**, Mârza, I., Gherman, Monica (1991) New evidence on the presence of cineritic material in the salt deposits of Ocna Dej (Cluj district). In vol.: The volcanic tuffs from the Transylvanian Basin, Romania. 201-206, Cluj-Napoca.
62. Mârza, I., Anastasiu, N., Seghedi, I., Szakács, Al., Kovacs, M., **Ghergari, Lucreția**, Bedelea, I., Șeclăman, M., Nicolescu, St., Ștefănescu, N., Voiculescu, L. (1991) On the nomenclature and classification of pyroclastites. In vol.: The volcanic tuffs from the Transylvanian Basin, Romania. 447-464, Cluj-Napoca.
63. **Ghergari, Lucreția**, Mészáros, N., Hosu, Al., Filipescu, S., Chira, Carmen (1991) The gypsiferous formation at Cheia (Cluj county). *Studia Univ. Babeş-Bolyai*, Ser. Geol.-Geogr., **XXXVI**, 1, 13-28.

64. **Ghergari, Lucreția**, Strusievicz, R., O. (1991) Low-temperature quartz from Baia Sprie and spatial distribution of the crystallographic forms; *Studia Univ. Babeş-Bolyai, Ser. Geol.-Geogr.*, **XXXVI**, 1, 41-47.
65. **Ghergari, Lucreția**, Strutinski, C., Nicolescu, Șt. (1991) Petrometallogenetical data on the Crystalline Formations from the Ic Ponor Area (Apuseni Mountains). I. The petrogenesis of Metamorphic rocks; *Studia Univ. Babeş-Bolyai, Ser. Geol.-Geogr.*, **XXXVI**, 1, 29-40.
66. **Ghergari, Lucreția**, Nica, D., B., Bedeleian, H., Hobincu, R. (1991) The petrography and mineralogy of the X-th coal-bed bearing formation (Lower Romanian) from the Jilț - Cojmănești Area (Oltenia); *Studia Univ. Babeş-Bolyai, Ser. Geol.-Geogr.*, **XXXVI**, 1, 49-56.
67. Mârza, I., **Ghergari, Lucreția**, Bodea-Popa, M., Dordea, D. (1991) Red granitoid elements in some Miocene conglomerates from the Transylvanian Basin and the "Hidden Mountains" (in Romanian, in original: Galeții de granitoide roșii din unele conglomerate Miocene din Bazinul Transilvaniei și "Munții Ascunși"). *Analele St. Univ. "Al. I. Cuza" Iași*, **XXXVII**, s. II-b, Geol., 69-78.
68. Mârza, I., **Ghergari, Lucreția** (1992) L'adulairisation des volcanites de Roșia Montană (Massif Cetate - Carpates Occidentales). *Rev. Roum. Geol. Geophys. Geogr.*, Ser. Geol., **36**, 15-23, București.
69. **Ghergari, Lucreția**, Mârza, I., Bodolea, A., Schiau, S. (1992) Observations cristallographiques et génétiques sur les mégascalénoèdres de calcite de la Grote de Valea Firii (Monts de Bihor, Roumanie); *Karstologia*, **20**, 49-53.
70. Mârza, I., Bedeleian, I., **Ghergari, Lucreția**, Mureșan, I., Ianoliu, C. (1992) Détermination des minéraux argileux contenus dans les échantillons des roches carottées de quelques structures gazéifères de Transylvanie (I). *V-a Conf. Naț. a Gr. Român pentru Studiul Argilelor*, Vol. sp., 99-121, București.
71. **Ghergari, Lucreția**, Onac, B., P. (1992) Crisite - a new mineral species found in the Bolhac Cave (Pădurea Craiului Mountains, Romania). *Proceedings of the European Conference on speleology*; vol. 2, Physical Speleology and Karstology, Helecine, Belgia.
72. **Ghergari, Lucreția**, Nicolescu, Șt., Strutinski, C. (1992) Petrometallogenetical data on the Crystalline Formations from the Ic Ponor Area (Apuseni Mountains). II. Metallogeny of metamorphic rocks. *Studia Univ. Babeş-Bolyai, Ser. Geol.-Geogr.*, **XXXVII**, 1, 4-11.
73. Mârza, I., **Ghergari, Lucreția**, Plăceanu-Marian, L., Marian, D. (1992) Note sur le quartz exogène du conglomérat de Râpa Dracului (Badénien), situé sous le tuf de Dej, dans la localité Vale (Département de Cluj). *Studia Univ. Babeş-Bolyai, Ser. Geol.-Geogr.*, **XXXVII**, 1, 13-21.
74. **Ghergari, Lucreția**, Lenart, C., Mârza, I., Pop, D. (1992) Anorthitic composition of plagioclases, criterion for parallelizing volcanic tuff horizons in the Transylvanian Basin). *Studia Univ. Babeş-Bolyai, Ser. Geol.-Geogr.*, **XXXVII**, 1, 31-40.
75. Mureșan, I., **Ghergari, Lucreția**, Bedeleian, I. (1992) The geology, mineralogy and rheological properties of quartz-kaolinitic sands from Popești (Cluj district) (in Romanian, in original: Geologia, mineralogia și proprietățile reologice ale nisipurilor cuarțoase-caolinoase de la Popești (județul Cluj)). *Studia Univ. Babeş-Bolyai, Ser. Geol.-Geogr.*, **XXXVII**, 1, 63-81.
76. Mészáros, N., **Ghergari, Lucreția**, Strusievicz, E. (1992) Contributions to the knowledge of the lithology and stratigraphy of the Miocene deposits from the Ocna Mureș zone (Transylvanian Basin); *Studia Univ. Babeş-Bolyai, Ser. Geol.-Geogr.*, **XXXVII**, 2, 64-70.
77. Mészáros, N., **Ghergari, Lucreția**, Chira, C. (1992) The Badenian deposits at Borzești; *Studia Univ. Babeş-Bolyai, Ser. Geol.-Geogr.*, **XXXVII**, 2, 71-78.
78. Mârza, I., **Ghergari, Lucreția**, Nicolescu, Șt., (1992) Les implications du phénomène d'impact météoritique sur l'épicroute terrestre; *Studia Univ. Babeş-Bolyai, Ser. Geol.-Geogr.*, **XXXVII**, 2, 83-90.
79. **Ghergari, Lucreția**, Strusievicz, R., O., Frățilă, Gh., Sîntămărian, A. (1993) Crystallogenesis of Calcite from Piatra Altarului Cave, (Bihor Mts., Romania). *Rom. Jour. Mineral.*, **76**, 1, 87-96.
80. **Ghergari, Lucreția**, Hanga, M. (1993) Contribution to the knowledge of copper acetates [in Romanian, in original: Contribuții la cunoașterea acetaților de cupru (II)]. *Conf. Naț. Chimie și Ing. Chimică, București*, 29-31 oct. 1993.
81. Mârza, I., **Ghergari, Lucreția**, Mărieș, M. (1993) Procès pneumatolytiques-hydrothermaux associés aux apogranites de Valea Drăganului (Monts Vlădeasa); *Studia Univ. Babeş-Bolyai, Ser. Geol.-Geogr.*, **XXXVIII**, 1, 7-22.
82. Mureșan, I., **Ghergari, Lucreția**, Bedeleian, I. (1993) The mineralogy and rheology of quartz-kaolinitic sands from the Stoguri-Aghireș perimeter (Cluj district) (in Romanian, in original: Mineralogia și reologia nisipurilor cuarțoase-caolinoase din perimetrul Stoguri - Aghireș, județul Cluj). *Studia Univ. Babeş-Bolyai, Ser. Geol.*, **XXXVIII**, 1, 43-62.
83. **Ghergari, Lucreția**, Bedeleian, I., Denuț, I. (1993) Some aspects concerning the mineralogy of metamorphites from the Catarama-Ivășcoaia perimeter (the Maramureș Mountains) [in Romanian, in original: Câteva aspecte privind mineralogia metamorfitelor din perimetrul Catarama-Ivășcoaia (Munții Maramureșului)]. *Studia Univ. Babeş-Bolyai, Ser. Geol.*, **XXXVIII**, 1, 75-80.

84. **Ghergari, Lucreția**, Onac, B. P. (1993) Crisite - a new mineral species found in the Bolhac Cave (Pădurea Craiului Mountains, Romania). *Bulletin de la Soc. Geogr. de Liege*, **29**, 97-104.
85. **Ghergari, Lucreția** (1993) A review: Mârza I. - The genesis of the deposits of magmatic origin – vol. III (in Romanian, in original; Recenzie: Mârza I. - Geneza zăcămintelor de origine magmatică, vol. III, Edit. Dacia, Cluj-Napoca), 382 p., 1992). *Studia Univ. Babeş-Bolyai*, Ser. Geol., **XXXVIII**, 1, 108.
86. **Ghergari, Lucreția**, Onac, B., P., Sîntămărian, A. (1994) Mineralogy of moonmilk formation in some Romanian and Norwegian caves. *Theoretical and Applied Karstology*, **6**/1993, 107-120, Bucureşti.
87. Mészáros, N., Mârza, I., **Ghergari, Lucreția**, Chira, C., Toth, L. (1994) Contributions to the stratigraphy of the Lueta - Mereşti region (the Herghita county) and to the age of the volcanic tuffs. In vol.: The Miocene from the Transylvanian Basin, Romania. 171-178.
88. Silvestru, E., **Ghergari, Lucreția** (1994) On the paleokarst in the cave Gheţarul de la Scărişoara (Bihor Mountains, Romania). *Theoretical and Applied Karstology*, **7**/1994, 155-161, Bucureşti.
89. **Ghergari, Lucreția**, Strusievicz, R.O., Dumitrescu, S. (1994) Minamiite in the hydrothermal alteration zone of the Fâncel-Lăpuşna caldera (Gurghiu Mts., East Carpathians): first record in Romania. *Studia Univ. Babeş-Bolyai*, Ser. Geol., **XXXIX**, 1-2, 93-103.
90. Bedeleian, I., **Ghergari, Lucreția**, Bedeleian, H. (1994) Hypergenetic processes in the rock of magmatic origin from the Şimleu Basin. *Studia Univ. Babeş-Bolyai*, Geol.-Geogr., **XXXIX**, 1-2, 105-114.
91. Mârza, I., **Ghergari, Lucreția**, Har, N. (1994) Contribution to the knowledge of lithospheric enclaves included in Pliocene - Quaternary alkaline basalts from the Perşani Mountains. *Studia Univ. Babeş-Bolyai*, Geologia, **XXXIX**, 1-2, 115-125.
92. **Ghergari, Lucreția**, Forray, F., Gál, A., Fărcaş, T. (1994) Arsenic minerals from Săcărâmb ore deposit: arsenic, arsenolite, villyaellenite and krautite (Transylvania, Romania). *Studia Univ. Babeş-Bolyai*, Ser. Geol., **XXXIX**, 1-2, 127-140.
93. **Ghergari, Lucreția**, Chira, C. (1994) A new outcrop of Eggenburgian Deposits in the NW of the Transylvanian Basin: Borzeşti (Cluj County). *Studia Univ. Babeş-Bolyai*, Ser. Geol., **XXXIX**, 1-2, 263-268.
94. Mârza, I., **Ghergari, Lucreția**, Ionescu, C., Har, N. (1995) The presence of some calcite pseudomorphs after Aragonite in Cacova Ierii Deposit, Băişoara, Apuseni Mts.; *Rom. Jour. Mineral.*, **76**, 2, 85-90.
95. **Ghergari, Lucreția**, Nicolescu, Şt., Mârza, I., Gruescu, C. (1995) Microchemical data on andradite garnets with inclusions from Ocna de Fier, Banat, Romania. *Rom. Jour. Mineral.*, **76**, 2, 95-100.
96. Onac, B., **Ghergari, Lucreția**, Gál, A. (1995) Crystallographical studies on gypsum crystals in Ponoraş Cave (Pădurea Craiului Mts., Romania); *Theoretical and Applied Karstology*, **8**, 63-68, Bucureşti.
97. **Ghergari, Lucreția**, Onac, B. (1995) The crystallogenesis of gypsum flowers. *Cave and Karst Science*, **22** (3), 119-122.
98. Mârza, I., **Ghergari, Lucreția**, Forray, F., Tămaş, C. (1995) Glauch-glamm formation, associated to the hydrothermal deposits from Apuseni Mountain - genetic and metallogenetic mechanism. *Studia Univ. Babeş-Bolyai*, Ser. Geol., **XL**, 1, 185-194.
99. **Ghergari, Lucreția**, Lazo, C. (1996) The mineralogy of Refractory Clays from Şuncuiuş (Bihor Mountains, Romania). *Studia Univ. Babeş-Bolyai*, Ser. Geol., **XLI**, 1, 145-164.
100. **Ghergari, Lucreția**, Forray, F., Andrei, V. (1996) Contributions to the Petrographic and Mineralogical Study of the Rock Salt from Cacica (Suceava County). *Studia Univ. Babeş-Bolyai*, Ser. Geol., **XLI**, 1, 165-180.
101. Gorea, M., Gagea, L., **Ghergari, Lucreția** (1997) The study of ceramic BaTiO₃ with BaSnO₃, BaZrO₃ and Sb₂O₃ microstructures synthesized beforehand (in Romanian, in original: Studiul microstructurii ceramicii de BaTiO₃ cu BaSnO₃, BaZrO₃ şi Sb₂O₃ sintetizaţi în prealabil). *Ses. Comunic. Ştiinţ. ale Univ. "Aurel Vlaicu", Arad*, Ediţia a IV-a, 30-31 oct. 1997, vol. III, Fizică-Chimie, 165-171, Arad.
102. **Ghergari, Lucreția**, Onac, B., Frăţilă, G. (1997) Mineralogy of crusts and efflorescences from Humpleu cave system. *Proceedings of the 12th International Congress of Speleology*, I, 231-234, La Chaux de Fonds, Switzerland, 10-17 August 1997.
103. Vieman, I., **Ghergari, Lucreția**, Onac, B. (1997) Crystallographical observations on calcite rafts from three Romanian Cave. *Proceedings of the 12th International Congress of Speleology*, I, 227-230, La Chaux de Fonds, Switzerland, 10-17 August 1997.
104. **Ghergari, Lucreția** (1997) A review: B. P. Onac - Stalagmitic formations in the caves of Pădurea Craiului (in Romanian, in original: Recenzie: Bogdan-Petroniu Onac - Formaţiuni stalagmitice în peşterile Pădurii Craiului, Edit. Acad., Bucureşti, 175 p. 1997). *Natura Silvaniae*, **1**, 235, Jibou.
105. **Ghergari, Lucreția**, Tămaş, T., Damm, P., Forray, F. (1997) Hydrothermal paleokarst in Peştera Valea Rea (Bihor Mountains, Romania). *Theoretical and Applied Karstology*, **10**/1997, 105-114, Bucureşti.
106. Mârza, I., Tămaş, C., **Ghergari, Lucreția** (1997) Low Sulfidation Epithermal Gold Deposits from Roşia Montană, Metaliferi Mountains, Romania. *St. Cerc. Geol., Geofiz. Geogr.*, Ser. Geol., **42**, 3-12.
107. **Ghergari, Lucreția**, Gorea, M. (1997) The influence of microstructure and phase composition on the dielectric properties of some ceramics systems. I. BaTiO₃ - TiO₂ ± BaZrO₃, BaSnO₃, Sb₂O₃. *Studia Univ. Babeş-Bolyai*, **XLII**, Ser. Geol., 27-42.

108. **Ghergari, Lucreția**, Tarkany, F., Goronea, T. (1997) Mineralogical and petrographic studies on the building materials of the XV-th century church from Lupșa (Alba County). *Studia Univ. Babeş-Bolyai, Ser. Geol.*, **XLII**, 1, 105-117.
109. Constantin, D.M., **Ghergari, Lucreția**, Rus, E.M., Oniciu, L. (1997) Characteristics of the active anodic mass from the alkaline Ni-Cd accumulator (in Romanian, in original: Caracteristici ale masei active anodice din acumulatorul alcalin Ni-Cd). *Simp. Știința Modernă și energia; Producerea, transportul și utilizarea energiei*, Ediția a XVI-a, 195-200, mai 1997, Cluj-Napoca.
110. Constantin, D.M., **Ghergari, Lucreția**, Rus, E.M., Oniciu, L. (1997) Characteristics of the active cathodic mass from an alkaline Ni-Cd accumulator (in Romanian, in original: Caracteristici ale masei active catodice din acumulatorul alcalin Ni-Cd). *Simp. Știința Modernă și energia; Producerea, transportul și utilizarea energiei*, Ediția a XVI-a, 201-206, mai 1997, Cluj-Napoca.
111. **Ghergari, Lucreția** (1998) A review: B. P. Onac - Stalagmitic formations in the caves of Pădurea Craiului (in Romanian, in original: Recenzie: Bogdan-Petroniu Onac - Formațiuni stalagmitice în peșterile Pădurii Craiului, Edit. Acad., București, 175 p. 1997). *Speomond*, **3**, 40, Cluj-Napoca.
112. **Ghergari, Lucreția**, Onac, B. P., Vremir, M., Strusievicz, R. (1998) La cristallogenèse des spéléothèmes de la grotte Lithophagus (Monts Pădurea Craiului, Roumanie); *Karstologia*, **31**, 19-26, Paris.
113. Constantin, D.M., Rus, E.M., Oniciu, L., **Ghergari, Lucreția** (1998) The influence of some additives on the electrochemical behaviour of sintered nickel electrodes in alkaline electrolyte. *Journal of Power Sources*, **74/2**, 188-197, Lausanne, Switzerland.
114. Gorea, Maria, **Ghergari, Lucreția** (1998) The obtaining and characterisation of BaTiO₃ used as raw material for Dielectric Ceramics (in Romanian, in original: Obținerea și caracterizarea BaTiO₃ utilizat ca materie primă pentru ceramica dielectrică. *Studia Univ. Babeş-Bolyai, Ser. Geol.*, **XLIII**, I, 3-13.
115. **Ghergari, Lucreția**, Tămaș, T. (1999) Huntite Formed under Supergene Conditions in Valea Rea Cave (Bihor Mountains). *Rom. Jour. Mineral.*, **79**, 151-158 (pentru 1998), București.
116. Ionescu, C., **Ghergari, Lucreția** (1999) The hydrogarnets - An unsolved problem? *Rom. Jour. Mineral.*, **79**, 101-104, (pentru 1998), București.
117. **Ghergari, Lucreția**, Tămaș, T. (1999) Mineralogy of cave deposits from Bihor Mountains (Romania). *Contribución del estudio científico de las cavidades kársticas al conocimiento geológico*. B. Andreo, F. Carrasco y J.J. Durán (Eds.), 243-255, Nerja (Málaga).
118. **Ghergari, Lucreția**, Gorea, M. (1999) The influence of microstructure and phase composition on dielectric properties of some ceramics. *VIth ECERS-Conference & Exhibition of the European Ceramic Society*, Vol. I, **60**, 97-98, London.
119. Ozunu, A., Literat, L., Gagea, L., **Ghergari, Lucreția** (1999) Kinetic models of dehydrated gypsum building materials (in Romanian, in original: Modele cinetice ale deshidratării gipsului). *Materiale de construcție*, **XXIX**, 1, 34-38.
120. **Ghergari, Lucreția**, Lazarovici, Gh., Ionescu, C., Tămaș, T. (1999) Geoarchaeological studies on some Romanian ceramic artefacts from the Early Neolithic Lunca – Poiana Slatinii site, Neamț District (in Romanian, in original: Studii geoarheologice asupra unor artefacte ceramice din Neoliticul Timpuriu din România: Stațiunea de la Lunca - Poiana Slatinii, județul Neamț). *Angvstia*, **4**, 1-7, Sfântu Gheorghe.
121. Boboș, I., **Ghergari, Lucreția** (1999) Conversion of smectite to ammonium illite in the hydrothermal system of Harghita Băi, Romania: SEM and TEM investigations. *Geologica Carpathica*, **50**, 5, 379-387.
122. **Ghergari, Lucreția**, Gorea, M., Tămaș, T., Susan, I., Șamșudean, C. (1999) A study on the microstructures and defects of household porcelain. *Studia Univ. Babeş-Bolyai, Ser. Geol.*, **XLIV**, I, 39-48.
123. Tămaș, T., Manolache, E., **Ghergari, Lucreția**, Drăgan-Bularda, M. (2000) Mineralogy and Microbiology of black deposits from Cuciulata Pothole (Bihor Mountains, Romania). In vol. "Proceedings of the joint meeting of Friends of Karst Theoretical and Applied Karstology, and IGCP 448", Edited by Bogdan P. Onac & Tudor Tămaș, 140-147, July 14-22, 2000.
124. **Ghergari, Lucreția**, Ionescu, C., Rusu, V., Gorea, M. (2000) Mineralogical considerations on Roman ceramics from Napoca archaeological site (in Romanian, in original: Considerații mineralogice asupra ceramicii romane din situl arheologic Napoca). *CONSILOX VIII*, vol. I, 115-122, Alba Iulia.
125. **Ghergari, Lucreția**, Ionescu, C., Gorea, M., Țentea, O., Toadere, M. (2000) The provenance of raw materials used for Dacic ceramics from the Roman site of Gilău (in Romanian, in original: Proveniența materiilor prime utilizate pentru ceramica dacică din situl roman Gilău). *CONSILOX VIII*, vol. I, 123-129, Alba Iulia.
126. **Ghergari, Lucreția**, Gorea, M., Ionescu, C., Zoltan, I. (2000) Processes at the glaze-ceramic body interface in different types of ceramics (in Romanian, in original: Procese de interfață glazură-ciob în diferite tipuri de ceramică). *CONSILOX VIII*, vol. I, 37-44, Alba Iulia.
127. Ionescu, C., **Ghergari, Lucreția**, Rusu, V., Gorea, M. (2000) Microscopic aspects of the Roman coarse ceramics from the archaeological site of Napoca (in Romanian, in original: Aspecte microscopice ale

- ceramicii grosiere romane din situl arheologic Napoca). In vol. Achievements and perspectives in the Quaternary studies in Romania. Univ. Babeş-Bolyai Cluj-Napoca, 48-50, Cluj-Napoca.
128. Ionescu, C., **Ghergari, Lucreția**, Tămaş, T. (2000) New data on the Neolithic ceramics from some Romanian archaeological sites. In vol. Achievements and perspectives in the Quaternary studies in Romania. Univ. Babeş-Bolyai Cluj-Napoca, 51-52, Cluj-Napoca.
 129. **Ghergari, Lucreția**, Ionescu, C. (2000) Roman fine ceramics from the Napoca site: mineralogical and geoaechaeological considerations (in Romanian, in original: Ceramica fină romană din situl Napoca: considerații mineralogice și geoarheologice). In vol. Achievements and perspectives in the Quaternary studies in Romania. Univ. Babeş-Bolyai Cluj-Napoca, 53-54, Cluj-Napoca.
 130. **Ghergari, Lucreția**, Onac, B. P. (2000) Mineralogy of clay horizons from within the peat deposit of Preluca Țiganului (Munții Gutâi). In vol. Achievements and perspectives in the Quaternary studies in Romania. Univ. Babeş-Bolyai Cluj-Napoca, 56-58, Cluj-Napoca.
 131. Ionescu, C., **Ghergari, Lucreția** (2000) Gem accumulations in Quaternary deposits from Romania (in Romanian, in original: Acumulări de geme în depozite cuaternare din România.) In vol. Achievements and perspectives in the Quaternary studies in Romania. Univ. Babeş-Bolyai Cluj-Napoca, 63-65, Cluj-Napoca.
 132. **Ghergari, Lucreția**, Ionescu, C., Tămaş, T., Onac, B. P. (2000) The weathering of clay minerals in early Holocene soil in north-western Romania. In vol. Achievements and perspectives in the Quaternary studies in Romania. Univ. Babeş-Bolyai Cluj-Napoca, 59-60, Cluj-Napoca.
 133. Tămaş, T., Causse, C., Blamart, D., von Grafenstein, U., **Ghergari, Lucreția** (2000) U-Th TIMS dating and stable isotope analyses on speleothems from V11 Cave (Bihor Mountains, NW Romania). In vol. Achievements and perspectives in the Quaternary studies in Romania. Univ. Babeş-Bolyai Cluj-Napoca, 17-19, Cluj-Napoca.
 134. **Ghergari, Lucreția**, Ionescu, C., Codrea, V (2000) Zircon, Ti-minerals with Ti content in the Quaternary deposits related with weathered-granodiorite environment. In vol.: Natural and technogenic placer and weathered rock deposits at the turn of the Millennium, 63-65, *Russian Academy of Science*, Moscow.
 135. Ionescu, C., **Ghergari, Lucreția** (2000) Brucite-bearing deposits in NW Romania: genetical considerations. *Polskie Towarz. Mineralog.-Sp. vol.*, **17**, 167-170, 2000, Krakow, Polonia.
 136. Ionescu, C., **Ghergari, Lucreția** (2000) The mineralogy of the brucite from predazzites in the Apuseni Mountains (NW Romania). *Polskie Towarz. Mineralog., Sp. vol.*, **17**, 171-174, Krakow, Polonia.
 137. **Ghergari, Lucreția**, Ionescu, C., Balintoni, I., Indrieş, R. (2000) Contact phenomena associated to quartzo-feldspatic injections in marbles and other metamorphic rocks from NW Romania. *Polskie Towarz. Mineralog., Sp. vol.*, **17**, 141-143, Krakow, Polonia.
 138. **Ghergari, Lucreția**, Țentea, O., Marcu, F. (2000) Mineralogical aspects of hand-made ceramics from the Roman castle of Gilău (in Romanian, in original: Aspectele mineralogice ale ceramicii lucrate cu mâna din castrul roman de la Gilău). *Acta Musei Apulensis*, **XXXVII/1**, 401-416, Alba Iulia.
 139. **Ghergari, Lucreția**, Ionescu, C. (2000) The hydrograndite and magnesioferite in the Budureasa Area (Apuseni Mountains, Romania): genetical implications. *N. Jb. Miner. Mh.*, **11**, 481-495. Stuttgart, Germania.
 140. **Ghergari, Lucreția**, Ionescu, C. (2000) The application of mineralogical methods of analysis in archaeology (in Romanian, in original: Aplicații ale metodelor mineralogice de analiză în arheologie). *Banatica*, **15**, I, 261-270, Reșița.
 141. **Ghergari, Lucreția**, Ionescu, C., Mariş, C. (2000) New data regarding the mineralogical compositions of the clayish silt from Dumbrava, Cluj District (in Romanian, in original: Date noi privind compoziția mineralogică a siltului lutitic de la Dumbrava, Județul Cluj). *Studia Univ. Babeş-Bolyai*, Ser. Geol., **XLV**, I, 23-33.
 142. **Ghergari, Lucreția**, Onac, B.P. (2001) Late Quaternary paleoclimate reconstruction based on clay minerals association from Preluca Țiganului (Gutâi Mountains, Romania). *Studia Univ. Babeş-Bolyai*, Ser. Geol., **XLVI**, I, 15-28.
 143. Wohlfarth, B., Hannon, G., Feurdean, A., **Ghergari, Lucreția**, Onac, B.P., Possnert, G. (2001) Reconstruction of climatic and environmental changes in NW Romania during the early part of the last deglaciation (~15,000-13,600 cal yr BP). *Quaternary Science Reviews*, **20**, 1897-1914.
 144. Ionescu, C., **Ghergari, Lucreția** (2001) SiO₂-gems associated to Mesozoic magmatites in Buru-Cheile Turzii area (Apuseni Mts, Romania). *Polskie Towarz. Mineralog., Sp. vol.*, **19**, 73-75, Polonia.
 145. **Ghergari, Lucreția**, Ionescu, C. (2001) Wall-rock xenoliths in Upper Cretaceous-Lower Paleogene granodiorites of Vlădeasa tephrolite (Apuseni Mts., Romania). *Pol. Tow. Mineral. Prace. Spec.*, **19**, 49-51, Krakow, Polonia.
 146. Ionescu, C., **Ghergari, Lucreția** (2001) Hydrothermal transformations of phlogopite: a case study from the Apuseni Mountains (Romania). *Mineralogia Polonica*, **32**, 1, 27-34, Polonia.

147. Ionescu, C., **Ghergari, Lucreția** (2002) Modeling and firing technology – reflected in the textural features and the mineralogy of the ceramics from Neolithic sites in Transylvania (Romania). *Geologica Carpathica*, **53**, (CD), Bratislava.
148. Tertîş, L., Bucur, I., Săsăran, E., **Ghergari, Lucreția** (2002) New remarks on the lithology and biostratigraphy of the Cretaceous deposits from Istria depression (Black sea shelf). *Studia Univ. Babeş-Bolyai*, Ser. Geol., **XLVII**, 2, 75-84.
149. Lazarovici, Gh., **Ghergari, Lucreția**, Ionescu, C. (2002) Ceramic artifacts from the Middle Neolithic in Transylvania. CCTLNI culture from Zau archeological site [in Romanian, in original: Artefacte ceramice din Neoliticul mijlociu în Transilvania: cultura CCTLNI din stațiunea Zau (județul Mureș)]. *Angvstia*, Ser. Arheologie, **7**, 7-18.
150. Ionescu, C., **Ghergari, Lucreția** (2002) Gemological identification of the components of a thread of beads from the V-VIth Centuries, found in Cordoş Area (Cluj-Napoca, Romania) [in Romanian, in original: Identificarea gemologică a componentelor unui şirag de mărgelă din secolele V-VI, descoperit în zona Cordoş (Municipiul Cluj-Napoca, România)]. *Angvstia*, Ser. Arheologie, **7**, 295-302, 2002.
151. Ionescu, C., **Ghergari, Lucreția** (2002) Methodologies of probing samples from archaeological sites, for the mineralogical, petrographical and pedological analysis (in Romanian, in original: Metodologii de recoltare a eşantioanelor şi probelor mineralogice, petrografice şi pedologice din siturile arheologice). *Angvstia*, Ser. Arheologie, **7**, 351-354, 2002.
152. Mârza, I., **Ghergari, Lucreția**, Ţarină, D. (2002) Nontronite from Lechința, Satu Mare County, *Analele Univ. Bucureşti*, Ser. Geol., Bucureşti, 2002.
153. Balintoni, I., **Ghergari, Lucreția**, Băbuţ, T. (2002) The Arieşeni nappe, or the Moma and Poiana nappes? *Studia Univ. Babeş-Bolyai*, Ser. Geol., **XLVII**, 2, 19-26.
154. Gál, J., **Ghergari, Lucreția** (2003) Pollutant elements in the Săcel Valley – the lower course (the Apuseni Mountains) (in Romanian, in original: Elemente poluante pe valea Săcel - curs inferior (Munții Apuseni)). In vol. Environment & Progress, (Mediul - Cercetare, Protecție şi Gestiune), Univ. Babeş-Bolyai Cluj-Napoca, 245-249.
155. Laczko, A., **Ghergari, Lucreția**, Toth, A. (2003) Sources of pollution in the area of cinnabar ore, at Sântimbru Băi, Harghita District (in Romanian, in original: Surse de poluare în perimetrul mineralizației cinabrifere de la Sântimbru Băi, județul Harghita). In vol. Environment & Progress, (Mediul - Cercetare, Protecție şi Gestiune), Univ. Babeş-Bolyai Cluj-Napoca, 245-249, 2003.

ABSTRACTS

1. **Ghergari, Lucreția**, Ionescu, C., Marincea, Şt. (1998) Sepiolite in the contact aureoles associated with Laramian intrusions from Budureasa, Apuseni Mts. Romania. *XVI Congress CBGA (Carp.-Balk. Geol. Assoc.)*, *Abstr. vol.*, p. 183, Vienna, Austria.
2. **Ghergari, Lucreția**, Ionescu, C. (1999) Zoning in vesuvianite crystal in Vața area (Apuseni Mts., Romania). *Analele Univ. Bucureşti*, Ser. Geologie, **XLVIII**, 24-25, Bucureşti.
3. Mârza, I., **Ghergari, Lucreția**, Ionescu, C., Constantina, C., Bodnariuc, A., Fodor, Şt. (1999) Gemological value of the area between the Ampoi and Geoagiu Valley (Apuseni Mountains). *Rom. Jour. Mineral.*, Suppl. 1, *Abstr. Vol.*, **79**, p. 45, Bucureşti.
4. Ionescu, C., **Ghergari, Lucreția**, (1999) Dypingite - product of the brucite alteration In the Budureasa-Pietroasa Area (Bihar Mts., Romania). *Rom. Jour. Mineral.*, Suppl. 1, *Abstr. Vol.*, **79**, p. 41, Bucureşti.
5. **Ghergari, Lucreția**, Ionescu, C. (1999) Genetic considerations on jasper in the Brad area (Apuseni Mts., Romania). *Rom. Jour. Mineral.*, Suppl. 1, *Abstr. Vol.*, **79**, p. 32, Bucureşti.
6. **Ghergari, Lucreția**, Ionescu, C., Püspöki, Zs. (1999) Jaspis occurrence from Brad, Apuseni Mts. (Romania). *Gems and Gemology*, **XXXV**, 139-140, Carlsbad Ca, USA.
7. Ionescu, Corina, **Ghergari, Lucreția**, Vădan, A. (1999) Gem-quality vesuvianite from Vata, Apuseni Mts. (Romania). *Gems and Gemology*, **XXXV**, p. 139, Carlsbad Ca, USA.
8. **Ghergari, Lucreția**, Onac, B., Ionescu, C., Bodnariuc, A., Onac, L. (1999) The mirroring of the paleoenvironment on the clay minerals in Quaternary deposits from North and West of Romania. In vol. "The environmental background to hominid evolution in Africa". *Book of Abstr.*, p. 68, Durban, South Africa.
9. Ionescu, C., **Ghergari, Lucreția**, (1999) Gems in the Quaternary deposits from Romania: blue quartz from Trestia-Maramureş. In vol. "The environmental background to hominid evolution in Africa". *Book of Abstr.*, 86-87, Durban, South Africa.
10. **Ghergari, Lucreția**, Ionescu, C., Lazarovici, Gh., Tămaş, T. (1999) Geoarchaeological studies of the Neolithic ceramics from Romania: raw materials (source, composition) and techniques of prewheel pottery processing. In vol. "The environmental background to hominid evolution in Africa". *Book of Abstr.*, 67-68, Durban, South Africa.

11. **Ghergari, Lucreția**, Ionescu, C., (2000) Gem-quality materials associated with Laramian magmatites in the Valea Chioarului area, Transylvania (Romania). *31st International Geological Congress*, Abstr. vol.-CD, Rio de Janeiro, Brazilia.
12. Ionescu, C., **Ghergari, Lucreția**, Fărcaș, T. (2000) Gems in the Baia Mare area (Romania). *31st International Geological Congress*, Abstr. vol.-CD, Rio de Janeiro, Brazilia.
13. Filipescu, S., **Ghergari, Lucreția**, Farcaș, T., Ionescu, C. (2000) Paleoenvironment and associated fossils in the Brad jasper (Miocene) from the Southern part of the Apuseni Mountains. *31st International Geological Congress*, Abstr. vol.-CD, Rio de Janeiro, Brazilia.
14. **Ghergari, Lucreția**, Ionescu, C. (2002) Structural and mineralogical changes of raw materials used for the Roman erthen-lamps from Potaissa site (in Romanian, in original: Modificări mineralogice și structurale ale materiilor prime utilizate pentru opaițele romane de la Potaissa). Symp. *Realizări și Persp. în Geol. și Paleont. Rom.* Abstr. vol., Iunie 2002 Cluj-Napoca, p.23.
15. Ionescu, C., **Ghergari, Lucreția** (2002) Neolithic Ceramics from Balta Sărată (Banat) (in Romanian, in original: Ceramica neolitică de la Balta Sărată (Banat): caracteristici mineralogice). Symp. *Realizări și Persp. în Geol. și Paleont. Rom.* Abstr. vol., Iunie 2002 Cluj-Napoca, p. 26.
16. **Ghergari, Lucreția**, Ionescu, C., Toth, A. (2002) The mineralogy of the alteration products of some iron sulphurs from Corund, Harghita District (in Romanian, in original: Mineralogia produșilor de alterare ai unor sulfuri de fier de la Corund, județul Harghita). In vol. *Mediul – cercetare, protecție și gestiune*, Univ. Babeş-Bolyai Cluj-Napoca, p. 49.
17. Ionescu, C., **Ghergari, Lucreția**, Forray, F. (2002) Data concerning the mineralogical composition damp No. 1 from Căpuș (Cluj District) [in Romanian, in original: Date privind compoziția mineralogică a sterilului din iazul nr.1 Căpuș (jud. Cluj)]. In vol.: *Mediul – cercetare, protecție și gestiune*, Univ. Babeş-Bolyai Cluj-Napoca, 56-57.
18. Ionescu, C., **Ghergari, Lucreția**, Culic, A., Grovu, P. (2002) The sterile from the Căpuș damp No. 2 (Cluj District): source of raw materials for building (in Romanian, in original: Sterilul din iazul de decantare nr. 2 Căpuș (jud. Cluj): sursă de materii prime pentru construcții). In vol. > *Mediul – cercetare, protecție și gestiune*, Univ. Babeş-Bolyai Cluj-Napoca, 57-58.
19. Forray, F., **Ghergari, Lucreția**, Ionescu, C. (2002) The mineralogy of the solid material from water stream in the Arieș – Abrud – Roșia area and their role in the metals transport (in Romanian, in original: Mineralogia materialului aflat în suspensie în cursuri de apă din zona Arieș-Abrud-Roșia și rolul acestuia în transportul metalelor). In vol.: *Mediul – cercetare, protecție și gestiune*, Cluj-Napoca, 45-46.
20. Gál, J., **Ghergari, Lucreția** (2002) Polluting elements in the lower course of the Săcel Valley (the Apuseni Mountains) [in Romanian, in original: Elemente poluante în zona cursului inferior al văii Săcel (Munții Apuseni)]. In vol.: *Mediul – cercetare, protecție și gestiune*, Cluj-Napoca Oct. 2002, p. 47.
21. **Ghergari, Lucreția**, Gál, J. (2002) The Mineralogz of Pollutants Formed in the Exploitation Mașca area (the inferior Basin of the Iara Valley) (in Rumanian, in original: Mineralogia produșilor poluanți formați în zona exploatării Mașca (bazinul inferior al văii Iara)). *Mediul – cercetare, protecție și gestiune*, Cluj-Napoca Oct. 2002, p. 48-49.
22. Laczko, A., **Ghergari, Lucreția**, Toth, A. (2002) Sources of pollution in the perimeter of cinabrifere Mineralization in Sântimbru Băi, Harghita District (in Rumanian, in original: Surse de poluare în perimetrul mineralizației cinabrifere de la Sântimbru Băi, județul Harghita). *Mediul – cercetare, protecție și gestiune*, Cluj-Napoca Oct. 2002, 60-61.
23. Stan, O., Ardelean, L., Ionescu, C., **Ghergari, Lucreția**, (2002) Mineralogy of chalcedonies in Căpuș area (Apuseni Mts., Romania). In vol. *4th European Mineralogical Union Symposium on Energy Modelling in Minerals*, Budapesta (Ungaria), 2002.
24. Ardelean, L., Ionescu, C., **Ghergari, Lucreția**, (2002) Chalcedonies, opals and jaspers from Ilba-Seini area (Romania). In vol. *4th European Mineralogical Union Symposium on Energy Modelling in Minerals*, Budapesta (Ungaria), 2002.
25. **Ghergari, Lucreția**, Ionescu, C., Lazăr, C. (2003) The mineralogy of the Neolithic ceramics from Ungurului cave (Șuncuiș, România). *Acta Min.-Petrogr.* Abstr. series, Szeged, **38**, 2003.
26. **Ghergari, Lucreția**, Ionescu, C., Rusu-Bolindeț, V. (2003) Geoarchaeological study on local fine ceramics from II-II century (Napoca site, Romania). *Acta Min.-Petrogr.* Abstr. series, Szeged, **39**, 2003.
27. Tămaș, T., **Ghergari, Lucreția** (2003) Hydronium jarosite from Iza cave (Rodnei Mts., Romania). *Acta Min.-Petrogr.* Abstr. series, Szeged, **39**, p. 102, 2003.

INDEX OF AUTHORS

Balintoni, I.C.	Naud, J. .
Bardocz, Z.	Negoită, V.
Bedelean, H.	Nikolin, B.
Bedelean, I.	O'Connor, G.
Benea, M.	Onac, B.P.
Blănaru, D.C.	Pertlik, F.
Borovikova, E.Yu.	Phetpu, K.
Breban, R.	Pintea, I.
Caracas, R.	Polonic, G.
Cauuet, B.	Pomârleanu, V.
Chirienco, M.	Pop, D.
Ciobanu, L.C.	Puşte, A.
Cook, J.N.	Rădăşanu, S.
Costin, D. .	Radu, D.M.
Damian, F.	Rosic, A.
Damian, Gh.	Sajó, I.
Dărăban, L.	Schoenbeck, T.
Diaconu, G.	Seghedi, I. .
Domnişoru, D.	Self, C.
Dumitraş, D.G.	Simon, V.
Gál, A.	Sreckovic-Batocanin, D.
Gatter, I.	Stan, R.
Ghergari, L.	Stremţan, C.
Gorea, M.	Strutinski, C.
Har, N. .	Stumbea, D.
Hercot, O.	Szabó, Cs.
Hercot, S.	Szakács, Al.
Hess, J.W.	Szakáll, S.
Hess, J.W.	Szentesy, C.
Hîrtopan, P.	Tămaş, C.
Iancu, O.G.	Thanasuthipitak, P.
Ionescu, C. .	Thanasuthipitak, T.
Kasper, H.U.	Udubaşa, Gh.
Kearns, J.	Vaskovic, N.
Kouzmina, E.	Vereş, D.Ş.
Laczkó, A.A.	Viczián, I.
Marincea, Şt.	Viehman, I.
Mariş, C.	Vlad, Ş.N.
Matovic, V.	White, W.B.
Mârza, I.	Zugrăvescu, D.
Minuţ, A.	
Moazzen, M.	
Molnár, F.	
Mosonyi, E.	
Murariu, T.	
Năstase, R.	