Monographic study of the Late Cretaceous representatives of the bolivinoidid benthic foraminifera

Marius Dan Georgescu

Department of Geoscience, University of Calgary, 2500 University Drive NW, Calgary, AB T2N 1N4, Canada

Received: January 2018; accepted February 2018

Available online 13 April 2018

DOI: <https://doi.org/10.5038/1937-8602.62.1.1299>

Abstract. A high resolution taxonomy and classification framework is developed for the representatives of the calcareous benthic foraminiferal family Bolivinoididae; this family consists of 24 species assigned to three genera. This new classification includes data pertaining of the test ultrastructure, ornamentation and porosity, which are obtained with the aid of careful and extensive study of the specimens under the scanning electron microscope. Two new taxa are described: *Bolivinoides antiquum* – n. sp. and *Swiecickina* – n. gen. A biostratigraphical framework consisting of eleven biozones based on bioevents given by species of the genera *Bolivinoides* and *Swiecickina* is developed for the middle Santonian-Maastrichtian stratigraphical interval; its resolution is more than double when compared to those of the previously developed biostratigraphical frameworks based on the representatives of this group.

Keywords: Late Cretaceous, foraminifera, Bolivinoides, classification, biostratigraphy.

INTRODUCTION

The Cretaceous representatives of the genus *Bolivinoides* Cushman, 1927a received attention from taxonomists and biostratigraphers in the last seventy years, almost more than any other genus of Cretaceous benthic foraminifera. A variety of factors contributed to the scientist's interest and amongst them it should be mentioned the fast pace of evolution, interesting and often enigmatic test morphology, and especially in the last half century usefulness in biostratigraphical studies. Studies on *Bolivinoides* were related to the development of the first reliable chronological framework of the Cretaceous (Van Hinte, 1967) and contributed significantly to the benthic-planktic biostratigraphical framework correlations (Petters, 1976, 1977; Robaszynski et al., 1984). Other studies mark true milestones in the development and evolution of some schools of foraminiferal micropaleontology such as Edgell (1954) in Australia, Pożaryska (1954) in Poland, and Reiss (1954) in Israel.

The earliest species of *Bolivinoides* were formally described in the nineteenth century and were assigned to the genus *Bolivina* d'Orbigny, 1839: *B. draco* (Marsson, 1878) and *B. decorata* (Jones in Wright, 1886); they are today two of the most frequently cited species of this genus. The assignment to *Bolivina* was determined by the biserial chamber arrangement, and this perspective is apparent in all the reports prior to the description of the genus *Bolivinoides* by Cushman (1927a) during a period of considerable proliferation of genera. The methodology of study and description of new species based on gross test architecture feature continued nearly three decades after this event, until Reiss (1954) realized a major breakthrough by describing *B. pustulata* based especially

on the occurrence of pustulose ornamentation. In the same methodological stream is the study by Hofker (1957) who described *B. granulata*, one junior synonym of *B. pustulata* as shown by Barr (1966a). Another step forward was realized by Petters (1977) who based on well-preserved material from the New Jersey coastal plain reported one interesting ultrastructure feature, namely elongate pores in the species *B. culverensis*. Despite this evident trend of advancing the knowledge of the test ultrastructure and high-detail test morphology, most of the studies remained focused on the gross test architecture. This can be explained by a certain inclination towards practicality amongst those specialists that encountered the representatives of this genus.

The trend to include ultrastructure features and high resolution data on test morphology was boosted again by Georgescu et al. (2011) who described *Bolivinoides mirabilis* from the upper Campanian of the New Jersey coastal plain based primarily on the occurrence of pore mounds at the test surface; in addition, from the same stratigraphical level *Elongateporeia elongatoporosa*, which is another bolivinoidid genus, was described mainly by the occurrence of pore mounds at the test surface and slit-like pores. These proved the first valid species of *Bolivinoides* as acknowledged herein, after 45 years of almost continuous reports, and in which only species that are junior synonyms of the pre-existing ones were described. Despite these advances, test ultrastructures and high resolution morphological features were misunderstood in two subsequent studies (Beniamovsky and Kopaeovich, 2016; Dubicka and Peryt, 2016).

This work represents a complete review of the bolivinoidid taxa. Each species and genus was studied using the scanning electron microscopy (SEM) resulting in a reassessment of

*Correspondence: dgeorge@ucalgary.ca

the taxonomic validity of the respective taxa. By bringing in the public domain a variety of data on the test ultrastructures, including ornamentation and porosity features and reporting new and interesting morphological structures, it represents a continuation of the pioneering steps realized by Reiss (1954), Hofker (1957), Petters (1977), and Georgescu et al. (2011). Rather than representing a purely fundamental study, it proves its practicality by the development of a biostratigraphical framework with a number of biozones that exceeds more than two times that of the previous ones. Moreover, it provides a set of data can be further elaborated in paleobiogeographical and evolutionary studies, which will be presented shortly in a sequel study.

STUDIED MATERIAL

The fossil material used and presented in this article was collected over a time period of nearly three decades and practically had begun while learning from Prof. T. Neagu (University of Bucharest) the methodology of identifying benthic and planktic foraminifera. I can recall, and not without a certain nostalgia, the times when I had made the first steps in learning under his guidance to identify the species of *Bolivinoides* from the upper Santonian-middle Campanian chalk deposits of southeastern Romania: *B. strigillata* (Chapman, 1892), *B. culverensis* Barr, 1966a, and *B. decorata* (Jones in Wright, 1886).

The material used in this study comes from two major sources: Deep Sea Drilling Project/Ocean Drilling Program (DSDP/ODP) sites or holes drilled worldwide and collections from different museums and universities. Amongst other sources I mention, inland wells drilled for scientific purposes, industry wells drilled inland and offshore, inland sections and material sent to me by other foraminiferologists. Some of this material was published in two previous studies (Georgescu et al., 2011; Georgescu, 2012).

DSDP/ODP sites and holes

Tests of bolivinoidid benthics presented in this article were recorded in six DSDP/ODP sites and holes, which are given in stratigraphical order: DSDP Site 511 (middle Santonian-lower

Campanian), ODP Leg 174AX (upper Santonian-lower upper Campanian), ODP Hole 762C (upper Campanian-Maastrichtian), DSDP Site 95 (lower Maastrichtian), ODP Hole 761B (Maastrichtian), and DSDP Hole 111A (upper Maastrichtian) (Fig. 1). The general stratigraphy and bolivinoidid taxa encountered at each of these locations are presented below.

DSDP Site 511 was drilled in the Southern Atlantic Ocean (Falkland Plateau). *Bolivinoides antiquum* – n. sp. is recorded from middle Santonian-lower Campanian dark grey to greenish claystone sediments, with the lowest recorded occurrence in Core 40; the tests are in excellent state or preservation, often pristine. The stratigraphy at this site is primarily according to the framework by Huber (1992) for the Austral Realm.

ODP Leg 174AX at Bass River site yielded in general well-preserved bolivinoidid tests, which were reported previously by Georgescu et al. (2011). The general stratigraphy at this site is according to Georgescu (2006). Bolivinoidid taxa occur at two stratigraphical levels: upper Santonian and lower upper Campanian; both intervals are represented by sediments accumulated in high water energy conditions during transgressive intervals and for this reason the tests are frequently broken especially in the terminal portions where the test wall is thinner. Upper Santonian is represented by the Merchantville formation, which yielded abundant and often well-preserved specimens of *B. culverensis*, and only rare and poorly preserved specimens of *B. strigillata* were encountered in this lithostratigraphic unit. At a higher stratigraphical level, the Marshalltown Formation of the lower upper Campanian yielded two species: *B. elongatoporosa* (Georgescu et al., 2011) and *B. mirabilis* (Georgescu et al., 2011).

DSDP Site 95 was drilled in the Yucatan outer shelf, offshore Mexico. According to the original biostratigraphical framework by McNeely (1973) the Santonian is overlain by lower Campanian white chalks. Nederbragt (1991) considered Core 13 of latest Santonian age. Georgescu (2010a, b) assigned the top (section 1) of Core 13 to the lower Campanian. A careful examination of the top section of Core 13 realized in the course of this study shows the occurrence of two bolivinoidid species, namely *Bolivinoides delicatula* Cushman, 1927a and *Swiecickina senonicus* (Dain,

Fig. 1. Geographical locations of the DSDP/ODP sites and holes that were investigated in the course of the present study. Base map is after Hay et al. (1999).

1961), which indicate a higher stratigraphical level, lower Maastrichtian. Therefore, it is evident that at this site there is a gap that encompasses the Campanian and possibly the lowermost Maastrichtian stratigraphical interval.

ODP Hole 762C was drilled in the Exmouth Plateau in the East Indian Ocean as part of the ODP Leg 122. The biostratigraphical framework used herein is that of Georgescu (2014a). The middle Campanian-Maastrichtian sediments at this location consist of soft white and yellowish nannofossil chalks, and they yielded an excellent record of two species: *Bolivinoides delicatula* and *B. doreeni* Finlay, 1940; one other species, namely *B. gigantea* Hiltebertmann and Koch, 1950 occurs in one single sample (Fig. 2). The specimens collected are in general well preserved and rarely fragmented demonstrating that during most times there were quiet conditions at the basin floor, with low water energy.

ODP Hole 761B was drilled in the Wombat Plateau of the East Indian Ocean. The biostratigraphical framework used at this location is that given by Georgescu (2014b). *Bolivinoides delicatula* is the only bolivinoidid species encountered at this site, in the soft white chalks of late Campanian-Maastrichtian age.

DSDP Hole 111A was drilled in the Orphan Knoll of the North Atlantic Ocean. Bolivinoidid taxa are known from the upper Maastrichtian white and yellow chalks, where assemblages consisting of *Bolivinoides peterssoni* Brotzen, 1944, *B. draco* (Marsson, 1878), *B. doreeni*, *B. vistulae* Pożaryska, 1954 and *Swiecickina paleocenica* (Brotzen, 1948) are encountered.

Collections

Four collections that were examined in the course of this study provided specimens of bolivinoidid taxa: Ehrenberg Collection (Museum of Natural History, Berlin), Cushman Collection (National Museum of Natural History, Smithsonian Institution, Washington, D.C.), Petters Collection (Rutgers, The State University of New Jersey), and McGugan Collection (University of Calgary).

Ehrenberg Collection (Museum of Natural History, Berlin), which is also referred to as C.G. Ehrenberg Collection, was studied in detail especially for planktic foraminifera of Santonian-Maastrichtian age (Georgescu, 2012, 2103). For this study were used only the specimens collected from the bulk samples. Only a small number of bulk samples provided bolivinoidid taxa, as follows: *Bolivinoides laevigata* Marie, 1941 and *B. praelaevigata* Barr, 1966 (Sample 2217c, upper Campanian-Maastrichtian, Meudon, France), *B. australis* Edgell, 1954, *B. decorata* (Jones in Wright, 1886) and *B. peterssoni* (Sample 2404a, lower Maastrichtian, Rügen Island, Germany), *B. draco* (Marsson, 1878), *B. australis* and *B. gigantea* (Sample 2404b, lower Maastrichtian, Rügen Island, Germany), and *Swiecickina clavatus* (Plotnikova, 1967) (Sample 2344a, proximity of the Campanian/Maastrichtian boundary, Moen Island, Denmark).

Cushman Collection (National Museum of Natural History, Smithsonian Institution, Washington, D.C.) was primarily used to document the morphology of *Bolivina watersi* Cushman, 1927a. The results were published by Georgescu et al. (2011) and resulted in the emendation of the genus *Trachelinella* Montanaro Gallitelli, 1956.

Petters Collection (Rutgers, The State University of New Jersey) was examined in the course of the study of the planktic

ODP Leg 122, Hole 762C (core number-core section, sample depth)	Sample depth (meters below sea floor)	Stages	Biozonation	<i>Bolivinoides delicatula</i>	<i>B. doreeni</i>	<i>B. gigantea</i>
43-1, 67-68 cm	555.17	upper Maastrichtian	<i>N. hariaensis</i>			
43-2, 70-71 cm	556.70					
43-3, 72-73 cm	558.22					
43-4, 68-69 cm	559.68		<i>A. mayaroensis</i>			
43-5, 66-67 cm	561.16					
44-1, 70-71 cm	564.70			x		
44-2, 72-73 cm	566.22					
44-3, 70-71 cm	567.70			x	x	
44-4, 70-71 cm	569.20				x	
44-5, 70-71 cm	570.70				x	x
44-6, 63.5-65 cm	572.14				x	
45-1, 70.5-71.5 cm	574.21					
45-2, 69.5-70.5 cm	575.70					
45-3, 70-71 cm	577.20					
45-4, 67-68 cm	578.67					
45-5, 68.5-69.5 cm	580.19					
46-1, 67.5-68.5 cm	583.61					
46-2, 65-66 cm	585.15					
46-3, 64.5-65.5 cm	586.65					
46-4, 73-74 cm	588.23	lower Maastrichtian	<i>G. cuvillieri</i>			
47-1, 70-71 cm	593.20					
47-2, 71-72 cm	594.71					
47-3, 69.5-70.5 cm	596.20			x	x	
47-4, 71-72 cm	597.71				x	
47-5, 70-71 cm	599.20					
47-6, 70-71 cm	600.70		<i>G. subpetaloidea</i>			
48-1, 70-71 cm	602.70				x	
48-2, 71-72 cm	604.21					
48-3, 71-72 cm	605.71					
48-4, 70-71 cm	607.20	upper Campanian	<i>G. rajagopalani</i>			
48-5, 70.5-71.5 cm	608.71					
48-6, 72-73 cm	610.22					
49-1, 70-71 cm	612.20					
49-2, 71-72 cm	613.71			x		
49-3, 71-72 cm	615.21			x		
49-4, 70.5-71.5 cm	616.71					
49-5, 64.5-65.5 cm	618.15					
50-1, 70-71 cm	621.70			x		
50-2, 72-73 cm	623.22			x		
50-3, 72-73 cm	624.72			x		
50-4, 69-70 cm	626.19			x		
50-5, 63-64 cm	627.63					
50-6, 74.5-75.5 cm	629.25					
51-1, 77.5-78.5 cm	631.28			x		
51-2, 68-69 cm	632.68					
51-3, 71.5-72.5 cm	634.22			x		
51-4, 69.5-70.5 cm	635.70			x		
51-5, 75-76 cm	637.25			x		
51-6, 65.5-66.5 cm	638.66			x		
52-1, 73.5-74.5 cm	640.74			x		
52-2, 70-71 cm	642.20					
52-3, 73.5-74.5 cm	643.74					
52-4, 72-73 cm	645.22					
52-5, 70-71 cm	646.70			x		
52-6, 73-74 cm	648.23					
53-1, 69.5-70.5 cm	650.20			x		
53-2, 71-72 cm	651.71					
53-3, 73-74 cm	653.23					
53-4, 60-61 cm	654.60					
53-5, 70-71 cm	656.20					
53-6, 62.5-63.5 cm	657.63	middle Campanian	<i>G. ventricosa</i>			
54-1, 70-71 cm	659.70					
54-2, 68-69 cm	661.29			x		
54-3, 69-70 cm	662.80			x		
54-4, 71-72 cm	664.32			x		
54-5, 65-66 cm	665.76					
55-1, 69-70 cm	669.19					
55-2, 70-71 cm	670.70					
55-3, 68-69 cm	672.18					
56-1, 70-71 cm	678.69					
56-2, 70-71 cm	680.20					
56-3, 67-68 cm	681.67			x		
56-4, 71-75 cm	683.24			x		
56-5, 71.5-72.5 cm	684.72					
56-6, 71-72.5 cm	686.21					
57-1, 73.5-74.5 cm	688.24					
57-2, 67-68 cm	689.67					

Fig. 2. Stratigraphical distribution of the *Bolivinoides* species encountered in the middle-Campanian-Maastrichtian sediments in ODP Hole 762B.

foraminiferal assemblages from the nearby ODP Leg 174AX at Bass River site. The samples from the well Dickinson 1 were examined and all the species recognized and reported by S.W. Petters in 1977 could be examined. This well of particular importance for the *Bolivinoidea* assemblage is significantly richer than that of the coeval sediments of ODP Leg 174AX due to the deeper waters.

McGugan Collection (University of Calgary) includes picked samples and carefully selected specimens from a variety of localities in Western Europe. Most of the samples are from the chalks of Northern Ireland and England. In addition, there are specimens from Germany, The Netherlands and Denmark. Many of the specimens are well-preserved and for this reason were selected to illustrate no less than eleven species in the present study, representing about one half of the total number of bolivinoidid species: *Bolivinoidea pustulata* Reiss, 1954, *B. praelaevigata*, *B. laevigata*, *B. decorata*, *B. miliaris* Hiltebrandt and Koch, 1950, *B. sidestrandensis* Barr, 1966a, *B. peterssoni*, *B. australis*, *B. gigantea*, *B. regularis* Reiss, 1954 and *B. doreeni*. In general, the specimens of this collection proved of paramount importance in assessing the test ultrastructure features and ornamentation.

TEST MORPHOLOGY AND TERMINOLOGY

Test morphology of *Bolivinoidea* and associated genera was never presented in detail in the previous studies. Due to the character of the present review work, it was considered necessary to make such a presentation to provide an accurate and consistent terminology throughout the numerous species of this genus. Test morphological features are grouped into four categories: (i) gross architecture, (ii) chamber backward extensions (CBE), (iii) periphery and peripheral structures and (iv) wall ultrastructure, ornamentation and porosity.

Gross architecture, orientation and symmetry

The representatives of *Bolivinoidea* and associated genera present the same pattern of chamber addition throughout their life cycle: chambers are added one at a time and alternately with respect to the test growth axis resulting in a biserial appearance (Fig. 3). The earliest chamber of the test is the proloculus, which present a spherical or subspherical shape, and all the subsequently added chambers are elongate, with the longest axis intersecting that of test growth at a narrow angle. Test portion with the proloculus is the posterior one, whereas chambers are added at the opposite side of the test, which is considered the anterior portion; the center of the proloculus and the midpoint of the suture between the last-formed and penultimate chambers define the test growth axis and antero-posterior direction. Test part with the proloculus is often referred to as the earlier or older portion of the test, whereas the opposite one the late or younger one.

Each chamber bears one aperture and the only apertural opening visible at the test exterior is that of the last-formed chamber. Chambers are mostly low and overlap the previous ones at a variable and in general low rate. Test earlier chambers are simple, but the last-formed ones present backward extensions, which give the so characteristic test appearance of the genus. For each of the chambers, the portion towards the direction in which new chambers are added during ontogeny is referred to as distal, whereas the opposite one is proximal.

Fig. 3. Test orientation, views and gross architecture features of a bolivinoidid test.

The test is laterally compressed, and each of the two wider faces is referred to lateral side. The two views from above periphery and perpendicular to the test growth axis are the edge views. Edge view that shows the aperture is also known as the apertural view, whereas the opposite one is the anti-apertural one; because the apertural view was the only one used by most of the researchers dealing with *Bolivinoidea* and tests with resembling morphology, it is frequently referred to in the literature of specialty simply as edge view.

Bolivinoidea was considered traditionally a bilaterally symmetrical genus, with the plane of symmetry passing through the proloculus, growth axis, and midlines of the last-formed chamber, aperture and penultimate chamber. New high-resolution data indicate that slightly asymmetrical tests occur, and this feature is mostly apparent in the differences in the sculpture of the two lateral side and a discrete curvature of the test growth axis in the earlier portion of the test as seen in edge view.

Chamber backward extensions

This feature was and still is considered one the most important in defining the genus *Bolivinoidea*; such structures are also used extensively in the taxonomy and classification at species level. The terminology under which they occur in literature varies significantly from author to author. For example, earlier studies mention the chamber backward extensions as "... prominent oblong tubercles which are arranged in oblique rows" (Jones in Wright, 1886, p. 330) and "oblique raised ridges, sometimes broken into tubercles" (Chapman, 1892, p. 515); such descriptions indicate that this feature was considered pertaining of the test ornamentation. In the original description of the genus, Cushman (1927a, p. 89) referred to this feature as "costae or raised lobes of the test"; this description is somehow broad for the term "costae" pertains of ornamentation whereas expression "raised lobes of

the test” of chamber shape. The duality in describing the CBE as both ornamentation and chamber shape features persisted in the next decades and is apparent in some most influential works (Hiltermann and Koch, 1950; Edgell, 1954; Reiss, 1954; Barr, 1966; etc). In the *Treatise on Invertebrate Paleontology*, Loeblich and Tappan (1964, p. C549-C552) regarded these features completely in the range of ornamentation. The transition from the cameral backward lobes to ornamentation features was more rarely documented and exemplified herein through the description of the genus *Bolivinoidea* in *Foraminiferal Genera and Their Classification*, which mentions “... basal margin of the chambers may have short and proximally directed projections over the preceding septa that merge into longitudinal costae...” (Loeblich and Tappan, 1987, p. 500). In this study, the CBE are considered features that pertain completely of the chamber shape. The transition from CBE to ornamentation structures is documented only in a small number of species. More rarely was used the expression of “ornamentation lobes”, which mixes and unifies one feature pertaining of chamber shape and the test ornamentation. Such terminology and practice are not accepted herein for the two features or sets of features have distinct evolutionary trends.

The morphology of the CBE in the posterior part of the last-formed chambers of the test is directly related to the formation of the inter-CBE space that separates every two nearby such CBE. Both CBE and inter-CBE spaces can present different width on the same chamber and therefore, should be used with care in taxonomy and classification. In the past, the number of CBE was considered paramount in in the taxonomy at species level by most authors, but there can be found significant differences in the number of CBE reported for the same species by different authors. One interesting idea recognized herein with potential in stabilizing the species descriptions in the future, occurs in the description of *B. decorata* by Barr (1966a, p. 232), where the lobes are separated into groups occurring towards the inner portions of the chambers and outer lobes towards the periphery. Further developing this idea, three types of CBE are recognized in the terminology adopted in this study: apertural, median and peripheral (Fig. 4). Apertural CBE is usually the thickest of these posteriorly-oriented structures; it is situated in the innermost position of the chamber, and the portion between the two apertural CBE of one chamber (one on each lateral side) bears the aperture. Peripheral CBE is the outermost one and mostly the thinnest amongst the CBE of one chamber; there is only one P on each chamber presenting backward extensions, and this contrasts with the all the other backward extensions that are symmetrically developed on each lateral side of the chambers of the adult stage. All the CBE between the apertural and peripheral ones, and are termed median and labelled consecutively from apertural to peripheral: median 1, median 2, median 3, etc. (Fig. 5). A distinct terminology for the inter-CBE spaces is not developed yet for the lack of practicality, but it can be developed in the future as its usefulness is demonstrated.

References to the shape of the CBE and inter-CBE spaces between them occur sporadically in different studies, but without receiving a considerable and consistent weight in taxonomy and classification. This setting is changes in this study, where two types of CBE, which proved useful in recognizing the evolution of the species of *Bolivinoidea*,

Fig. 4. Terminology of the chamber backward extensions of a chamber of one bolivinoidid test.

are recognized (Fig. 5). Type 1, which is also referred to as “wide”, includes the CBE with a broadly rounded posterior end; the CBE are separated by narrow inter-CBE spaces, which occasionally present a cul-de-sac outline. Type 2 is also termed “narrow” consists of CBE with rounded posterior end, which are separated by inter-CBE with different outline. Type 1 occurs especially in the species that are close to the bolivinoid ancestor, whereas Type 2 in those species positioned towards the distal portions of the bolivinoidid lineages. Therefore, Type 1 is primitive whereas Type 2 evolved. Notably, the transition from Type 1 to Type 2 is gradual. The occurrence of Type 1 CBE throughout one lineage demonstrates its primitive character, whereas Type 2 CBE occurs throughout the more evolved lineages. This classification proves useful for it is consistent with the directional character of the evolution process and directly reflected in evolutionary classification.

Periphery and peripheral structures

Test periphery can be evaluated both in lateral and edge views. Periphery outline lobation can be evaluated in lateral view, whereas in edge view there can be studied the periphery shape and peripheral structures.

Four types of peripheral outline as seen in lateral view are recognized herein: simple, lobate, subdentate and dentate (Fig. 6). The simple outline occurs in species in which the sutures are indistinct conferring the test outline an entire appearance. Lobate outline occurs in species in which the sutures are only slightly constricted and peripheral CBE is attached to the previous chamber in the longitudinal row on its whole length; this type is apparent only in the adult portion of the test. The subdentate type presents distinctly indented sutures, and occasionally with an incipient detachment of peripheral CBE from the periphery in the last-formed one or two chambers of the test. Dentate outline is encountered in the taxa in which the peripheral CBE is distinctly detached from the periphery; this type occurs in the adult portion of the test and in such cases the periphery can present a zig-zag outline.

Fig. 5. Types and terminology of the chamber backward extensions (CBE) of selected species of the genus *Bolivinoidea*. Abbreviations: A-apertural, M-median and P-peripheral.

Periphery shape as seen in edge view can vary from subrounded to subangular and in some taxa is acute. This feature reflects always the lateral compression of the test that occurs in the entire group's species. It is to be noted the fact that subrounded periphery occurs in the species closest or closer to the bolivinid ancestors, whereas the acute one in those of the distal parts of the bolivinoidid lineages; therefore, the subrounded periphery is regarded as primitive, acute one as evolved and subangular one intermediate between the other two. Notably, periphery shapes together with the test lateral

compression were studied in the past in a view taken from anterior part of the test along the growth axis (Hiltermann and Koch, 1950; Edgell, 1954; Reiss, 1954, Barr, 1966a, etc).

In most of the bolivinoidid species the test periphery is simple, without peripheral structures. Peripheral structures occur in a small number of species, and are the result of the development of one CBE, namely the peripheral one. In these taxa, the peripheral CBE extends over that of the previous chamber of the row. In some species the extensions of successive chambers result in the formation of a true keel, which can be visible in lateral and edge views. Notably, the keeled periphery in a *Bolivinoidea* species was for the first time mentioned by Edgell (1954, p. 73).

Wall ultrastructure, ornamentation and porosity

Test wall ultrastructure terminology was provided in the case of Cretaceous planktic foraminifera by Georgescu (2016), and the course of this study it was evident that it can be applied in the case of the Cretaceous representatives of the genus *Bolivinoidea*. Therefore, it is followed herein. In most of the species the test wall is simple, and only in a small number is incipiently reticulately-ridged. Notably, the incipiently reticulately-ridged wall ultrastructure was recognized only on the apertural faces of the last-formed chambers.

Chamber surface can be smooth or ornamented, and two types of ornamentation are recognized herein: primary and secondary. Primary ornamentation is that which formed through the changes in the wall ultrastructure, whereas the secondary one evolved later in the group's evolution and formed through the transformation of some CBE (median and peripheral) into ornamentation structures (Fig. 7). The beginnings of the study of these high resolution morphological structures can be traced back at the works of Reiss (1954) and Hofker (1957) who described the first taxa based on features that include but are not restricted to test ornamentation. Although the SEM was introduced in the current micropaleontological practice in the latest sixties, no other advance in implementing ornamentation features in the group's taxonomy and classification until the description of the first species with pore mounded chamber surface by Georgescu et al. (2011).

The smooth chamber surface is the simplest type, occurs in primitive species and in some lineages consistently occurs in all the species that form them. Primary ornamentation can consist of pustules, pore mounds, or a combination of the two; short longitudinal costae, which are parallel to the test growth axis or periphery, are recognized in only one species. Pustules are imperforate dome-like structures, with a circular or elongate outline as seen from above chamber surface; the longest axis of the elongated pustules is mostly parallel or subparallel to the test growth axis. Pore-mounded ornamentation is more complex and Georgescu et al. (2011) developed a classification and terminology of the pore mounds. These structures can be circular or elliptical according to the outline as seen from above the chamber surface and incipiently developed and well-developed according to their prominence (Fig. 8). Secondary ornamentation occurs in a number of species and begins with the alignment of the CBE of two or more successive chambers in one of the two series of

Fig. 6. Types of the test outline as seen in lateral view (above) and peripheral structures (below) of the bolivinoidid tests.

Fig. 7. Main types of primary and secondary ornamentation in bolivinoiid tests. Abbreviation: CBE-chamber backward extensions.

chambers of one test. Two longitudinal ridges can be further developed in the proximity of the central suture between the two rows of chambers (Fig. 7). The secondary ornamentation on the test lateral sides is developed from the median CBE. Other secondary ornamentation structures can occur at the test periphery though the modification of the peripheral CBE and result in the formation of keels.

Fig. 8. Types of pore mounds according to outline and degree of development.

Pores of the tests of the study group are small-sized. In an early classification of these structures according to their shape at the chamber surface, Georgescu et al. (2011) recognized three categories: circular, elliptical and slit-like (Fig. 9). Notably, slit-like pores occur only in the tests that evolved well-developed pore mounds with elliptical outline. Pores with irregular outline occur, but more data are necessary in order to confer the status of distinct category for the 'irregular' outline; the occurrence of such pores is mentioned in the species description wherever they were observed in the course of this study.

CLASSIFICATION

Family Bolivinoiidae was erected by Loeblich and Tappan (1984), and the only genus included within it was *Bolivinoides* (Loeblich and Tappan, 1987). In this study the validity of Bolivinoiidae is accepted and it includes three genera: *Bolivinoides* Cushman 1927a, *Trachelinella* Montanaro Gallitelli 1956 and *Swiecickina* - new genus. An emendation of Bolivinoiidae appears necessary especially to accommodate the new data

on the test ultrastructural morphology. All the type specimens are deposited in the Willi Karl Braun Micropaleontology Collection (WKB) of the University of Calgary.

Family Bolivinoiidae Loeblich and Tappan, 1984 - emended

Synonymy

Bolivinoiidae LOEBLICH AND TAPPAN 1984, p. 42.

Bolivinoiidae Loeblich and Tappan 1984. Loeblich and Tappan 1987, p. 500.

Diagnosis. Calcareous benthic foraminifera with the last-formed chambers with backward extensions

Description. Test is biserial throughout and consists of proloculus followed by chambers alternately added with respect to the test growth axis. Last-formed chambers present three or more backward projections, which are oblique to the test growth axis and often cover the suture with the previous chamber of the same row. Sutures are depressed or lined with narrow ridges that can be anastomosed or not towards the central zigzag suture. The test is laterally compressed, with a subrounded to angular and keeled periphery. Aperture is elliptical, mostly on the apertural face of the last-formed chamber or at its base, and presents a thin imperforate outer rim and a short internal toothplate; in some taxa the aperture is rimmed and produced on a short neck. Test wall is calcitic hyaline and finely perforate; chamber surface can be smooth or ornamented with pustules, pore mounds or ridges.

Genera included

Bolivinoides CUSHMAN 1927a, p. 89.

Trachelinella MONTANARO GALLITELLI 1956, p. 38.

Swiecickina n. gen.

Stratigraphical range: Upper Cretaceous (middle Santonian)-Paleogene (Paleocene).

Fig. 9. Types of pores according to outline and general aspect.

Genus *Bolivinoides* Cushman, 1927a – new emendation

Type species: *Bolovina draco* Marsson, 1878

Synonymy

Bolivinoides CUSHMAN 1927a, p. 89.

Bolivinoides Cushman. Cushman 1927b, p. 61.

Bolivinoides Cushman. Galloway 1933, p. 351.

Bolivinoides Cushman. Cushman 1950, p. 257.

Bolivinoides Cushman. Hiltermann and Koch 1950, p. 597.

Bolivinoides Cushman. Hofker 1952, p. 377. Emended.

Bolivinoides Cushman. Montanaro Gallitelli 1957, p. 145.

Bolivinoides Cushman. Pokorný 1958 (1963), p. 329.

Bolivinoides Cushman. Dain 1961, p. 30.

Bolivinoides Cushman. Beckmann and Koch 1964, p. 37.

Bolivinoides Cushman. Loeblich and Tappan 1964, p. 549.

Bolivinoides Cushman. Barr 1966a, p. 227.

Bolivinoides Cushman. Loeblich and Tappan 1987, p. 500.

Bolivinoides Cushman. Revets 1996, p. 9.

Bolivinoides Cushman. Georgescu et al. 2011, p. 114. New emendation.

Elongateporeia GEORGESCU, ARZ, MACAULEY, KUKULSKI, ARENILLAS and PÉREZ-RODRÍGUEZ 2011, p. 117 (see the 'Remarks' section below).

Diagnosis. Bolivinoidids with the aperture on or at the base of the apertural face of the last-formed chamber and test surface smooth, pustulose, pore-mounded or finely striate.

Description. Test is biserial throughout, consisting of proloculus followed by chambers added alternately with respect to the test growth axis. Earlier chambers are subglobular, and the last-formed ones elongate, with the longest axis oblique to the test growth axis forming anteriorly an acute angle. Chambers increase in size as added throughout the ontogeny resulting in a flaring aspect of the test as seen in lateral view. Chamber posterior side present a looped wall that result in the occurrence of backward extensions, which can present a regular or less regular aspect. There is one apertural CBE that bears the aperture, one peripheral CBE at the periphery and one to several median CBE between them; in general the apertural CBE is the largest of these structures. Vestigial portions of the CBE can occur over the lateral sides of the test early portion of some species. Sutures are straight to slightly curved and often obscured by the CBE of the chambers added subsequently during the ontogeny. Test is laterally compressed, resulting in a subrounded to acute

periphery. Periphery can be subrounded to acute. In most of the species the periphery is simple, without peripheral structures; one peripheral keel occurs in a small number of taxa. Aperture has an elongate outline and occurs at the base or on the apertural face of the last-formed chamber; it is bordered on much of its circumference by a narrow imperforate lip, and presents a short toothplate towards the chamber interior. Test wall is calcitic, hyaline and perforate. Chamber surface can be smooth or ornamented. Two types of ornamentation can occur. Primary ornamentation consists of pore mounds and pustules that can present a circular or elongate outline and different degree of development; secondary ornamentation results from the modification of the CBE; it consists of ridges that are oblique or parallel to the test growth axis, and developed from the anterior towards the posterior side of the test. Pores are simple or elevated at the top of a pore mound, and can have circular or elongate outline and more rarely slit-like aspect.

Remarks. The emendation proposed herein is to accommodate the significant advances in understanding the wall ultrastructure, ornamentation and porosity. Identification of the two types of ornamentation, one generated by the evolution of the wall and the other by the modification in the CBE is a major advance in understanding the morphology and evolutionary changes in amongst the species of the genus *Bolivinoides*, and warrants the necessity of such taxonomic act. *Elongateporeia* was described by Georgescu et al. (2011) to accommodate a bolivinoidid lineage of late Santonian-Campanian age. Having consistently chamber backward extensions, the tests included in this lineage are all bolivinoidid, and for this reason are herein included within the synonymy list of *Bolivinoides*. *Elongateporeia* is a valid unit in evolutionary classification, where it has the rank of directional lineage.

Species included (in stratigraphical order).

Bolivinoides antiquum Georgescu, 2018 – n. sp.

B. strigillata (Chapman, 1892)

B. culverensis Barr, 1966a

B. angulata Reiss, 1954

B. pustulata Reiss, 1954

B. praelaevigata Barr, 1966a

B. laevigata Marie, 1941

B. decorata (Jones in Wright, 1886)

B. delicatula Cushman, 1927a

B. mirabilis Georgescu et al., 2011

B. elongatoporosa (Georgescu et al., 2011)

B. miliaris Hiltermann and Koch, 1950

B. sidestrandensis Barr, 1966a

B. peterssoni Brotzen, 1944

B. australis Edgell, 1954

B. giganteus Hiltermann and Koch, 1950

B. draco (Marsson, 1878)

B. regularis Reiss, 1954

B. doreeni Finlay, 1940

B. vistulae Pożaryska, 1954

Stratigraphical range: Upper Cretaceous (middle Santonian)-Paleogene (Paleocene).

Bolivinoides antiquum Georgescu – n. sp.
(Figs. 10 and 11)

Holotype. Specimen WKB 010205.

Holotype dimensions. Length: L=0.199 mm; width: W=0.111

Fig. 10. Three specimens of *Bolivinoides antiquum* – n. sp. from sediments of the South Atlantic Ocean (DSDP Site 511). 1: holotype, Sample 79-511-28-6, 20-24 cm (lower Campanian). 2: paratype, Sample 79-511-34-2, 14-19 cm (lower Campanian). 3: paratype, Sample 79-511-32-1, 19-22 cm (lower Campanian).

Fig. 11. Paratype of *Bolivinoides antiquum* – n. sp. from sediments of the South Atlantic Ocean (DSDP Site 511) and magnifications of the chamber surface showing the primary ornamentation consisting of pore mounds (1b, 1c).

mm; W/L=0.558; thickness: T=0.077 mm; T/L=0.387.

Paratypes. Five specimens, WKB 010206-010210.

Paratype dimensions. Length: L=0.179-0.212 mm; width: W=0.098-0.117 mm; W/L=0.530-0.585; thickness: T=0.054-0.084 mm; T/L=0.272-0.396.

Type locality. Deep Sea Drilling Project Site 511, South Atlantic Ocean, Falkland Plateau; geographical coordinates: 51° 00.28'S and 46° 58.30'W.

Type level. Holotype was collected from the grey claystones of early Campanian age of Sample 71-511-28-6, 18-26 cm

from a depth below the sea floor of 231.18-231.22 m. The stratigraphy at type locality is after Huber (1992) and Huber et al. (1995).

Name origin. Species named with the neutral form of the Latin word *antiquus* (ancient).

Diagnosis. Small-sized species of *Bolivinoides* with one median CBE and occasionally ornamented with pore mounds.

Description. Test consists of the proloculus followed by chambers alternately added with respect to the test growth axis resulting in a biserial arrangement. Chamber increase in size as added and their longer axis is tilted forming an acute angle with the test growth axis along a posterior-to-anterior direction. Up to four pairs of chambers present distinct and short backward extensions that are not extended over the previous chamber of the row; for every chamber there are one apertural CBE, one peripheral CBE and two median CBE, one on each lateral side of the test, so in lateral view there are in total only three CBE. Peripheral CBE is the smallest of them, whereas the apertural CBE is slightly larger or of equal size with the median CBE; median CBE is relatively narrow and bordered in general by two broad inter-CBE spaces that can present equal width or not. Sutures are depressed and straight to gently curved anteriorly, often obscured by the CBE of the subsequently added chamber. Periphery is subrounded and simple, without peripheral structures. Aperture has the shape of a medium high arch and is situated at the base of the apertural face of the last-formed chamber; periapertural structure is asymmetrical and consists of a narrow imperforate rim that curves towards interior and borders the aperture on one side. Test wall is calcitic, hyaline, simple and finely perforate. Small-sized pore mounds with a measured diameter of 0.0010-0.0020 mm, occur in some specimens (Fig. 11), but specimens with completely smooth chamber surface are also known; specimens ornamented with pore mounds occur in equal proportion with those that present a completely smooth test surface. Pores are small, have a circular to elliptical outline and present a measured diameter or maximum dimension of 0.0002-0.0006 mm; they are widely spaced, at a distance of the nearby pores that often exceeds around twenty times their diameter or maximum dimension, a display that confers a simple and solid aspect to the chamber wall. Elongate pores do not present a preferential arrangement.

Remarks. *Bolivinoides antiquum* differs from all the other species of the genus by having only one median CBE on each side of the test, and ornamentation consisting of pore mounds that occur only in some specimens. Sliter (1977a, pl. 12, fig. 8) reported as *B. laevigatus* Marie specimens with similar gross test architecture with *B. antiquum* from the upper Campanian sediments of DSDP Hole 327A drilled in the Maurice Ewing Bank. Rare such specimens were also recovered from the Maastrichtian sediments at the type location of *B. antiquum* (Fig. 12). Examination of the ultrastructural features of these specimens shows that the ornamentation consists of low and thin anastomosing ridges that are more prominent over the earlier portion of the test and the wall is densely perforated; these features contrast with those observed in *B. antiquum*. More material is needed in order to clarify the taxonomic status and stratigraphical range of these specimens with obvious bolivinoidid aspect.

Stratigraphical range: Middle Santonian-lower Campanian.

Geographical distribution. South Atlantic Ocean; this species is known only from the type locality.

Fig. 12. Specimen almost identical to that illustrated as *Bolivinoidea laevigatus* by Sliter (1977a) collected from the upper Campanian sediments of South Atlantic Ocean (DSDP Site 511), Sample 79-511-23-1, 38-41 cm.

Bolivinoidea strigillata (Chapman, 1892)
(Fig. 13)

Synonymy

Bolivina strigillata CHAPMAN 1892, p. 515, pl. 15, fig. 10.
Bolivinoidea austiniana CUSHMAN 1937, p. 104, pl. 15, fig. 10.
Bolivinoidea strigillata (Chapman). Williams-Mitchell 1948, p. 106, pl. 9, fig. 3. Hiltermann and Koch 1950, p. 614, figs 2-4 (1-13, 64), 5 (10, 68). Hofker 1952, fig. 1. Edgell 1954, p. 70, pl. 13, fig. 8, pl. 14, fig. 8. Hofker 1955, fig. a. Hofker 1957a, fig. 323: a. Hofker 1958a, pl. 1, figs 1-2. Pokorný 1958/1963, fig. 329: center figure. Witwicka 1958, p. 196, pl. 9, fig. 8. Barr and Cordey 1964, p. 308, pl. 49, figs 1-3. Goel 1965, p. 79, pl. 7, fig. 6. Barr 1966a p. 228, pl. 34, figs 8-9, pl. 37, figs 7-9.
Bolivinoidea decorata cf. *delicatula* Cushman. Edgell 1954, p. 71, pl. 13, fig. 7, pl. 14, fig. 7.
Bolivinoidea austinianus Cushman. Frizzell 1954, p. 111, pl. 16, fig. 16. Hiltermann 1963, pl. 1, figs 11, 15.
Bolivinoidea strigillatus (Chapman). Bykova 1959, fig. 1: 1. Vassilenko 1961, p. 186, pl. 39, fig. 7. Hiltermann and Koch 1962, pl. 48, figs 10-11. Hiltermann 1963, pl. 1, figs 12-13, 16, 18-19, 22-24. Küpper 1963, p. 609, pl. 1, fig. 6. Barr 1970, p. 651, pl. 100, figs 6-7. Voloshina 1974, p. 21, pl. 2, fig. 9. Heller 1975, pl. 4, fig. 11. Govindan 1977, p. 469, pl. 1, figs 1-8, pl. 2, fig. 13, text-fig. 10: a-c. Koch 1977, p. 44, pl. 12, figs 7-8. Petters 1977, p. 1035, pl. 1, fig. 9. Sliter 1977b, pl. 5, figs 5-6. Rozumeiko 1978, p. 135, pl. 20, fig. 4. Kaptarenko-Chernousova et al. 1979, p. 147, pl. 55, fig. 8. Swiecicki 1980, p. 190, pl. 6, figs 7-8. Gawor-Biedowa 1984, pl. 1, fig. 1. Hart et al. 1989, p. 324, pl. 7.5, fig. 8. Schönfeld 1990, p. 80, pl. 3, figs 12-13. Neagu 1992, p. 75, pl. 8, figs 1-8. Machaniec and Zapałowicz-Bilan 2005, fig. 3: 1. Beniamovsky and Kopaevich 2016, fig. 2: 12-15. Dubicka and Peryt 2016, p. 79, figs 3: 1-3. Dubicka in Walaszczyk et al. 2016, text-fig. 5: 2. Dubicka et al. 2017, fig. 6: S.
Bolivinoidea strigillata strigillata (Chapman). Belford 1960, p. 61, pl. 15, figs 14-18.
 Transitional specimens *Bolivinoidea strigillatus*-*B. decoratus*. Hiltermann 1963, pl. 1, figs 2-3, 7-9.
 Transitional specimens *Bolivinoidea strigillatus*-*B. austinianus*. Hiltermann 1963, pl. 1, fig. 14.
Bolivinoidea cf. *austinianus* (Cushman). Hiltermann 1963, pl. 1, figs 17, 20.

Fig. 13. Four specimens of *Bolivinoidea strigillata* (Chapman, 1892) from the upper Santonian of Germany. 1-2: specimens previously illustrated by Schönfeld (1990, pl. 3, figs 12-13); © Schweizerbart Science Publishers; published with permission. 3-4: specimens previously illustrated by Koch (1977, pl. 12, figs 5, 8); © Schweizerbart Science Publishers; published with permission.

Bolivinoidea cyrenaicus BARR 1970, p. 646, pl. 99, figs 8-9, pl. 100, figs 4-5.

Bolivinoidea culverensis Barr. Neagu 1992, p. 75, pl. 7, fig. 1-11, pl. 8, figs 9-18. Dubicka and Peryt 2016, p. 79, fig. 3: 6-8. Dubicka in Walaszczyk et al. 2016, text-fig. 5: 3. Transitional forms between *Bolivinoidea strigillatus* and *B. culverensis*. Dubicka and Peryt 2016, fig. 3: 4-5.

Bolivinoidea cf. *culverensis* Barr. Beniamovsky and Kopaevich 2016, fig. 2: 16-18.

Diagnosis. Species of *Bolivinoidea* with two to three CBE on each lateral side of one chamber and with a deep longitudinal groove in the central portion of each lateral sides of the test.

Description. Test consists of the proloculus followed by chambers added alternately with respect to the test growth axis, resulting in a biserial arrangement. Chambers increase in size as added, are relatively high and with the longer axis obliquely oriented and forming an acute angle with the growth axis. The last-formed four or five pairs of chambers present well-developed narrow backward extensions that pass over the suture with the previous chamber of the row. Each chamber presents one apertural CBE, one peripheral CBE and four to six median CBE, which are symmetrically arranged on the two lateral sides; therefore, there are two to three

median CBE visible on each lateral side of one chamber. Median CBE are in general larger than the apertural CBE, and the peripheral CBE is the smallest of them. Sutures are depressed, straight or slightly curved in the direction of growth, and often covered by the CBE of the subsequently added chambers. Periphery is subrounded and simple, without peripheral structures. Aperture has the shape of an arch situated at the base of the apertural face of the last-formed chamber, and in some specimens occurs higher on the apertural face, with a pseudosuture at the base. The aperture is partly bordered by a narrow imperforate rim. Test wall is calcitic, hyaline, simple and perforate. Chamber surface is smooth, but the earlier portion of the test on the lateral sides can have an ornamented aspect due to the occurrence of prominent vestiges of CBE, which occasionally present a nodular appearance. A well-defined and deep groove can occur in the central portion of the lateral sides of some specimens as result of a loose arrangement of the CBE of successive chambers. Pores are small, with a circular or elliptical outline; there is no preferential arrangement of the elongate pores.

Remarks. *Bolivinoides strigillata* differs from *B. antiquum* mainly by having two to three rather than one median CBE on each lateral side of the last-formed chambers; it is also a much larger species (Hiltermann and Koch, 1950; Edgell, 1954; Barr, 1966a), with the average length that is more than double when compared to that of *B. antiquum*. The variability of the position of the aperture was well-documented and illustrated in the classical works of Hiltermann and Koch (1950), Edgell (1954) and Barr (1966a); such variability seemingly indicates the aperture shift from the base of the apertural face on a direction towards the apex of the test. As the aperture advances in a higher position on the apertural face, the bordering walls joint along a pseudosuture at the base of the apertural face; the term pseudosuture was proposed by Georgescu (2009) in the case of the genus *Praeplanctonia* of Early Cretaceous (late Albian) age. Wall porosity could not be observed directly in the course of this study because all the examined specimens presented the test surface affected by diagenesis; Beniamovsky and Kopaevich (2016, fig. 4: d) illustrated a high-detail micrograph of the test surface of a relatively well-preserved specimen that presents a mixture of circular and elongate pores. No data on pores dimensions are available.

Stratigraphical range: Upper Santonian-lower Campanian.

Geographical distribution. UK (England), Australia, Africa (Libya), Asia (India, Kazakhstan), EU (France, Germany, Poland, Romania, The Netherlands), Ukraine, and USA (Alabama, New Jersey, Texas).

Bolivinoides culverensis Barr, 1966a
(Figs. 14 and 15)

Synonymy

Bolivinoides hiltermanni BARR 1966a, p. 229, pl. 36, figs 7-8, pl. 37, figs 1-3.

Bolivinoides culverensis BARR 1967, p. 136 (new name).

Bolivinoides cyrenaicus BARR 1970, p. 646, pl. 99, figs 8-9, pl. 100, figs 4-5.

Bolivinoides sirticus BARR 1970, p. 651, pl. 100, figs 8-12.

Bolivinoides culverensis Barr. Petters 1977, p. 1031, pl. 1, figs

10-12. Swiecicki 1980, p. 170, pl. 6, figs 5-6. Hart et al. 1989, p. 322, pl. 7.4, figs 10-11.

Bolivinoides angulatus Reiss. El-Nady 2006, p. 674, pl. 1, figs 1-2.

Elongateporeia culverensis (Barr). Georgescu et al. 2011, p. 118, pl. 2, figs 3-13, pl. 3, figs 1-3.

Elongateporeia (Bolivinoides) culverensis (Barr). Beniamovsky and Kopaevich 2016, fig. 4: d, f.

Diagnosis. Species of *Bolivinoides* with wide CBE and primary ornamentation consisting of a mixture of circular and ellipsoidal pore mounds.

Description. Test consists of the proloculus followed by chambers alternately added with respect to the test growth axis resulting in a biserial arrangement. Earlier chambers are subglobular, the elongate; chamber longer axis is tilted and forms with the test growth axis an acute angle along a

Fig. 14. Three specimens of *Bolivinoides culverensis* Barr, 1966a from the upper Santonian of the New Jersey Coastal Plain (ODP Leg 174AX). 1: specimen from Sample 174AX, 505.05-505.08 m previously illustrated by Georgescu et al (2011, pl. 2, figs 5-7). 2: specimen from Sample 174AX, 505.05-505.08 m. 3: specimen from Sample 174AX, 505.05-505.08 m.

Fig. 15. Wall ultrastructure, primary ornamentation and porosity of three specimens of *Bolivinooides culverensis* Barr, 1966a from the upper Santonian of the New Jersey Coastal Plain (ODP Leg 174AX). All the three specimens were collected from Sample 174AX, 505.05–505.08 m.

posterior-anterior direction. Chamber increase in size as added, conferring thus the test a flaring aspect in the direction of growth. The last-formed three to four pairs of chambers present well-developed, wide backward extensions, which extend over the previously added chamber of the same row. Each chamber has one apertural CBE, one peripheral CBE and four to six median CBE; the median CBE are symmetrically developed on the two lateral sides of the test and therefore, two to three occur on each lateral side. Apertural CBE is the largest of them and the peripheral CBE the smallest one; the median CBE gradually decrease in width from the aperture towards the periphery. All the CBE are separated by narrow inter-CBE spaces, and occasionally two such nearby CBE can be adjacent. Sutures are depressed, straight or slightly curved in the direction of growth and often covered by the backward extensions of the subsequently added chamber of the row. Test is strongly compressed in edge view. Periphery is subrounded to subangular and simple, without peripheral structures. Aperture has an elongate shape, with the longer axis corresponding with that of chamber elongation; it is situated on the apertural face of the last-formed chamber and in some specimens is adjacent to the suture between the last and penultimate of the test chambers. One imperforate peripheral rim, which is rarely preserved, borders the aperture on one side and extends towards the chamber interior as a short toothplate. Test wall is calcitic, hyaline, simple and perforate. Primary ornamentation consists of pore mounds, which occur especially over the chambers of the earlier portion of the test. Secondary ornamentation occurs over the earlier chambers and consists of vestigial CBE with a circular or elliptical outline. Pore mounds have an elliptical outline and are in incipient stage of development; they present a measured longer axis of 0.0032–0.0059 mm and shorter axis of 0.0016–0.0033 mm; in general they occur as scattered distinct structures, more rarely in clusters of two, three or more. Pores can present a circular diameter with a measured diameter of 0.0010–0.0019 mm or can be elongate, with a measured longer axis of 0.0015–0.0022 mm and shorter axis of 0.0003–0.0006 mm; strongly elongate pores have a slit-like aspect with the measured longer axis of 0.0018–0.0028 mm and shorter axis of 0.0007–0.0020 mm. Elongate and slit-like pores present a preferential arrangement, which is parallel to the test growth axis; pores with circular outline occur mostly over the earlier portion of the test.

Remarks. This species differs from *B. antiquum* and *B. strigillata* by having wider CBE, more median CBE (two to

three occur on each lateral side rather than only one), elongate pores presenting a preferential arrangement with the longer axis parallel to the growth direction, and primary ornamentation consisting of pore mounds that can be in incipient stage of formation or well-developed. Petters (1977, p. 1031, pl. 1, fig. 11) reported for the first time elongate pores in *B. culverensis* from the New Jersey Coastal Plain, which were termed “slit-like pore” and considered a characteristic of this species; their occurrence was later confirmed by Georgescu et al. (2011) based on well-preserved specimens from the same region. In the present study it is shown that elongate pores can occur in other species of *Bolivinooides*, but *B. culverensis* is one of the few in which they present a preferential arrangement parallel with the test growth axis. A description of the test ultrastructure (ornamentation and pore characteristics and distribution patterns), which is followed herein, was given by Georgescu et al. (2011). The evolutionary occurrence of this species below the Santonian/Campanian boundary was reported in several studies and confirmed herein (Barr, 1966a; Van Hinte, 1967; Georgescu et al., 2011) and confirmed herein; cases in which the lowermost occurrence is situated above the Santonian/Campanian boundary are considered local occurrences (e.g., Hampton et al., 2007; Bailey et al., 2009; etc). The two species described by Barr (1970) from the upper Santonian-lower Campanian rocks of Libya, namely *B. sirticus* and *B. cyrenaicus* are regarded as end-members in the spectrum of variability of *B. culverensis*, the former having a slower increase in chamber size through the ontogenetic development.

Stratigraphical range: Upper Santonian-lower Campanian.

Geographical distribution. UK (England), Africa (Egypt, Libya), and USA (New Jersey).

Bolivinooides angulata Reiss, 1954
(Fig. 16)

Synonymy

Bolivinooides angulata REISS 1954, p. 155, pl. 28, figs 1–4.

Bolivinooides angulatus Reiss. Hiltermann 1963, pl. 1, fig. 21.

Diagnosis. Species of *Bolivinooides* with variable number of CBE on the lateral sides of the test and well-developed central groove in the posterior portion of the lateral sides.

Description. Test consists of the proloculus followed by chambers alternately added with respect to the test growth axis resulting in a biserial arrangement. Chamber longer axis is tilted forming an acute angle with the test growth axis along a posterior-anterior direction. The last four to five pairs of chambers present narrow, straight to anastomosing backward extensions that can be extended over the previous chamber of the row or not. Each chamber bears one apertural CBE, which is the largest, one peripheral CBE that is the smallest one, and a variable number of median CBE; each chamber has two to six median CME, which are symmetrically arranged on the two lateral sides, so there are one to three visible on each lateral side. Sutures are depressed and obscured by the backward extensions of the subsequently added chamber of the row. Test is compressed in edge view, with the lateral sides gently concave in a view along the growth axis and from above the apertural face. Periphery is rounded and simple, without peripheral structures. Aperture has the shape of a medium high to high arch and is situated at the base of the last-formed

Fig. 16. Five specimens of *Bolivinoides angulata* Reiss, 1954 from the lower Campanian of Israel. 1: holotype previously illustrated by Reiss (1954, pl. 28, fig. 1); © Cushman Foundation; published with permission. 2-3: paratypes illustrated by Reiss (1954, pl. 28, figs 2 and 4 respectively); © Cushman Foundation; published with permission. 4-5: specimens illustrated by Hiltermann (1963, pl. figs 21a and 21b respectively); © Elsevier; published with permission.

chamber. The successive addition of CBE determines the formation of two longitudinal grooves, which are more visible along the tests growth axis on the lateral sides over the earlier portion of the test.

Remarks. No specimens of this rare species could be studied and therefore, the description is based entirely on the original figures of Reiss (1954) and those from the review of Hiltermann (1963). The two reports show a considerable variability in the test size: 0.47 mm in the case of the holotype (Reiss, 1954, pl. 1, fig. 1) and 0.28 mm for one of the specimens illustrated by Hiltermann (1963, pl. 1, fig. 21). In addition, in the original description and figuration by Reiss (1954) not only there can be two to three CBE on each side of the last-formed chamber, but these structures can also have an irregular and anastomosing aspect, whereas in the photographs given by Hiltermann (1963) there is only one median CBE on one lateral side of the last-formed chambers. No data on the test wall ultrastructural characteristics are known, but from the photographs of Hiltermann (1963) the test surface appears smooth. Notably, SEM micrographs on this species are not available yet and a restudy of

the type specimens and additional material from the original localities in the light of the modern study techniques would be most helpful. *Bolivinoides angulata* was considered a junior synonym of *B. culverensis* by Dubicka and Peryt (2016) but no arguments were brought by these authors to support such synonymy; this synonymy is not accepted herein due to the sharp differences in the test morphology of the two species.

Stratigraphical range: Lower Campanian, possibly the lower part of the Middle Campanian.

Geographical distribution. Middle East (Israel).

Bolivinoides pustulata Reiss, 1954
(Figs. 17 and 18)

Synonymy

(?) *Bolivinoides texana* CUSHMAN 1937, p. 104, pl. 15, fig. 12.

(?) *Bolivinoides texanus* Cushman. Frizzell 1954, 112, pl. 16, fig. 20. Hiltermann 1963, pl. 2, fig. 14, 16, 18.

Bolivinoides pustulata REISS 1954, p. 156, pl. 29, figs 9-10.

Bolivinoides granulata HOFKER 1957, p. 250, figs 303 (d-f), 307, 310.

Bolivinoides granulata primitiva Hofker 1957, p. 251, fig. 303: a-c.

Bolivinoides pustulata Reiss. Barr 1966a, p. 236, pl. 37, fig. 4.

Bolivinoides granulatus Hofker. Hiltermann and Koch 1962, pl. 51, figs 8-9. Hiltermann 1963, pl. 2, figs 5, 11-13.

Beckmann and Koch 1964, p. 39, pl. 5, figs 4-6. Hanzliková

1972, p. 82, pl. 19, fig. 13. Koch 1977, p. 52, pl. 13, figs 8-9.

Dailey 1983, pl. 2, fig. 10. Schönfeld 1990, p. 85, pl. 3, fig. 9.

Hofmann et al. 1999, pl. 1, fig. 12. Ohmert 2011, pl. 3, fig. 22.

Dubicka 2015, (?) fig. 9: E, G.

Bolivinoides pustulatus Reiss. Hiltermann 1963, pl. 2, fig. 17.

Swiecicki 1980, p. 185, pl. 6, fig. 9. Hart et al. 1989, p. 324,

Fig. 17. Four specimens of *Bolivinoides pustulata* Reiss, 1954 from the upper lower Campanian of Misburg, east of Hannover (Germany); all the specimens are from the McGugan Collection, University of Calgary.

Fig. 18. Micrographs of the chamber surface of two specimens of *Bolivinoides pustulata* Reiss, 1954 from the upper lower Campanian of Misburg, east of Hannover (Germany) showing the pustulose primary ornamentation; note in specimen 1 that pustules can fuse and form short costae in the peripheral region. Both specimens are from the McGugan Collection, University of Calgary.

pl. 7.5, fig. 6. Gawor-Biedowa 1992, p. 106, pl. 19, figs 7-8. Frenzel 2000, p. 144, pl. 24, fig. 10.

Bolivinoides granulatus emersus BECKMANN and KOCH 1964, p. 40, pl. 5, figs 1-3.

Bolivinoides praelaevigatus Barr. Petters 1977, p. 1034, pl. 1, figs 7-8.

Bolivinoides granulatus emersus Beckmann and Koch. Bolli et al 1994, p. 130, fig. 35: 4.

Diagnosis. Species of *Bolivinoides* with pustulose primary ornamentation, with the pustules fused occasionally forming short ridges.

Description. Test consists of the proloculus followed by chambers alternately added with respect to the test growth axis resulting in a biserial arrangement. Chamber longer axis is tilted, forming an acute angle with the test growth axis along a posterior-anterior direction. Chambers increase in size throughout the ontogenetic development resulting in a flaring aspect of the test; in some specimens the rate of last-formed pair of chambers size increase is considerably smaller than those between the earlier chambers. Up to five pairs of chambers present wider backward extensions, which frequently are closely spaced separated by narrow inter-CBE spaces, and more rarely appear adjacent to the nearby ones; the backward extensions are short and can be extended in some specimens over the previous chamber of the row. There is one apertural CBE, one peripheral CBE and four to six median CBE symmetrically distributed on the two lateral sides of one chamber; therefore, there are two to three median CBE on each lateral side of a chamber. Apertural CBE is the largest of these structures and the peripheral one the smallest. Sutures are depressed, straight to gently curved in the direction of growth, and mostly obscured by the backward extensions of the subsequent chamber in a row. Test is strongly compressed in edge view. Periphery is subangular; the peripheral CBE can present a continuous pattern resulting in the development of a peripheral rim or weak keel. Aperture has an elongate shape on a direction parallel with the chamber main elongation axis, is situated on the apertural face of the last-formed chamber and in rare specimens a pseudosuture can be observed between the aperture base and the suture between the last-formed two chambers of the test. Test wall is calcitic, hyaline, simple and finely perforate. Primary ornamentation consists of pustules that can have a circular or elliptical outline and present a measured diameter or maximum dimension of 0.0035-0.0139 mm; the primary ornamentation elements are denser and more prominent over

the earlier portion of the test. Pustules are frequently fused or partly fused towards the peripheral region resulting in the formation of longitudinal ridges with pustulose aspect. Secondary ornamentation occurs over the earlier portion of the lateral sides of the test, and consists of vestigial chamber backward extensions forming nodules and tubercles; in some specimens the larger pustules of the primary ornamentation can have the same sizes with the secondary ornamentation structures, which makes them more difficult to study. Most of the pores present a circular outline, with measured diameters of 0.0030-0.0055 mm. Pores with an elongate or irregular outline can occur, and in this case their measured maximum dimension is of 0.0043-0.0091 mm; the pores with elongate outline are randomly oriented and do not present a preferential arrangement.

Remarks. *Bolivinoides pustulata* differs from *B. culverensis* mainly by the primary ornamentation features, which is dominated by pustules and does not present pore mounds, and porosity features in which the elongate pores lack a preferential arrangement with the longer axis parallel to the test growth axis; in addition slit-like pores that occur consistently in *B. culverensis*, are absent in *B. pustulata*. The obvious resemblances between the gross test architecture of the two species together with the considerable morphological differences in the test ultrastructures indicate that SEM observations are necessary for their correct identification. The status of this species was differently considered by various authors. Hiltermann (1963) noted the similarities between *B. texanus*, *B. pustulata* and *B. granulata*; Barr (1966a) considered that *B. pustulata* and *B. granulata* are synonyms and the name of the former species has priority; subsequently, Schönfeld (1990) noted that such a synonymy is insufficiently documented and used *B. granulatus* as valid name for this taxon. In the actual study, *B. pustulata* and *B. granulata* are considered synonyms and the earliest described one is valid according to the Principle of Priority. Dubicka and Peryt (2016) included in the synonymy of *B. granulatus* the species described and named *B. pustulata* and *B. granulata* by Reiss (1954) and Hofker (1957) respectively; the priority in naming species follows the chronological order in which they were and are described across the biological and paleontological sciences and therefore, the anti-chronological order used by Dubicka and Peryt (op. cit.) cannot be accepted.

Stratigraphical range: Upper lower Campanian-lower upper Campanian.

Geographical distribution. USA (New Jersey, Texas), Caribbean Region (Trinidad), EU (Austria, Czech Republic, Germany, The Netherlands, Poland), Middle East (Israel), UK (England), and South Atlantic Ocean.

Bolivinoides praelaevigata Barr, 1966a
(Fig. 19)

Synonymy

Bolivinoides praelaevigata Barr 1966a, p. 237, pl. 38, figs 7-9.
Bolivinoides praelaevigatus Barr. Salaj and Maamouri 1982, pl. 4, fig. 9.

Bolivinoides praelaevigata Barr. Jaff et al. 2014, pl. 2, figs 11-12.

Bolivinoides laevigatus Marie. Dubicka and Peryt 2016, fig. 10: 1.

Fig. 19. Two specimens of *Bolivinoides praelaevigata* Barr, 1966a from the lower Maastrichtian of the surroundings of Norfolk (UK, England); both specimens are from the McGugan Collection, University of Calgary.

Bolivinoides granulatus Hofker. Dubicka in Walaszczyk et al. 2016, text-fig. 5: 9.

Diagnosis. Species of *Bolivinoides* with smooth chamber surface and weakly developed CBE that present a low relief.

Description. Test consists of the proloculus followed by chambers alternately added with respect to the growth axis resulting in a biserial arrangement. Chamber longer axis is tilted with respect to the test growth axis, forming an acute angle with it along a posterior-anterior direction; chamber s increase gradually in size as added. Up to six pairs of chambers of the adult stage present short and narrow to wide backward extensions with low relief, which are short and extend over the previously formed chamber of the row. Each of the last-formed chambers presents six to eight CBE as follows: one apertural CBE, which is often the largest one, one peripheral CBE, which can be very small and often the smallest of them, and four to six median CBE; median CBE are symmetrically arranged on the two lateral sides of each chamber, so on each lateral side there can be two to three CBE. In general there is a decrease in the median CBE size from the aperture towards the periphery, but specimens that present sharp differences from this pattern are encountered; in such specimens the term irregular can be applied to degree of development of the median CBE. In some specimens the CBE of the successively added chambers can form two weakly developed ridges parallel with the test growth axis over the lateral sides of the earlier portion of the test. Sutures are depressed and straight to gently curved in the direction of growth, and often obscured by the backward extensions of the subsequently added chambers. The test is strongly compressed in edge view, with parallel sides. Periphery is subrounded to subangular and simple, without periapertural

structures. Aperture is simple, with elongate outline and situated on the apertural face of the last-formed chamber; one weak pseudosuture can occur between the aperture base and the suture between the last-formed and penultimate chambers of the test as seen in edge view. One imperforate periapertural rim, which is rarely preserved, borders the aperture at least on one side and extends towards the chamber interior as a short neck. Test wall is calcitic hyaline, simple and perforate. Chamber surface is smooth, without primary or secondary ornamentation. Pores are small-sized, with a circular or more rarely elongate outline and a measured diameter or maximum dimension respectively of 0.0003-0.0009 mm.

Remarks. *Bolivinoides praelaevigata* presents in general large sizes, with a maximum measured length of 0.523 mm. The backward extensions are narrower when compared to those of *B. culverensis* and *B. pustulata*, but are separated by somehow wider inter-CBE spaces. The occurrence of one pseudosuture between the aperture base and the suture between the last-formed and penultimate chamber in some specimens is evident in the original figuration of this species (Barr, 1966a, pl. 38, figs 7b and 9b). Another interesting morphological feature of this species is the complete absence of the primary ornamentation; in addition, the secondary ornamentation is represented only by two weakly developed ridges parallel to the growth axis in the posterior side of the test on the two lateral sides. *Bolivinoides praelaevigata* differs from *B. strigillata* mainly by the weakly developed CBE, more compressed test with a slower rate of thickness increase resulting in parallel sides in edge view, and larger size. The evolutionary occurrence of this species was well-documented by Barr (1966a) and all the subsequent reports indicate higher stratigraphical levels within the species range.

Stratigraphical range: Uppermost lower Campanian-lower Maastrichtian.

Geographical distribution. UK (England), Africa (Tunisia), EU (France, Poland), and Middle East (Iraq).

Bolivinoides laevigata Marie, 1941
(Fig. 20)

Synonymy

Bolivinoides decorata (Jones) *laevigata* MARIE 1941, p. 189, fig. 281.

Bolivinoides decorata laevigata Marie. Hiltermann 1952, p. 63, text-figs 4-5. Hagn 1953, p. 75, pl. 6, fig. 23.

Bolivinoides praecursor REISS 1954, p. 156, pl. 30, figs 4-7.

Bolivinoides laevigata Marie. Hiltermann and Koch 1955, p. 367, pl. 27, figs 13-15. Hofker 1957, p. 252, fig. 304: c. Goel 1965, p. 83, pl. 7, fig. 19. Barr 1966a, p. 237, pl. 34, fig. 7, pl. 38, figs 1-5.

Bolivinoides praecursor Reiss. Bieda 1958, p. 41, fig. 12. Goel 1965, p. 79, pl. 7, fig. 8. Plotnikova 1967, p. 94, pl. 15, fig. 4. Kaptarenko-Chernousova et al. 1979, p. 146, pl. 55, fig. 10.

Bolivinoides laevigatus laevigata Marie. Vassilenko 1961, p. 195, pl. 40, figs 2-3.

Bolivinoides laevigatus finitima VASSILENKO 1961, p. 196, pl. 40, figs 5-6, 9.

Bolivinoides laevigatus Marie. Hiltermann and Koch 1962, pl. 51, fig. 11. Hiltermann 1963, pl. 2, figs 6-10. Plotnikova 1967, p. 93, pl. 14, fig. 7. Sliter 1968, p. 89, pl. 13, fig. 2.

Fig. 20. Four specimens of *Bolivinoidea laevigata* Marie, 1941 from the lower upper Campanian of Höver, east of Hannover (Germany); all the specimens are from the McGugan Collection, University of Calgary.

Voloshina 1974, p. 21, pl. 2, fig. 11. Koch 1977, p. 52, pl. 13, fig. 7. Petters 1977, p. 1033, pl. 1, fig. 4. Kaptarenko-Chernousova et al. 1979, p. 146, pl. 55, fig. 19. Hart et al. 1989, p. 324, pl. 7.5, figs 1-2. Schönfeld 1990, p. 81, pl. 3, figs 5-6. Rescher 1991, pl. 17, fig. 1. Gawor-Biedowa 1992, p. 102, pl. 17, fig. 2. Ohmert 2011, pl. 3, fig. 23. Sheldon et al. 2012, fig. 3. P. Jaff et al. 2014, pl. 1, figs 13-14, pl. 2, fig. 8. Dubicka 2015, fig. 8: B, D. Dubicka and Peryt 2016, p. 86, fig. 9: 8-11. Dubicka in Walaszczyk et al. 2016, text-fig. 5: 11. Transitional specimens *Bolivinoidea laevigatus*-*B. granulatus*. Hiltermann 1963, pl. 2, fig. 4.

Bolivinoidea hiltermanni GOEL 1965, p. 83, pl. 6, figs 17-18. *Bolivinoidea laevigatus laevigatus* Marie. Swiecicki 1980, p. 177, pl. 6, fig. 10.

Diagnosis. Species of *Bolivinoidea* with smooth chamber surface, in general with low relief CBE and secondary ornamentation weakly developed over the posterior portion of the test lateral sides.

Description. Test consists of the proloculus followed by chambers alternately added with respect with the test growth axis resulting in a biserial arrangement. Chamber longer axis is tilted, forming an acute angle with the test growth axis along a posterior-anterior direction; there is a gradual increase in chamber size resulting in a flaring test outline in lateral view. The last-formed five to seven pairs of chambers, mostly six, present chambers with wide to narrow backward extensions, which extend over the previous chambers of the row; the backward extensions present in general a low relief, and are separated by narrow inter-CBE spaces. Each chamber presents in general ten backward extensions as follows: one apertural CBE, which in many specimens is also the largest, one peripheral CBE that is the smallest and six median CBE symmetrically distributed on the two lateral sides of the

chamber; therefore, there are three median CBE on each lateral side of one of the last-formed chambers, and there are rare specimens where the number is reduced to two. In general the M1 CBE is the largest of them and there is a gradual decrease of the median CBE from the apertural face of one chamber towards the periphery; in some specimens the M1 CBE is larger than the apertural CBE. Sutures are straight to anteriorly curved and obscured by the backward extensions of the previously added chamber of the row. Test is laterally compressed, with a slow increase in chamber thickness throughout the ontogeny. Periphery is subrounded to subangular and simple, without peripheral structures. Aperture is elongate and situated on the apertural face of the last-formed chamber; aperture longer axis is in general parallel to the chamber longest axis; periapertural structure is rarely preserved and consists of one imperforate rim that borders the aperture on one side and extends towards the chamber interior as a short toothplate. In rare specimens, one pseudosuture can occur between the aperture base and the suture between the last-formed and penultimate chamber in edge view. Test wall is calcitic, hyaline, simple and perforate. Chamber surface is smooth, without primary ornamentation. Secondary ornamentation

can occur over the earlier portion of the test on the two lateral sides as vestigial CBE, but specimens showing this feature are rare; two longitudinal rows resulted from the loose alignment of the CBE can occur frequently on the lateral sides of the earlier portion of the tests. Pores are small, with a circular or more rarely elongate outline and a measured diameter or maximum dimension respectively of 0.0002-0.0009 mm.

Remarks. *Bolivinoidea laevigata* differs from *B. praelaevigata* by having more prominent CBE on the two sides of the test, a slow but constant chamber increase in thickness and therefore, lacking parallel test sides in edge view, and more regularly developed CBE especially in the last-formed chambers where the decrease in size from the aperture to periphery has a high degree of constancy; in addition, the number of median CBE is rather constant, three on each lateral side on one chamber, rather than variable, two to three as is in *B. praelaevigata*. Dubicka and Peryt (2016) considered *B. praelaevigata* a junior synonym of *B. laevigata*, without bringing any argument to support this synonymy; the high resolution data on the test morphology on the two species as presented herein, indicate that the two species are both distinct and valid; therefore, the perspective of Barr (1966a) is considered factual and paramount in the development of a high resolution classification. One possibility, which I consider distinct, is that Dubicka and Peryt (2016) misinterpreted the test morphology of *B. praelaevigata*, which was illustrated as drawings by Barr (1966sa, pl. 38, figs 7-9) for such technique can occasionally provide a distorted perspective on the features prominence.

Stratigraphical range: Uppermost lower Campanian-lower Maastrichtian.

Geographical distribution. EU (Denmark, France, Germany, Poland, The Netherlands), Asia (Russia), Middle East (Iraq,

Israel), UK (England), Ukraine, and USA (California, New Jersey, Texas).

Bolivinoides decorata (Jones in Wright, 1886)
(Fig. 21)

Synonymy

Bolivina decorata JONES in Wright 1886, p. 330, pl. 27, figs 7-8.

Bolivina decorata Jones. Heron-Allen and Earland 1910, p. 409, pl. 7, figs 1-2. Plummer 1931, p. 181, pl. 10, fig. 10. Macfadyen 1932, pl. 35, fig. 20. Applin and Applin 1944, pl. 5, fig. 9. Shimer and Shrock 1944, pl. 8, fig. 6. Moore et al. 1952, fig. 2-9: 6.

Bolivina lattacea CARSEY 1926, p. 27, pl. 4, fig. 9.

Bolivinoides decorata delicatula Cushman. Cushman 1931, p. 308, pl. 35, fig. 13. Cushman 1932, p. 338, pl. 51, fig. 6. Cushman 1933, pl. 26, fig. 19. Marie 1941, p. 188, pl. 29, fig. 280. Cushman 1944, p. 11, pl. 2, fig. 22. Cushman 1946, p. 113, pl. 48, figs 11-12. Cushman and Renz 1946, p. 36, pl. 6, fig. 4. Cushman 1948, p. 259, pl. 24, fig. 15.

Bolivinoides decorata (Jones). Sandidge 1932, p. 196, pl. 19, fig. 16. Dain 1934, p. 33, pl. 3, fig. 34. Cole 1938, pl. 4, fig. 9. Schijfsma 1946, p. 70, pl. 7, fig. 8. Cushman 1948, p. 259, pl. 24, fig. 14. Hofker 1955, fig. b. Hofker 1957a, fig. 323: b. Hofker 1958a, pl. 1, figs 3-10. Hofker 1958b, fig. 1. Yeremeyeva and Belousova 1961, p. 86, pl. 24, fig. 6. Barr 1966a, p. 231, pl. 34, figs 2-6, 12, pl. 35, figs 6-9, pl. 36, figs

1-5. Barr 1966b, p. 500, pl. 77, figs 4-5. Hofker 1966, pl. 3, fig. 50. Robaszynski et al. 1985, pl. 6, figs 1-2. Robaszynski and Christensen 1989, pl. 2, figs 1-3. Neagu 1992, p. 75, pl. 8, figs 19-24.

Bolivinoides decoratus (Jones). Keller 1939, pl. 2, fig. 3. Maslakova 1959, p. 115, pl. 14, figs 4-5. Akimetz 1961, p. 188, pl. 18, figs 17-18. Kaptarenko-Chernousova et al. 1963, p. 109, pl. 25, fig. 4. Baranovskaya and Bulinnikova 1964, p. 291, pl. 64, figs 1-8. Van Hinte 1967, pl. 1, figs 2-3. Hanzliková 1969, p. 31, pl. 5, fig. 10. Hanzliková 1972, p. 80, pl. 19, figs 7-9. Gorbenko 1974, p. 52, pl. 9, fig. 1. Petters 1977, p. 1032, pl. 1, fig. 2. Villain 1977, p. 69, pl. 5, fig. 14. Rozumeiko 1978, p. 136, pl. 20, fig. 6. Kaptarenko-Chernousova et al. 1979, p. 145, pl. 55, fig. 12. Neagu 1979, pl. 53, fig. 38. Swiecicki 1980, p. 172, pl. 6, figs 2-4. Salaj and Maamouri 1982, pl. 4, fig. 4. Gawor-Biedowa 1984, pl. 1, fig. 4. Olsson and Nyong 1984, pl. 4, fig. 10. Liszkowa and Morgiel 1985, pl. 2, fig. 1. Aliev et al. 1988, p. 106, pl. 11, fig. 9, pl. 12, fig. 1. Hart et al. 1989, p. 324, pl. 7.4, fig. 12. Bless et al. 1990, fig. 5: h, l-o. Gawor-Biedowa 1992, p. 99, pl. 20, fig. 4. Gasiński 1993, pl. 1, fig. 8. Hewaidy and El Ashwah 1993, pl. 4, fig. 14. Bolli et al. 1994, p. 129, fig. 34: 35. Ismail and El Saadany 1995, p. 195, fig. 11: 4. Speijer and van der Zwaan 1995, p. 46, pl. 9, fig. 2. Witte and Schuumann 1996, pl. 1, fig. 16. Widmark 1997, p. 34, fig. 13: G. Khalil and Mashaly 2004, pl. 6, fig. 4. El Ashwah 2005, pl. 1, fig. 10. Machaniec and Zapałowicz-Bilan 2005, fig. 3: 2. El-Nady 2006, p. 674, pl. 1, figs 5-7. Niazi et al. 2013, pl. 4, fig. 2. Jaff et al. 2014, pl. 1, fig. 6. Dubicka 2015, fig. 8: F, H. Dubicka and Peryt 2016, p. 80, fig. 4: 1-4. Dubicka in Walaszczyk et al. 2016, text-fig. 5: 5.

Bolivinoides draco (Marsson). Wicher 1949, pl. 5, fig. 5, pl. 7. Nomura 1995, pl. 2, fig. 22.

Bolivinoides decorata decorata (Jones). Hiltermann and Koch 1950, p. 606, figs 2-4 (14-25, 27-31, 35-38, 42-45), 5 (36, 71). Hagn 1953, p. 74, pl. 6, fig. 22. Pożaryska 1954, p. 154, fig. 5. Reiss 1954, p. 155, pl. 28, figs 5-8, 13. Hiltermann and Koch 1955, p. 365, pl. 28, fig. 4. Bieda 1958, p. 32, fig. 6. Pokorný 1958/1963, fig. 329: left figure. Witwicka 1958, p. 198, pl. 9, fig. 9. Ansary et al. 1962, pl. 4. Van Hinte 1963, p. 105, pl. 14, fig. 1. McGugan 1974, pl. 1, fig. 3.

Bolivinoides polonica POŻARYSKA 1954, p. 252, fig. 1.

Bolivinoides decoratus decoratus (Jones). McGugan 1957, p. 339, pl. 32, figs 10-15. Hiltermann and Koch 1962, p. 315, pl. 46, fig. 7. Hiltermann 1963, pl. 3, fig. 15, pl. 4, figs 6, 8-14. Beckmann and Koch 1964, p. 43, pl. 6, figs 11-15. Medizza 1967, p. 21, pl. 1, fig. 11. Heller 1975, pl. 4, fig. 12. Koch 1977, p. 51, pl. 12, figs 5-6. Jordan and Gasse 1986, p. 15, pl. 5, figs 1-3. Rescher 1991, pl. 17, fig. 2. Frenzel 2000, p. 136, pl. 25, figs 2-5. Georgescu 2012, pl. 1, figs 15-18.

Bolivinoides decoratus decorata (Jones). Bykova 1959, fig. 1: 2. Vassilenko 1961, p. 189, pl. 39, fig. 8-10. Plotnikova 1967, p. 89, pl. 14, figs 5-6.

Bolivinoides polonica Pożaryska. Hofker 1960c, figs 2-5. Pożaryska 1965, p. 100, pl. 14, fig. 2.

Transitional specimens *Bolivinoides draco miliaris*-*B. decoratus*. Hiltermann 1963, pl. 4, fig. 15.

Bolivinoides cf. *decorata decorata* (Jones). Van Hinte 1963, p. 105, pl. 14, fig. 2.

Fig. 21. Three specimens of *Bolivinoides decorata* (Jones in Wright, 1886) from the lower Maastrichtian of the surroundings of Norfolk (UK, England) (1-2) and upper Campanian of the surroundings of Lüneburg (Germany); all the three specimens are from the McGugan Collection, University of Calgary.

Bolivinoides rhombodecorata GOEL 1965, p. 80, pl. 7, figs 9-10.

Bolivinoides pustulodecorata GOEL 1965, p. 80, pl. 7, fig. 11.

Bolivinoides cf. *miliaris* Hiltermann and Koch. Van Hinte 1965, p. 81, pl. 1, fig. 4.

Bolivinoides cf. *decoratus* (Jones). Barr 1970, p. 647, pl. 100, figs 1-3.

Bolivinoides australis Edgell. Sliter 1977b, pl. 5, figs 2-3. Swiecicki 1980, p. 168, pl. 6, fig. 1.

Bolivinoides gr. *decoratus* (Jones). Tronchetti 2001, pl. 4, figs 1-2.

Bolivinoides gr. *miliaris* Hiltermann and Koch. Tronchetti 2001, pl. 4, figs 3-5.

Bolivinoides culverensis Barr transitional to *B. decoratus* (Jones). Dubicka 2015, fig. 6: F.

Bolivinoides decoratus (Jones) transitional to *B. sidestrandensis* Barr. Dubicka 2015, fig. 6: H.

Transitional between *Bolivinoides strigillatus* to *B. decoratus*. Dubicka and Peryt 2016, fig. 3: 9-12.

Bolivinoides culverensis Barr. Dubicka et al. 2017, fig. 6: P-R.

Diagnosis. Species of *Bolivinoides* with smooth surface and well-developed CBE.

Description. Test consists of the proloculus followed by chambers alternately added with respect to the growth axis resulting in a biserial chamber arrangement. Chambers are low and tilted, with the elongation axis obliquely oriented with respect to the test growth axis, with which it forms an acute angle. Chambers increase in size as added throughout ontogeny resulting in a flaring test; the last-formed four to six pairs of chambers present well-developed, narrow backward extensions that have a marked in relief aspect, and extend over the previous chamber of the row. The backward extensions are separated by inter-CBE spaces; in general the width of the inter-CBE spaces is almost equal to that of the adjacent backward extensions. Each chamber has ten to more rarely twelve CBE as follows: one apertural CBE, which can be the largest or not, one peripheral CBE that in general is the smallest one and six to more rarely eight median CBE; the median CBE are symmetrically distributed on the two lateral side of each chamber and therefore, three to more rarely four are visible on one lateral side. Median CBE present in general comparable sizes but rarer specimens, in which the M1 CBE is the largest and there is a decrease in backward extensions size from the aperture towards the periphery, are also known. Sutures are depressed and straight, often covered by the CBE of the successively added chamber of the row. Test wall is compressed in edge view, with a slight increase in thickness through ontogeny. Periphery is subangular, and the fused peripheral CBE of the last-formed chambers can confer the test a weakly keeled aspect in the adult portion. Aperture is elongate and situated on the apertural face of the last-formed chamber, with the elongation axis parallel or superimposed over the chamber elongation axis. Periapertural structure consists of one imperforate peripheral rim that borders the aperture on one side and extends towards the chamber interior as one short toothplate. Test wall is calcitic, hyaline, simple and perforate. The primary ornamentation is completely absent and therefore, the chamber surface appears smooth. Secondary ornamentation consists of vestigial structures with circular or elliptical outline of the CBE and occurs over the lateral sides of the earlier portion of the test. Most of the pores

have a circular outline, more rarely elliptical or irregular, and present a measured diameter or maximum dimension respectively of 0.0004-0.0011 mm; the pores with elliptical outline do not present a preferential orientation pattern.

Remarks. *Bolivinoides decorata* differs from *B. laevigata* mainly by having more prominent chamber backward extensions, which are narrower and separated by wider inter-CBE spaces. In addition the pores are slightly larger: 0.0004-0.0011 mm rather than 0.0002-0.0009 mm. This species differs from *B. strigillata* especially by having more regularly developed CBE over the last-formed chambers and less developed secondary ornamentation; the two species present numerous morphological resemblances, which determined some authors to illustrate specimens they considered transitional between *B. strigillata* and *B. decorata* (Hiltermann, 1963; Dubicka and Peryt, 2016). The observations in this study indicate that there is no overlapping between the stratigraphical ranges of the two species as previously shown for example by Barr (1966a) and Petters (1977); therefore, such alleged 'transitional' specimens are assigned either to *B. strigillata* or *B. decorata*.

Stratigraphical range: Middle Campanian-lower Maastrichtian.

Geographical distribution. UK (England, Northern Ireland), Africa (Egypt, Libya, Tunisia), Asia (Azerbaijan, Belarus, Kazakhstan, Iran, Russia), Caribbean Region (Trinidad), EU (Austria, Belgium, Czech Republic, France, Germany, Italy, Ireland, Poland, Romania, The Netherlands), Middle East (Iraq), Ukraine, USA (Alabama, Florida, Georgia, Maryland, New Jersey, Texas), East Indian Ocean, and South Atlantic Ocean.

Bolivinoides elongatoporosa (Georgescu, Arz, Macauley, Kukulski, Arenillas and Pérez-Rodríguez, 2011)
(Figs. 22 and 23)

Synonymy

Elongatoporeia elongatoporosa GEORGESCU, ARZ, MACAULEY, KUKULSKI, ARENILLAS and PÉREZ-RODRÍGUEZ 2011, p. 120, pl. 3, figs 4-14.

Bolivinoides granulatus Hofker. Dubicka and Peryt 2016, p. 86, fig. 9: 1-6.

Diagnosis. Species of *Bolivinoides* with the primary ornamentation consisting of ellipsoidal pore mounds, and main porosity represented by slit-like pores.

Description. Test consists of the proloculus followed by chambers alternately added with respect to the test growth axis resulting in a biserial arrangement. Chamber longer axis is tilted and forms an acute angle with the test growth axis along the posterior-anterior direction. There is a gradual increase of chamber size throughout ontogeny, resulting in a flaring aspect of the test in lateral view. The last-formed one or two pairs of chambers present wide and low relief backward extensions. Each of the last-formed chambers present four to ten such backward extensions as follows: one apertural CBE, one peripheral CBB that is the smallest of them and two to six median CBE symmetrically arranged on the two lateral sides of the test; therefore, on each lateral side there are one to two, rarely three CBE. In general the M1 CBE is larger than the apertural CBE. Based on the two reports of this species it appears reasonable to infer a reduction in the number of median CBE from two to three towards the base of its stratigraphical

Fig. 22. Four specimens of *Bolivinoides elongatoporosa* (Georgescu et al., 2011) from the upper Campanian sediments of the New Jersey Coastal Plain (ODP Leg 174AX). 1: holotype from Sample 174AX, 435.56-435.57 m previously illustrated by Georgescu et al. (2011, pl. 3, figs 4-5). 2: paratype from Sample 174AX, 435.56-435.57 m previously illustrated by Georgescu et al. (2011, pl. 3, figs 6-8). 3: paratype from Sample 174AX, 435.56-435.57 m previously illustrated by Georgescu et al. (2011, pl. 3, figs 10-11). 4: paratype from Sample 174AX, 435.56-435.57 m.

Fig. 23. Micrographs of the chamber surface of three specimens of *Bolivinoides elongatoporosa* (Georgescu et al., 2011) from the upper Campanian sediments of the New Jersey Coastal Plain (ODP Leg 174AX); all specimens are from Sample 174AX, 435.56-435.57 m. A detail of figure 1a was illustrated by Georgescu et al. (2011, pl. 3, fig. 9).

range to one towards the top. Sutures are depressed, straight or more rarely curved in the direction of growth and in general visible, without being obscured by the short CBE. Test is strongly compressed in edge view, with nearly parallel lateral

sides and a slight increase in thickness towards the last-formed chambers. Periphery is subangular and simple, without periapertural structures. Aperture is situated on the apertural face of the last-formed chamber, and has an elongate to more rarely subcircular outline; the aperture elongation axis matches that of the chamber elongation as seen in edge view. Periapertural structure is rarely preserved and consists of one imperforate rim that borders the aperture on one side and extends towards the chamber interior as a short toothplate. Test wall is calcitic, hyaline, simple and perforate. Primary ornamentation consists of scattered well-developed pore mounds with an elliptical outline with a measured length of 0.0053-0.0097 mm and a measured width of 0.0023-0.0034 mm; these structures are preferentially arranged with the longest axis parallel or subparallel to the test growth axis. In rare cases these structures can occur as clusters of two or three. Secondary ornamentation lacks completely in this species. Pores are slit-like, straight or undulated, with a measured length of 0.0034-0.0059 mm and a measured width of 0.0001-0.0003 mm.

Remarks. This species was first reported, described and formalized based on material from the Marshalltown Formation of the New Jersey coastal plain by Georgescu et al. (2011); the test ultrastructures are well-preserved and the ornamentation consisting entirely of well-developed elongate pore mounds was reported; the authors noted also that such ornamentation is not known in any other species of benthic foraminifera with biserial chamber arrangement (ibid.,

p. 122). The water energy during the time of accumulation of the Marshalltown Formation was relatively high and for this reason the last-formed chambers of the tests at the type locality are broken. Despite these difficulties the species was considered descendent from *B. culverensis* with which were included within the genus *Elongatoporeia* by Georgescu et al. (2011); notably, this genus that accommodates a directional lineage is a unit of evolutionary classification. It was recognized at that time a gap encompassing roughly the middle Campanian that separated the two species. But this gap was filled recently by Dubicka and Peryt (2016) who reported *B. elongatopora* as *B. granulatus* from Poland; the report indicates the occurrence of a zone in which the stratigraphical ranges of the ancestral *B. culverensis* overlap with that of the descendant *B. elongatopora* and therefore, validating the directional lineage *Elongatoporeia*. Dubicka and Peryt (2016, p. 86) misinterpreted the ornamentation in *B. elongatopora* and *B. granulatus* and drew the conclusion that such ornamentation "... can occur sporadically in an almost all bolivinoidids taxa." This interpretation is proved inaccurate in the present study where is demonstrated that the ornamentation of *B. elongatopora* consists of elongate pore mounds, whereas that of *B. pustulata* of domical pustules, which are imperforate. In fact the two species seemingly evolved from the same ancestor, which is *B. culverensis*, which explains somehow the high chances to be confused for each other in case the ultrastructural features are not taken in consideration.

Stratigraphical range: Middle-lower upper Campanian.

Geographical distribution. USA (New Jersey) and EU (Poland).

Bolivinoides delicatula Cushman, 1927a
(Figs. 24 and 25)

Synonymy

Bolivina decorata Jones. Cushman 1926b, p. 586, pl. 15, fig. 11. White 1929, p. 43, pl. 5, fig. 1.

Bolivinoides decorata delicatula CUSHMAN 1927a, p. 90, pl. 12, fig. 8.

Bolivinoides decorata delicatula Cushman. Cushman and Jarvis 1928, p. 99, pl. 14, fig. 9. Cushman and Jarvis 1932, p. 42, pl. 13, fig. 2. Cushman 1946, p. 113, pl. 48, figs 10, 13-14. Wicher 1949, p. 85, pl. 5, fig. 4, pl. 7. Hiltermann and Koch 1950, p. 612, fig. 5 (65-67). Hiltermann 1952, p. 61, text-figs 4-5. Liszka 1955, p. 184, pl. 12, fig. 14.

Bolivinoides sp. (813) "traubig". Wicher 1953, fig. 1: 4.

Bolivinoides decoratus latticeus (Carsey). Frizzell 1954, p. 112, pl. 16, figs 18-19.

Bolivinoides "traubig" n.sp. Bettenstaedt and Wicher 1955, pl. 2, fig. 21.

Bolivinoides delicatulus Cushman. Said and Kenawy 1956, p. 140, pl. 3, fig. 44. Bykova 1959, fig. 1: 4. Maslakova 1959, p. 115, pl. 14, fig. 6. Rauzer-Chernousova and Fursenko 1959, fig. 834. Akimetz 1961, p. 189, pl. 18, fig. 19. Vassilenko 1961, p. 193, pl. 40, fig. 7. Hillebrandt 1962, p. 71, pl. 5, fig. 13. Graham and Church 1963, p. 51, pl. 5, fig. 24. Plotnikova 1967, p. 90, pl. 15, fig. 3. Hanzliková 1972, p. 82, pl. 19, fig. 12. Gorbenko 1974, p. 53, pl. 9, fig. 2. Voloshina 1974, p. 21, pl. 2, fig. 12. Govindan 1977, p. 473, pl. 2, figs 11-12, 14, 16-17, text-fig. 10: d-g. Rozumeiko 1978, p. 139, pl. 20, fig.

Fig. 24. Three specimens of *Bolivinoides delicatula* Cushman, 1927a from the lower Maastrichtian sediments of Yucatan outer shelf (DSDP Site 95). All the specimens are from Sample 10-95-13-1, 101.5-103 cm.

8. Kaptarenko-Chernousova et al. 1979, p. 145, pl. 55, fig. 17. Dailey 1983, pl. 2, fig. 5. Tjalsma and Lohmann 1983, p. 5, pl. 4, fig. 7. Liszkowa and Morgiel 1985, pl. 2, fig. 2. Van Morkhoven et al. 1986, p. 337, pl. 110, figs 1-2. De Klasz et al. 1988, p. 57, pl. 12, fig. 9. Salaj 1988, pl. 1, fig. 16. Thomas 1990, pl. 1, fig. 15. Nomura 1991, pl. 1, fig. 6. De Klasz and de Klasz 1993, p. 314, pl. 1, fig. 10. Boltovskoy et al. 1995, p. 133, pl. 1, fig. 11. Widmark 1997, p. 34, fig. 13: E. Alegret and Thomas 2001, p. 279, pl. 2, fig. 11. El-Dawy 2001, pl. 2, fig. 4. Howe et al. 2003, pl. 8, figs 33-34. Campbell et al., 2004, fig. 14: H-I. El-Nady 2006, p. 675, pl. 1, figs 8-10. Naidin and Beniamovski 2006, pl. 3, fig. 19. Alegret et al. 2009, pl. 1, fig. 6. Anan 2011b, p. 136, pl. 1, fig. 3. Holbourn et al. 2013, p. 72, figs 1-3. Olshanetskiy 2015, p. 696, pl. 2, fig. 6.

Bolivinoides delicatula Cushman. Hofker 1957, p. 251, figs 303 (g), 304 (a-b), 306, 308-309, 311. Bieda 1958, p. 38, fig. 10. Kavary and Frizzell 1963, p. 29, pl. 4, figs 1-2. Sztrákos 2005, pl. 6, fig. 1.

Bolivinoides decoratus delicatulus Cushman. Hiltermann and Koch 1962, p. 315, pl. 47, fig. 5.

Bolivinoides cf. *delicatulus* Cushman. Hiltermann 1963, pl. 2, fig. 15.

Bolivinoides delicatulus delicatulus Cushman. Hiltermann 1963, pl. 1, figs 1, 4-6, 10, pl. 2, fig. 19, pl. 3, figs 1-2, 5-6, 10, 13-14, 16-18. Beckmann and Koch 1964, p. 41, pl. 5, figs 11-21. Medizza 1967, p. 22, pl. 2, fig. 7. Bolli et al. 1994, p. 129, fig. 34: 37.

Fig. 25. Detail micrographs of specimens of *Bolivinoides delicatula* Cushman, 1927a from the lower Maastrichtian sediments of Yucatan outer shelf (DSDP Site 95) showing the ornamentation consisting of fine costae consisting of aligned and fused pustules. All specimens are from Sample 10-95-13-1, 101.5-103 cm.

Transitional specimens *Bolivinoides delicatulus delicatulus*-*B. delicatulus regularis*. Hiltermann 1963, pl. 3, fig. 19.

Diagnosis. Species of *Bolivinoides* with primary ornamentation consisting of short fine costae parallel to the periphery or test longitudinal axis.

Description. Test consists of the proloculus followed by chambers alternately added with respect to the test growth axis resulting in a biserial arrangement. Chambers are low and tilted, with the elongation axis tilted and forming an acute angle with that of growth axis along a posterior-anterior direction. Chambers increase slowly in size as added, resulting in a gently anteriorly flaring outline in lateral view. Up to seven pairs of the last-formed chambers present wide backward extensions, which are short and attach to the previous chamber of the row, and in some cases are wider posteriorly and narrower at the mid-length; the backward extensions are separated by narrow inter-CBE spaces and occasionally are wider anteriorly. There are one anterior CBE, which is often the largest, one peripheral CBE that is the smallest, and four to eight median CBE, which are symmetrically distributed on the two sides of the test; therefore, on one lateral side of a chamber there are two to four median CBE. In general there is a marked decrease in the CBE on a direction for the aperture towards the test periphery. Sutures are straight or gently curved in the direction of test growth and mostly covered by the backward extensions of the subsequently added chamber of the row. Test is compressed in edge view, with a slight increase in chamber thickness throughout ontogeny, which results in slightly divergent lateral sides. Periphery is subangular, and with a weakly keeled appearance given by the continuous aspect of the peripheral CBE of the successively added chambers. Aperture is an elongate opening on the apertural side on the last-formed chamber. Periapertural structure is delicate and therefore, rarely preserved; there is one asymmetrical imperforate rim that borders the aperture on one side and extends towards the chamber interior as a short toothplate. Test wall is calcitic, hyaline, simple and perforate. Primary ornamentation is absent in most of the studied specimens; rare scattered domical or elongate pustules with a measured diameter or maximum dimension of 0.0035-0.0084 mm can occur especially over the weak peripheral keel formed by the peripheral CBE. Short costae that can be aligned either parallel to the periphery or to the longitudinal axis can occur over certain portions of the chambers. Secondary ornamentation consists of nodular structures with circular or elliptical

outline, which occur over the lateral sides of the earlier portion of the test; CBE confer the test a knitted aspect on the two lateral sides of the test. Pores are simple, mostly with a circular outline with a measured diameter of 0.0028-0.0051 mm; elongate pores that do not present a preferential arrangement pattern occur rarely.

Remarks. *Bolivinoides delicatula* is the only species of the genus that presents the primary ornamentation consisting of short longitudinal costae over certain portions of the chambers. Another characteristic feature is given by the general knitted aspect of the test in lateral view. The occurrence of one weak peripheral keel resulted from the

alignment of the peripheral CBE is herein reported for the first time; this structure confer a weakly rimmed aspect to the test in the peripheral region as seen in lateral view. Notably, in some specimens the weak keel may appear undulated in edge view. *Bolivinoides delicatula* is one of the few species of the genus that survived the mass extinction event at the Cretaceous/Paleogene boundary, but no continuous occurrences across this boundary were reported; seemingly it was drastically affected but recovered shortly afterwards. The extinction of this species is in the Paleocene; lower Eocene occurrence present corroded and abraded tests, which might indicate possible reworking; additional data are necessary to clarify if *B. delicatula* crossed the Paleocene/Eocene boundary.

Stratigraphical range: Middle Campanian-Paleocene.

Geographical distribution. Mexico, Africa (Egypt, Senegal), Asia (Belarus, India, Iran, Kazakhstan, Russia), Caribbean Region (Trinidad, Yucatan outer shelf), EU (Czech Republic, France, Germany, Italy, Poland, Spain, The Netherlands), Ukraine, USA (California, Texas), East Indian Ocean, North Atlantic Ocean, and South Atlantic Ocean.

Bolivinoides mirabilis Georgescu, Arz, Macauley, Kukulski, Arenillas and Pérez-Rodríguez, 2011 (Figs. 26 and 27)

Synonymy

Bolivinoides mirabilis GEORGESCU, ARZ, MACAULEY, KUKULSKI, ARENILLAS and PÉREZ-RODRÍGUEZ 2011, p. 115, pl. 1, figs 1-10. *Bolivinoides* ex. gr. *B. decorata* (Jones). Georgescu et al. 2011, p. 116, pl. 1, figs 11-12, pl. 2, figs 1-2.

Diagnosis. Species of *Bolivinoides* with well-developed CBE and the primary ornamentation consisting of pore mounds that can be circular or elongate.

Description. Test consists of proloculus followed by chambers alternately added with respect to the test growth axis resulting in a biserial arrangement. Chambers are low and with the elongation axis tilted, forming an acute with the test growth axis along a posterior-anterior direction. The chambers present a variable rate of size increase through ontogeny, which is higher in the earlier stage and lower in adult and gerontic individuals; chamber size increase results in a test flaring outline in the direction of growth as seen in lateral view. The last-formed four to six pairs of chambers present well-developed narrow backward extensions, which are separated by wide inter-CBE spaces; in general the inter-CBE

spaces are wider than the chamber backward extensions. For each chamber there are: one apertural CBE, which can be the largest in some specimens, one peripheral CBE that is the smallest and three to four median CBE, which are symmetrically distributed on the two sides of each chamber; therefore, there are three to four CBE on each lateral side of one chamber, and there is a distinct tendency to increase the number from three to four during the ontogenetic development. The CBE decrease in size on a direction from the aperture to periphery, and the decrease rate can be constant or non-linear. Test is compressed in edge view, with a low rate of chamber increase in thickness. Periphery is subangular to subrounded and

Fig. 26. Three specimens of *Bolivinooides mirabilis* Georgescu, Arz, Macauley, Kukulski, Arenillas and Pérez-Rodríguez, 2011 from the lower upper Campanian sediments of New Jersey coastal plain (ODP Leg 174AX). 1: specimen designated as holotype from Sample 174AX, Sample 435.85-435.86 cm and previously illustrated by Georgescu et al. (2011, pl. 1, figs 1-3). 2: specimen designated as paratype from Sample 174AX, Sample 435.85-435.86 cm and previously illustrated by Georgescu et al. (2011, pl. 1, figs 6-8). 3: specimen from Sample 174AX, Sample 435.57-435.58 cm illustrated as *Bolivinooides* ex. gr. *B. decorata* (Jones) by Georgescu et al. (2011, pl. 1, figs 11-12).

Fig. 27. Detail micrographs of the chamber surface (1a, 2-3) and periapertural structure (1b) of three specimens of *Bolivinooides mirabilis* Georgescu, Arz, Macauley, Kukulski, Arenillas and Pérez-Rodríguez, 2011 from the lower upper Campanian sediments of New Jersey coastal plain (ODP Leg 174AX). 1a: specimen previously illustrated by Georgescu et al. (2011, pl. 1, fig. 5). 1b: specimen previously illustrated by Georgescu et al. (2011, pl. 1, fig. 4). 2: specimen previously illustrated by Georgescu et al. (2011, pl. 1, fig. 9). 3: specimen previously illustrated by Georgescu et al. (2011, pl. 2, fig. 2).

simple, without peripheral structures. Aperture has an elongate outline and is situated on the apertural face of the last-formed chamber; one pseudosuture between the apertural base and the suture between the last-formed and penultimate chambers of the test occurs in some specimens. Periapertural structure consists of one imperforate rim that borders the aperture on one side and extends towards the chamber interior as a short toothplate. Test wall is calcitic, hyaline, simple and perforate. Primary ornamentation consists of incipient to well-developed pore mounds and its characteristics vary considerably through ontogeny. In earlier stages the pore mounds have a circular outline with a measured diameter of 0.0039-0.0056 mm but in the adult stage they can merge and form larger structures with irregular outline and gentler slopes. There are no elements of secondary ornamentation recognized in this species. In the earlier stages of development the pores are circular, with a measured diameter of 0.0011-0.0017 mm and more rarely elliptical, with a measured maximum dimension of 0.0017-0.0019 mm; in the adult and gerontic tests most of the pores are elliptical, with a measured maximum dimension of 0.0019-0.0042 mm.

Remarks. The general test aspect and variability of *B. mirabilis* resemble those of *B. decorata*, including the dominance of two test growth stages, which are the juveniles on one hand and mature and gerontic on the other, with a small number of tests within the intermediate size range. Notably, Georgescu et al. (2011) described the two test varieties as *B. mirabilis* new species and *B. ex. gr. B. decorata*; this interpretation is corrected herein and the two are considered conspecific. *Bolivinooides mirabilis* differs from *B. culverensis* mainly by having narrower and more prominent CBE that are separated by wider inter-CBE spaces. It differs from *B. elongatoporosa* especially by having the primary ornamentation without

well-developed and elliptical pore mounds and pores that are not slit-like, but with a circular and/or elliptical outline. The pore mounded primary ornamentation in *B. mirabilis* differentiates this species from *B. decorata*, where the primary ornamentation is completely absent; in contrast, *B. mirabilis* lacks the secondary ornamentation, which occurs on the two lateral sides of the test earlier portion in *B. decorata*.

Stratigraphical range: Lower upper Campanian.

Geographical distribution. USA (New Jersey).

Bolivinoides miliaris Hiltermann and Koch, 1950
(Figs. 28 and 29)

Synonymy

Bolivinoides decorata (Jones). Cushman 1933, pl. 26, fig. 18. Cushman 1946, p. 113, pl. 48, figs 8-9.

Bolivinoides decorata (Jones) forma typica. Marie 1941, p. 188, pl. 29, fig. 279.

Bolivinoides trinitatis Cushman and Jarvis. Kikoïne 1948, p. 22, pl. 2, fig. 2.

Bolivinoides draco miliaris HILTERMANN and KOCH 1950, p. 604, figs 2-4 (26, 32-34, 39-41, 46-48) 5 (39).

Bolivinoides draco miliaris Hiltermann and Koch. Sigal 1952, pl. 17, fig. 2. Pozaryska 1954, p. 254, fig. 4. Reiss 1954, p. 155, pl. 28, figs 9-12, 14. Bieda 1958, p. 44, fig. 14. Witwicka 1958, p. 200, pl. 9, fig. 11. Ansary et al. 1962, pl. 4. Hiltermann and Koch 1962, p. 317, pl. 46, fig. 9. Hiltermann 1963, pl. 4, figs 21-23. Küpper 1963, p. 609, pl. 1, fig. 8. Van Hinte 1963, p. 106, pl. 13, figs 7-8. Beckmann and Koch 1964, p. 44, pl. 6, figs 21-23. Cati 1964, p. 239, pl. 38, fig. 18. Barr 1966a, p. 234, pl. 35, figs 4-5. Medizza 1967, p. 24, pl. 2, fig. 4. Scheibnerová 1968, p. 52, text-fig. 8. Sliter 1968, p. 89, pl. 13, fig. 1. Ansary and Tewfik 1969, pl. 2, fig. 2. McGugan 1974, pl. 1, fig. 4. Koch 1977, p. 56, pl. 12, fig. 4. Swiecicki 1980,

p. 165, pl. 5, figs 20-22. Dailey 1983, pl. 2, fig. 9. Busulini et al. 1984, fig. 5: j. Schönfeld 1990, p. 84, pl. 3, fig. 11. Bolli et al. 1994, p. 130, fig. 35: 2. Ayyad et al. 1997, fig. 8: k. Frenzel 2000, p. 140, pl. 25, figs 10-11. Ismail 2002, pl. 2, figs 1-2. El-Nady 2006, p. 678, pl. 1, fig. 15. Ohmert 2011, pl. 3, fig. 21.

Bolivinoides decoratus decoratus (Jones). Frizzell 1954, p. 111, pl. 16, fig. 17.

Bolivinoides miliaris Hiltermann and Koch. Hofker 1957a, fig 322: a-b. Vassilenko 1961, p. 200, pl. 40, fig. 4, pl. 41, fig. 1. Van Hinte 1967, pl. 1, fig. 4. Barr 1970, p. 650, pl. 99, fig. 1. Govindan 1977, p. 472, text-fig. 10: h, l-m. Petters 1977, p. 1034, pl. 1, fig. 3. Gawor-Biedowa 1984, pl. 1, fig. 2. Liszkowa and Morgiel 1985, pl. 2, fig. 3. Hart et al. 1989, p. 324, pl. 7.5, fig. 3. Gawor-Biedowa 1992, p. 103, pl. 20, fig. 6. Hewaidy and El Ashwah 1993, pl. 4, fig. 13. Campbell et al., 2004, fig. 14: J-K. Anan 2011b, p. 138, pl. 1, fig. 7.

Bolivinoides peterssoni Brotzen. Witwicka 1958, p. 199, pl. 9, fig. 11.

Bolivinoides cf. *rhomboidea* (Cushman). Beckmann 1960, p. 62, fig. 10.

Bolivinoides draco verrucosus BECKMANN and KOCH 1964, p. 46, pl. 16, figs 16-20.

Bolivinoides draco verrucosus Beckmann and Koch. Medizza 1967, p. 25, pl. 1, fig. 15, pl. 2, fig. 8. Bolli et al. 1994, p. 130, fig. 35: 3.

Intermediate form between *Bolivinoides miliaris* and *B. decoratus*. Van Hinte 1967, pl. 1, fig. 1.

Bolivinoides draco draco (Marsson). Keller 1988, pl. 1, figs 10-11.

Bolivinoides cf. *draco* (Marsson). Bless et al. 1990, fig. 5: c.

Bolivinoides praemiliaris Rexilius. Campbell et al., 2004, fig. 14: L-M (nomen nudum).

Bolivinoides intermedius DUBICKA and PERYT 2016, p. 82, figs 5: 5-9, 6: 1-4.

Transitional form between *Bolivinoides intermedius* and *B. draco*. Dubicka and Peryt 2016, fig. 6: 5.

Bolivinoides intermedius Dubicka and Peryt. Dubicka in Walaszczyk et al. 2016, text-fig. 5: 6.

Diagnosis. Species of *Bolivinoides* with a weak peripheral keel, smooth chamber surface and secondary ornamentation well-developed over the most surface of the lateral sides.

Description. Test consists of the proloculus followed by chambers alternately added with respect to the

Fig. 29. Detail micrographs of one specimen of *Bolivinoides miliaris* Hiltermann and Koch, 1950 from the upper upper Campanian of the surroundings of Lüneburg (Germany) showing the secondary ornamentation on the earlier portion of the lateral side (1a), porosity on the apertural face of the penultimate chamber (1b) and peripheral weak keel (1c); specimen from the McGugan Collection, University of Calgary.

Fig. 28. Two specimens of *Bolivinoides miliaris* Hiltermann and Koch, 1950 from the upper upper Campanian of the surroundings of Lüneburg (Germany); both specimens are from the McGugan Collection, University of Calgary.

growth axis forming a biserial arrangement. Chambers are low and their elongation axis forms an acute angle with the test growth axis along a posterior-anterior direction. Chambers increase in size throughout ontogeny, resulting in a test with flaring aspect in the direction of growth and distinct rhomboidal outline in edge view. The last-formed five or six pairs of chambers present narrow backward extensions separated by wide inter-CBE spaces; the backward extensions are relatively short, extend and attach to the previous chamber of the row, and occasionally are fused with similar structures of the previous chambers forming longer such structures with a continuous aspect. Each chamber presents several backward extensions as follows: one apertural CBE, which occasionally can be the largest, one peripheral CBE that is the smallest and six to eight median CBE, which are symmetrically distributed on the two chamber lateral sides; therefore, there are three to more frequently four CBE on one lateral side of a chamber. The median CBE present a well-marked decrease in size on a direction from the aperture towards the periphery, and frequently the M1 CBE is the largest of the backward extensions of one chamber. Sutures are straight or gently curved in the direction of growth and often obscured by the subsequently added chambers. Test is compressed in edge view, and chambers present a marked increase in thickness through ontogeny. Periphery is subangular to angular, with a weak keel formed by the peripheral CBE of the subsequently added chambers; the weak keel can have a continuous appearance in the portion of the earlier chambers of some specimens. Aperture has an elongate shape, with the longer axis corresponding to that of the respective chamber elongation; a prominent transversal wall inflexion can occur on the apertural face between the aperture base and suture between the last-formed and penultimate chambers of the test. Periapertural structure consists of one imperforate peripheral rim that borders the aperture on one side and extends towards the chamber interior as a short toothplate. Test wall is calcitic, hyaline and perforate. Primary ornamentation is completely absent in the representatives of this species. Secondary ornamentation consists of vestigial CBE with circular or elongate outline, which occur over 75% of the posterior part of the lateral sides of the test; the vestigial structures are aligned and form two ridges with discontinuous aspect, one on each side of the test growth axis on both lateral sides of the test. The weak peripheral keel is also a structure pertaining of the secondary ornamentation. Pores have a circular outline with a measured diameter of 0.0005-0.0016 mm; elongate pores that do not present a preferential alignment and/or pores with irregular outline are extremely rare. Test wall ultrastructure is simple or incipiently simple-ridged, one feature that can be observed especially on the apertural face of the last-formed chamber of well-preserved specimens.

Remarks. *Bolivinoidea miliaris* differs from *B. decorata* mainly by the angular periphery that bears a weak keel and secondary ornamentation, which is developed over most of the test lateral sides; in addition, the two ridges with discontinuous aspect developed on the two lateral sides symmetrically with respect to the test growth axis are more evident and prominent in the former species when compared to the latter. The chamber increase in thickness is at a higher rate in *B. miliaris* when compared to *B. decoratus*. *Bolivinoidea draco verrucosus* described by Beckmann and Koch (1964) and *B.*

intermedius of Dubicka and Peryt (2016) are based on small magnitude morphological differences and fit within the range of variability of *B. miliaris*; therefore, they are considered junior synonyms of this species.

Stratigraphical range: Upper upper Campanian-lower Maastrichtian.

Geographical distribution. EU (Austria, Belgium, Czech Republic, France, Germany, Italy, Slovakia, The Netherlands), Africa (Egypt, Libya, Tunisia), Asia (India, Kazakhstan), Caribbean Region (Trinidad), South America (Chile), UK (England, Northern Ireland), USA (California, New Jersey, Texas), and East Indian Ocean.

Bolivinoidea sidestrandensis Barr, 1966a
(Figs. 30 and 31)

Synonymy

Bolivinoidea curtus Reiss. Said and Kenawy 1956, p. 140, pl. 3, fig. 43.

Bolivinoidea sidestrandensis BARR 1966a, p. 239, pl. 34, figs 10-11, pl. 36, figs 6, 9, pl. 37, fig. 6.

Bolivinoidea sidestrandensis Barr. Sliter 1977b, pl. 5, fig. 4. Swiecicki 1980, p. 187, pl. 14, fig. 14. Hart et al. 1989, p. 324, pl. 7.5, fig. 7. Peryt 2000, pl. 1, fig. 4.

Fig. 30. Two specimens of *Bolivinoidea sidestrandensis* Barr, 1966a from the lower Maastrichtian of the surroundings of Norfolk (UK, England); both specimens are from the McGugan Collection, University of Calgary.

Fig. 31. Detail micrographs of two specimens of *Bolivinoides sidestrandensis* Barr, 1966a from the lower Maastrichtian of the surroundings of Norfolk (UK, England) showing the smooth chamber surface and CBE in the adult portion of the test (1a) and peripheral region (1b, 2); both specimens are from the McGugan Collection, University of Calgary.

diameter and/or maximum dimension of 0.0003–0.0008 mm.

Remarks. *Bolivinoides sidestrandensis* presents considerable morphological resemblances with *B. decorata* of which it differs mainly by having more CBE especially on the last-formed chambers of the test and more prominent secondary ornamentation that consists of two weak ridges consisting of loosely aligned CBE parallel to the test growth axis.

Stratigraphical range: Upper upper Campanian–lower Maastrichtian.

Geographical distribution. UK (England), Africa (Egypt), and EU (France, Poland).

Diagnosis. Species of *Bolivinoides* with well-developed narrow CBE separated by narrow inter-CBE spaces, and smooth chamber surface.

Description. Test consists of the proloculus followed by chambers alternately added with respect to the test growth axis resulting in a biserial arrangement. In lateral view the chambers are elongate, with the elongation axis tilted and forming an acute angle with the test growth axis along a posterior–anterior direction. Chambers increase in size as added through ontogeny, resulting in a test with flaring aspect as seen in lateral view; the last-formed four to five pairs of chambers present relatively narrow and short backward extensions, which are separated by narrower inter-CBE spaces and attach to the previous chamber of the row. There are several types of CBE, as follows: one apertural CBE, which is the largest one in most of the specimens, one peripheral CBE that is the smallest and six to ten median CBE, which are symmetrically arranged on the two lateral sides of the test; therefore, there are three to five median CBE on each lateral side of one chamber. There a distinct trend of increasing the number of median CBE in ontogeny. In all chambers, the median CBE decrease in size on a direction from the aperture to periphery, and in rare specimen the M1 CBE is slightly larger than the apertural CBE. Sutures are straight or slightly curved in the direction of growth, and are often obscured by the backward extensions of the subsequently added chambers in a row. Test is distinctly compressed in edge view, with nearly parallel lateral sides in the adult stage or with a slow increase in chamber thickness. Periphery is subrounded and simple, without peripheral structures. Aperture is situated on the apertural side and has an elongate outline, with the elongation axis superimposed over the chamber elongation axis as seen in edge view; the aperture does not come in contact with the suture between the last-formed two chambers of the test. Periapertural structure consists of one thin imperforate rim that is rarely preserved, which borders the aperture on one side and extends towards the chamber interior as a short toothplate. Test wall is calcitic, hyaline, simple and perforate. Chamber surface lacks primary ornamentation and therefore, has a smooth aspect. The secondary ornamentation consists of two ridges parallel to the test growth axis consisting of loosely aligned CBE and developed over the earlier two thirds of the lateral sides of the test. Pores have mostly a circular outline, more rarely elongate or irregular, and with a measured

Bolivinoides peterssoni Brotzen, 1944
(Fig. 32)

Synonymy.

Bolivinoides peterssoni BROTZEN 1944, p. 49, pl. 1, fig. 10.
Bolivinoides peterssoni Brotzen. Wicher 1953, fig. 1: 3.
 Pożaryska 1954, p. 256, fig. 8. Reiss 1954, p. 157, pl. 30, figs 12–14. Bettenstaedt and Wicher 1955, pl. 2, fig. 17.
 Hiltermann and Koch 1955, p. 366, pl. 28, figs 7–8. Hofker 1956a, p. 69, fig. 44. Hofker 1956b, fig. 27. Hofker 1957, p. 253, figs 305 (a–b, d), 312. Hofker 1960b, figs 8–9.
 Vassilenko 1961, p. 198, pl. 40, fig. 8. Hiltermann 1963, pl. 2, figs 1–3. Cati 1964, p. 240, pl. 38, fig. 19. Kiesel and Trümper 1965, p. 157, pl. 12, figs 5–6. Pringgoprawiro 1965, p. 35, pl. 7, fig. 1. Barr 1966a, p. 238, pl. 38, fig. 6. Plotnikova 1967, p. 95, pl. 15, fig. 5. Stenestad 1968, pl. 2, fig. 2. Barr 1970, p. 650, pl. 99, fig. 6. Al-Shaibani 1971, p. 109, pl. 3, fig. 1.
 Koch 1977, p. 59, pl. 13, fig. 6. Rozumeiko 1978, p. 137, pl. 20, fig. 5. Kaptarenko-Chernousova et al. 1979, p. 146, pl. 55, fig. 16. Swiecicki 1980, p. 183, pl. 6, fig. 11. Hart et al. 1989, p. 324, pl. 7.5, fig. 5. Schönfeld 1990, p. 82, pl. 3, figs 3–4. Gawor-Biedowa 1992, p. 105, pl. 19, fig. 9. Hewaidy and El Ashwah 1993, pl. 4, fig. 15. Frenzel 2000, p. 143, pl. 24, figs 11–15. El-Nady 2006, p. 679, pl. 1, figs 20–21. Anan 2011b, p. 138, pl. 1, fig. 10. Georgescu 2012, pl. 1, figs 19–20. Dubicka and Peryt 2016, p. 87, fig. 10: 2–5.
Bolivinoides compressa REISS 1954, p. 156, pl. 30, figs 1–3.
Bolivinoides compressa Reiss. Hofker 1956a, p. 70, fig. 45. Goel 1965, p. 82, pl. 7, fig. 16.
Bolivinoides draco miliaris Hiltermann and Koch. Said and Kenawy 1956, p. 140, pl. 3, fig. 40.
Bolivinoides mielnicensis BIEDA 1958, p. 35, fig. 8.
Bolivinoides delicatula Cushman. Goel 1965, p. 81, pl. 7, fig. 12.
Bolivinoides cf. *peterssoni* Brotzen. McGugan 1974, pl. 1, fig. 5.
Bolivinoides mielnicensis Bieda. Gawor-Biedowa 1992, p. 103, pl. 20, fig. 2.
Bolivinoides decoratus (Jones). Hofmann et al. 1999, pl. 1, fig. 16. Jaff et al. 2014, pl. 1, figs 3–5, 7–9, pl. 2, figs 1–5.
Bolivinoides paleocenicus Brotzen. Anan 2011b, p. 138, pl. 1, fig. 9. Beniamovsky et al. 2012, pl. 3, fig. 24.
Bolivinoides zikoi ANAN 2011b, p. 140, pl. 1, fig. 11.
Diagnosis. Species of *Bolivinoides* with smooth chamber surface and a relatively wide peripheral rim on the lateral side.

Fig. 32. Three specimens of *Bolivinoides peterssoni* Brotzen, 1944 from the lower lower Maastrichtian of the surroundings of Lüneburg (Germany) (1-2) and lower Maastrichtian of the Rügen Island (Germany) (3). 1-2: specimens from the McGugan Collection, University of Calgary. 3: specimen from the C.G. Ehrenberg Collection (Museum of Natural History, Berlin), Sample 2404a, which was previously illustrated by Georgescu (2012, pl. 1, fig. 19-20).

Description. Test consists of the proloculus followed by chambers alternately added with respect to the test growth axis resulting in a biserial arrangement. Chambers are low, with the elongation axis tilted and forming an acute angle with the test growth axis. Chambers increase in size as added through the ontogenetic development resulting in a flaring outline as the test is seen in lateral view. The last-formed four to six pairs of chambers present relatively wide backward extensions separated by narrow inter-CBE spaces; the backward extensions are short and attach to the previous chamber of the row. One chamber presents different types of backward extensions, as follows: one apertural CBE, which only in rare specimens is the largest, one peripheral CBE that is well-developed and attached to the previous such structures conferring the peripheral region a continuous appearance, and four to six median CBE, which are symmetrically arranged on the two lateral sides of the test; therefore, there are two to three median CBE on one lateral side of each chamber. There is a distinct tendency to increase the number of median CBE through the ontogeny; the M1 CBE is the largest in most of the specimens. Sutures are straight or gently curved in the direction of growth and mostly obscured by the backward extensions of the successively added chambers of the row. Test is compressed in edge view, with the chambers presenting a low rate of size increase in thickness and as a result, the two lateral sides may appear almost parallel or slightly divergent anteriorly. Periphery is subrounded to subangular and simple, without peripheral structures; the merged peripheral CBE form

a wide and smooth rim-like structure at the periphery, which is visible in the two lateral views, and is interrupted only by the proloculus that has a bulbous appearance. Aperture has an elongate shape and is situated on the apertural face of the last-formed chamber; its axis of elongations matches that of chamber elongation as seen in edge view. Periapertural structure consists of one imperforate peripheral rim that borders the aperture on one side and extends towards the chamber interior as a short toothplate. Test wall is calcitic, hyaline, simple and perforate. Chamber surface is smooth due to the absence of primary ornamentation. Secondary ornamentation consists of two closely-spaced or adjacent longitudinal ridges, one on each side of the test growth axis; the ridges are developed on the two lateral sides in the posterior one third to one half of the test length and are formed up by loosely aligned CBE that can fuse in some specimens. Pores are simple, mostly with a circular outline, extremely rarely elliptical or irregular in shape, and a measured diameter or maximum dimension respectively of 0.0003-0.0010 mm.

Remarks. *Bolivinoides peterssoni* differs from *B. laevigatus* mainly by having larger peripheral CBE that merge to form a rim-like structure, which in some specimens presents a considerable width, at the test peripheral region as seen in lateral views. Notably, the weak indentations that can be observed in lateral view at the periphery of some specimens are given by the junction between the peripheral CBE and previous chamber of the row. The wide rim-like structure visible in lateral views is the main feature that differentiates this species of *B. decorata*, in which is absent.

Stratigraphical range: Upper upper Campanian-Maastrichtian. This species became extinct at the Cretaceous/Paleogene boundary.

Geographical distribution. EU (Austria, Belgium, Denmark, France, Germany, Italy, Poland, Sweden, The Netherlands), Africa (Egypt, Libya), Asia (Kazakhstan, Russia), Middle East (Iraq, Israel), UK (England, Northern Ireland), and Ukraine.

Bolivinoides australis Edgell, 1954
(Figs. 33 and 34)

Synonymy

Bolivinoides decorata (Jones). Dampel 1934, p. 13, pl. 1, fig. 10.
Bolivinoides decorata delicatula Cushman. Kikoïne 1948, p. 21, pl. 1, fig. 10.

Bolivinoides decoratus (Jones). Hofker 1952, fig. 3. Sacal and Debourle 1957, p. 14, pl. 3, fig. 21. Rasheed and Ravindran 1980, p. 206, pl. 1, figs 1-4. Khalil 1998, fig. 8: 9.

Bolivinoides decorata australis EDGELL 1954, p. 71, pl. 13, figs 5-6, pl. 14, figs 5-6.

Bolivinoides australis Edgell. Hofker 1955, fig. d. Edgell. Hofker 1956b, figs 40-41. Hofker 1958a, pl. 1, figs 11-17, pl. 2, figs 18-23. Hofker 1958b, fig. 1. Hofker 1958c, pl. 6, figs 34-35. Hofker 1960a, fig. 8. Goel 1965, p. 82, pl. 7, figs 14-15. Barr 1966a, p. 238, pl. 34, fig. 1. Hofker 1966, pl. 10, figs 79-86, pl. 13, figs 37-42. Neagu 1979, pl. 53, figs 33-34. Robaszynski et al. 1985, pl. 6, figs 3-6. Robaszynski and Christensen 1989, pl. 2, figs 5-9. Bless et al. 1990, fig. 5: k. Felder and Bless 1993, fig. 6: S-W. Witte and Schuumann 1996, pl. 1, fig. 15. Frenzel 2000, p. 135, pl. 25, figs 7-9. Campbell et al., 2004, fig 14: F-G. Jaff et al. 2014, pl. 1, figs 1-2.

Fig. 33. Two specimens of *Bolivinoides australis* Edgell, 1954 from upper upper Campanian of the surroundings of Lüneburg (Germany); both specimens are from the McGugan Collection, University of Calgary.

Bolivinoides decorata australis Edgell. Bieda 1958, p. 33, fig. 7. McGugan 1974, pl. 1, fig. 2.

Bolivinoides cf. *doreeni* Finlay. Bieda 1958, p. 39, fig. 11.

Bolivinoides decoratus australis Edgell. Hiltermann 1963, pl. pl. 4, fig. 7. Mello 1969, p. 73, pl. 8, fig. 8. McNeil and Caldwell 1981, p. 229, pl. 18, fig. 7.

Bolivinoides decoratus delicatulus Cushman. Pringgoprawiro 1965, p. 34, pl. 7, fig. 2.

Bolivinoides giganteus Hiltermann and Koch. Stenestad 1968, pl. 2, fig. 1.

Bolivinoides miliaris Hiltermann and Koch. Salaj and Maamouri 1982, pl. 4, fig. 8. Dubicka and Peryt 2016, p. 82, fig. 5: 1-4.

Bolivinoides cf. *australis* Edgell. Tronchetti 2001, pl. 4, fig. 6.

Bolivinoides draco (Marsson). Wicher 1949, p. 85, pl. 5, fig. 7. Omaña 2006, fig. 4: 9.

Fig. 34. Detail micrographs of one specimen of *Bolivinoides australis* Edgell, 1954 from upper upper Campanian of the surroundings of Lüneburg (Germany) showing the CBE of the penultimate chamber (1a) and secondary ornamentation on one lateral side of the test (1b); specimen from the McGugan Collection, University of Calgary.

Bolivinoides giganteus Hiltermann and Koch. Dubicka and Peryt 2011, fig. 5: O. Dubicka and Peryt 2016, fig. 7: 5.

Diagnosis. Species of *Bolivinoides* with secondary ornamentation structures on the lateral sides forming a V-shaped pattern with the cusp opening in the direction of growth.

Description. Test consists of the proloculus followed by chambers alternately added with respect to the test growth axis forming a biserial arrangement. Chambers are elongate and low, with the elongation axis tilted and forming an acute angle with the test growth axis. Chambers increase in size as added throughout ontogeny, resulting in a flaring aspect of the test as seen in lateral view. The last-formed six to seven pairs of chambers present narrow backward extensions that attach to the previous chamber of the row and are separated by wide inter-CBE spaces. Each chamber presents different types of CBE, as follows: one apertural CBE, which is the largest amongst them, one peripheral CBE that is the smallest and ten to twelve median CBE that are symmetrically arranged on the two lateral sides of one chamber; therefore there are five to six median CBE on one lateral side of one chamber. The backward extensions of one chamber decrease in size on a direction from the aperture towards periphery. Sutures are straight or gently curved in the direction of growth, often obscured by the successively added chamber of the row and its backward extensions. Test is compressed in edge view, with the chambers presenting a slow but constant size increase resulting in slightly divergent lateral sides. Periphery is subangular and simple, without peripheral structures; in general it has a nodular and discontinuous appearance due to the peripheral CBE of the successively added chambers, which are not fused. Aperture is elongate and situated on the apertural face of the last-formed chamber; the aperture elongation axis matches the longest elongation axis of the chamber on which is situated. Periapertural structure consists of one delicate imperforate lip, which is rarely preserved; it borders the aperture on one side and extends towards the chamber interior as a short toothplate. Test wall is calcitic, hyaline, simple and perforate. Primary ornamentation is completely absent. Secondary ornamentation consists of interrupted CBE, which consists of structures with circular and elongate outline, is developed almost all over the test lateral sides except for the portion of the last two added chambers; the structures of secondary ornamentation are isolated, rarely fuse and form a more or less well-defined V-shaped pattern with the cusp opening in the growth direction. Pores are mostly circular and rarely present an elongate or irregular outline, and present a measured diameter or maximum dimension respectively of 0.0003-0.0009 mm.

Remarks. The secondary ornamentation consisting of circular and elongate structures that form a V-shaped pattern with the cusp opening in the direction of growth is a characteristic of this *B. australis*; this arrangement presents a considerable intra-specific variability.

Stratigraphical range: Uppermost upper Campanian-lower Maastrichtian.

Geographical distribution. Australia, Africa (Egypt, Tunisia), Asia (Kazakhstan, India), Canada (Manitoba), EU (Austria, Belgium,

Denmark, France, Germany, Poland, The Netherlands), Mexico, Middle East (Iraq), UK (England, Northern Ireland), USA (California, South Dakota), and East Indian Ocean.

Bolivinoides gigantea Hiltermann
and Koch, 1950
(Figs. 35 and 36)

Synonymy

Bolivinoides decorata (Jones). Cushman 1944, p. 10, pl. 2, fig. 23.

Bolivinoides decorata gigantea HILTERMANN and KOCH 1950, p. 610, figs 2-4 (49-51, 55-57, 61-63), 5 (50).

Bolivinoides decorata gigantea Hiltermann and Koch. Edgell 1954, p. 72, pl. 13, fig. 4, pl. 14, fig. 4. Pożaryska 1954, p. 255, fig. 6.

Bolivinoides gigantea Hiltermann and Koch. Hofker 1955, fig. c. Hofker 1956a, p. 69, fig. 49. Hofker 1956b, fig. 63. Hofker 1957, fig. 320: a-b. Hofker 1958a, pl. 2, figs 24-27. Hofker 1958b, fig. 1. Hofker 1960a, fig. 7. Hofker 1960b, figs 12-14. Hofker 1960c, fig. 1. Hofker 1962, text-fig. 9. Hofker 1966, pl. 18, fig. 13. Neagu 1979, pl. 53, figs 35-37.

Bolivinoides senonicus YEREMEYeva 1957, p. 12, pl. 1, fig. 8. *Bolivinoides giganteus* Hiltermann and Koch. Vassilenko 1961, p. 192, pl. 40, fig. 1. Barr 1970, p. 647, pl. 99, figs 4-5. Petters 1977, p. 1032, pl. 1, fig. 6. Gawor-Biedowa 1984, pl. 1, fig. 3. Bless et al. 1990, fig. 5: a-b, d. Gawor-Biedowa

Fig. 35. Two specimens of *Bolivinoides gigantea* Hiltermann and Koch, 1950 from uppermost Maastrichtian from the surroundings of Mariager (Denmark); both specimens are from the McGugan Collection, University of Calgary.

Fig. 36. Detail micrographs of two specimens of *Bolivinoides gigantea* Hiltermann and Koch, 1950 from uppermost Maastrichtian from the surroundings of Mariager (Denmark) showing the secondary ornamentation on one lateral side of the test (1) and porosity of the penultimate chamber (2); both specimens are from the McGugan Collection, University of Calgary.

1992, p. 102, pl. 20, fig. 7. Khalil 1998, fig. 8: 5. Howe et al. 2003, pl. 8, figs 35-36. Anan 2011a, p. 301, pl. 1, fig. 6. Anan 2011b, p. 137, pl. 1, fig. 6. Dubicka and Peryt 2016, p. 84, fig. 7: 6.

Bolivinoides senonicus Yeremeyeva. Yeremeyeva and Belousova 1961, p. 86, pl. 24, fig. 7.

Bolivinoides decoratus giganteus Hiltermann and Koch. Hiltermann and Koch 1962, p. 315, pl. 46, fig. 8. Hiltermann 1963, pl. 4, figs 1-5. Kiesel and Trümper 1965, p. 157, pl. 12, fig. 7. Medizza 1967, p. 22, pl. 1, figs 12-14. Hanzliková 1969, p. 31, pl. 5, fig. 9. Hanzliková 1972, p. 81, pl. 18, fig. 14. Koch 1977, p. 63, pl. 12, fig. 1. Salaj 1988, pl. 1, fig. 15. Frenzel 2000, p. 137, pl. 25, fig. 6. Naidin and Beniamovski 2006, pl. 3, fig. 14. Ohmert 2011, pl. 3, fig. 20.

Bolivinoides ellipsodecorata GOEL 1965, p. 81, pl. 7, fig. 13.

Bolivinoides draco miliaris Hiltermann and Koch. Nestler 1982, fig. 23. Schönfeld 1990, p. 84, pl. 3, fig. 10. Khalil 1998, fig. 8: 2, 8. Ismail 2012, p. 38, pl. 3, fig. 7.

Bolivinoides decoratus (Jones). Loeblich and Tappan 1987, pl. 551, figs 8-9.

Bolivinoides draco draco (Marsson). Ismail 2012, p. 38, pl. 3, fig. 6.

Transitional form between *Bolivinoides intermedius* and *B. giganteus*. Dubicka and Peryt 2016, fig. 7: 1-2.

Diagnosis. Large-sized species of *Bolivinoides* with the CBE of the successive chambers of one row in continuity forming long secondary ornamentation structures.

Description. Test consists of the proloculus followed by chambers alternately added with respect to the test growth axis resulting in a biserial arrangement. Chambers are low and elongate, with the elongation axis tilted and forming an acute angle with the test growth axis along a posterior-anterior direction. Chambers increase in size as added throughout the ontogenetic development, resulting in a test with flaring aspect as seen in lateral view; chamber rate of size increase is constant. The last-formed five to six pairs of last-formed chambers present well-developed narrow backward extensions that attach to the previous chamber of the row and are separated by wide inter-CBE spaces; frequently they attach to the backward extensions of the previous chamber of the row, forming elongate ridge-like structures with continuous appearance that can run to the central portion of the test lateral sides. There are different types of backward extensions, as follows: one apertural CBE of medium size, one peripheral CBE that is the smallest and ten to fourteen median CBE,

which are symmetrically arranged on the two lateral sides of one chamber; therefore, there are five to seven median CBE on one lateral side of one chamber. There is a distinct decrease in the median CBE of one chamber on a direction from the aperture to periphery; the increase in the number of median CBE through the ontogenetic development is another distinct trend in the individuals of this species. The M1 CBE is the largest of the backward extensions of one chamber. Test is compressed in edge view, commonly with the two lateral sides nearly parallel in the adult portion of the test; there is a slow increase in the rate of chamber increase in thickness. Periphery is angular, with a weak keel that has a nodular or interrupted appearance that confers this species a weakly keeled aspect; the weak keel can be observed in the lateral views as narrow rim-like structures at the periphery. Aperture has an elongate shape and is situated on the apertural side of the last-formed chamber; the aperture elongation axis corresponds to that of the chamber. Periapertural structure consists of one delicate imperforate rim, which is rarely preserved; the rim borders the aperture on one side and extends towards the chamber interior as a short toothplate. Test wall is calcitic, hyaline, simple and perforate. Primary ornamentation is completely absent and therefore, the chamber surface appears smooth. Secondary ornamentation consists of vestigial CBE that occur as structures with circular or elongate outline in the posterior third of the test lateral sides; the weak peripheral keel is also a structure that pertains of secondary ornamentation for it results from the alignment and overlapping of the peripheral CBE. Pores are simple, with a circular and more rarely irregular and elongate outline, and a measured diameter or maximum dimension of 0.0006-0.0013 mm; the backward extension surface of the last-formed chambers can lack porosity.

Remarks. *Bolivinoidea gigantea* is the largest species of this genus. It differs from *B. miliaris* mainly by having the secondary ornamentation structures restricted to the posterior third of the test lateral sides, a higher degree of alignment of the CBE of the successively added chambers in one row and a slightly higher number of median CBE in the last-formed chamber (five to seven median rather than five to six). In general the separation between the two taxa requires observations carried out on large numbers of specimens.

Stratigraphical range: Maastrichtian.

Geographical distribution. EU (Belgium, Czech Republic, Denmark, France, Germany, Italy, Poland, The Netherlands), Africa (Egypt, Libya, Tunisia), Asia (Kazakhstan, Russia), Australia, USA (New Jersey, Texas), East Indian Ocean, and North Atlantic Ocean.

Bolivinoidea draco (Marsson, 1878)
(Figs. 37 and 38)

Synonymy

Bolivina draco MARSSON 1878, p. 157, pl. 3, fig. 25.
Bolivina draco Marsson. Egger 1899, p. 47, pl. 16, figs 14-16.
Franke 1928, p. 151, pl. 14, fig. 2.

Fig. 37. Two specimens of *Bolivinoidea draco* (Marsson, 1878) from the lower Maastrichtian of the Rügen Island (Germany) (1) and upper Maastrichtian of the North Atlantic Ocean (Orphan Knoll) (2). 1: specimen from the C.G. Ehrenberg Collection (Museum of Natural History, Berlin), Sample 2404a. 2: specimen from Sample 12-111A-11-6, 116-117 cm.

Fig. 38. Detail micrographs of one specimen of *Bolivinoidea draco* (Marsson, 1878) from the lower Maastrichtian of the Rügen Island (Germany) showing the secondary ornamentation (1a, 1c) and peripheral keel (1b); specimen from the C.G. Ehrenberg Collection (Museum of Natural History, Berlin), Sample 2404a.

Bolivina rhomboidea CUSHMAN 1926a, p. 19, pl. 2, fig. 3.
Bolivinoidea rhomboidea (Cushman). Cushman 1927c, p. 158, pl. 28, fig. 12. Dampel 1934, p. 14, pl. 1, fig. 9. Cushman 1946, p. 113, pl. 48, fig. 15. Wicher 1949, p. 85, pl. 5, fig. 6, fig. 7.
Bolivinoidea rhomboidalis (Cushman). Cushman 1927b, pl. 13, fig. 6. Cushman 1950, pl. 21, fig. 9.
Bolivinoidea draco (Marsson). White 1929, p. 45, pl. 5, fig. 2. Galloway 1933, pl. 32, fig. 11. Kikoïne 1948, p. 21, pl. 2, fig. 1. Hofker 1957, fig. 321: a-b. Montanaro Gallitelli 1957, p. 145, pl. 33, figs 14-16. Sacal and Debourle 1957, p. 14, pl. 3, fig. 18. Eternod Olvera 1959, p. 80, pl. 2, figs 10, 12. Bykova 1959, fig. 1: 3. Maslakova 1959, p. 115, pl. 14, figs 7-8. Rauzer-Chernousova and Fursenko 1959, fig. 835A. Vassilenko 1961, p. 201, pl. 41, fig. 2. Kavary and Frizzell 1963, p. 29, pl. 3, fig. 30. Loeblich and Tappan 1964, fig. 435: 1-2. Baranovskaya and Bulinnikova 1964, p. 294, pl. 64, fig. 9. Hofker 1966, pl. 13, figs 34-36. Stenestad 1968, pl. 2, fig. 3. Bate and Bayliss 1969, pl. 4, fig. 4. Hanzliková 1969, p. 32, pl. 5, fig. 11. Barr 1970, p. 646, pl. 99, figs 2-3. Hanzliková 1972, p. 81, pl. 19, figs 10-11. Vaptzarova 1974, p. 47, pl. 4, figs

23-25. Govindan 1977, p. 473, pl. 2, fig. 15, text-fig. 10: n-o. Petters 1977, p. 1032, pl. 1, fig. 1. Rozumeiko 1978, p. 138, pl. 20, fig. 7. Kaptarenko-Chernousova et al. 1979, p. 145, pl. 55, fig. 11. Dailey 1983, pl. 2, fig. 8. Liszkowa and Morgiel 1985, pl. 2, fig. 5. Robaszynski et al. 1985, pl. 5, fig. 4. Butt 1986, pl. 1, fig. 2. Loeblich and Tappan 1987, pl. 551, figs 3-7. Aliev et al. 1988, p. 106, pl. 11, fig. 8. Hart et al. 1989, p. 324, pl. 7.4, fig. 13. Martínez 1989, pl. 1, fig. 9. Nomura 1991, pl. 1, fig. 5. Gawor-Biedowa 1992, p. 101, pl. 20, fig. 8. Gasiński 1993, pl. 1, fig. 7. Hewaidy and El Ashwah 1993, pl. 4, fig. 12. Ziko et al. 1993, fig. 4: 13. Boltovskoy et al. 1995, p. 133, pl. 1, fig. 4. Nomura and Brohi 1995, pl. 1, fig. 11. Revets 1996, pl. 4, figs 1-4. Widmark 1997, p. 34, fig. 13: F. Koutsoukos and de Klasz 2000, p. 397, fig. 6: 7. Howe et al. 2003, pl. 8, figs 31-32. Pérez Estrada et al. 2007, pl. 1, fig. 1. Alegret Badiola 2008, fig. 5: a. Anan 2011a, p. 301, pl. 1, fig. 6. Anan 2011b, p. 136, pl. 1, fig. 5. Jaff et al. 2014, pl. 1, fig. 10-12, pl. 2, figs 6-7. Michalski et al. 2016, fig. 9: B. Dubicka and Peryt 2016, p. 83, fig. 6: 6-10. Dubicka in Walaszczyk et al. 2016, text-fig. 5: 8.

Bolivina (Bolivinoides) draco (Marsson). Kalinin 1937, p. 46, pl. 5, figs 77-78.

Bolivinoides draco draco (Marsson). Hiltermann and Koch 1950, p. 598, figs 1 (72-73), 2-4 (52-54, 58-60) 5 (53, 69-70). Noth 1951, p. 63, pl. 9, fig. 10. Papp and Küpper 1953, p. 44, pl. 2, fig. 4. Edgell 1954, p. 73, pl. 13, figs 1-3, pl. 14, figs 1-3. Pożaryska 1954, p. 253, fig. 3. Reiss 1954, p. 155, pl. 29, figs 1-3. Bukowy and Georch 1956, pl. 18, fig. 16. Said and Kenawy 1956, p. 140, pl. 3, fig. 41. Pokorný 1958/1963, fig. 329: right figure. Ansary et al. 1962, pl. 4. Hiltermann and Koch 1962, p. 318, pl. 46, fig. 10. Hiltermann 1963, pl. 4, figs 16-19. Van Hinte 1963, p. 105, pl. 14, fig. 3. Beckmann and Koch 1964, p. 45, pl. 6, fig. 24. Pringgoprawiro 1965, p. 34, pl. 7, fig. 3. Medizza 1967, p. 23, pl. 2, figs 1-2. Scheibnerová 1968, p. 50, pl. 2, figs 8-10. Ansary and Tewfik 1969, pl. 2, fig. 1. Al-Shaibani 1971, p. 109, pl. 2, fig. 6. Beckmann 1978, p. 765, pl. 2, figs 12, 18-19. Neagu 1979, pl. 53, fig. 39. Salaj 1980, pl. 33, fig. 8. Swiecicki 1980, p. 162, pl. 5, fig. 19. Almogi-Labin et al., 1990, pl. 1, fig. 5. Bolli et al. 1994, p. 129, fig. 35: 1. Ismail and El Saadany 1995, p. 195, fig. 11: 6. Speijer and van der Zwaan 1995, p. 48, pl. 1, fig. 3. Ayyad et al. 1997, fig. 8: j. Frenzel 2000, p. 138, pl. 25, figs 12-14, text-fig. 75. Ismail 2002, pl. 1, figs 25-26. Abdelghany 2003, fig. 8: 5. El-Nady 2006, p. 678, pl. 1, figs 12-14. Ohmert 2011, pl. 3, fig. 24. Holbourn et al. 2013, p. 74, fig. 1. Shabestari et al. 2014, pl. 3, fig. 5. Anan 2017, p. 4, pl. 1, fig. 2.

Bolivinoides doreeni Finlay. Hofker 1956a, p. 70, fig. 48.

Bolivinoides draco doreeni Finlay. Ansary et al. 1962, pl. 4. Ismail and El Saadany 1995, p. 195, fig. 11: 5. El-Nady 2006, p. 675, pl. 1, fig. 11.

Bolivinoides draco doreeni Finlay. Martínez-Pardo 1965, text-fig. 3.

Bolivinoides miliaris (Hiltermann and Koch). Felder and Bless 1993, fig. 6: Q-R. Jaff et al. 2014, pl. 1, figs 15-16, pl. 2, figs 9-10.

Bolivinoides draco aegyptiacus ANAN 2017, p. 3, pl. 1, fig. 1.

Diagnosis. Species of *Bolivinoides* with peripheral keel and two longitudinal ridges in the posterior and central portions of the lateral sides.

Description. Test consists of the proloculus followed by chambers added alternately with respect to the test growth

axis forming a biserial arrangement. Chambers are low and elongate, with the elongation axis tilted and forming an acute angle with the test growth axis along a posterior to anterior direction. Chambers increase in size as added through ontogeny at a constant rate, resulting in a test sub-rhombical outline with flaring aspect in the direction of growth. The last-formed three to four pairs of chambers present well-developed and narrow backward extensions, which are attached to the previous chambers of the row. The backward extensions are separated by wide inter-CBE spaces. There are different types of CBE, as follows: One apertural CB, which is relatively small, one peripheral CBE that is also small and frequently fused with the similar such structures of the previous chambers in a row, and six to eight median CBE that are symmetrically arranged on the two lateral sides of a chamber; therefore, there are three to four median CBE on one later side of one chamber. There is a well-marked decrease in the median CBE size in one chamber on a direction from the aperture towards periphery; the M1 CBE is the largest backward extension of one chamber. Test is compressed in edge view, with a low rate of chamber increase in thickness resulting in distinctly diverging lateral sides in the direction of test growth. Periphery is angular and keeled; the keel is often continuous and in lateral view forms a rim-like structure at the periphery. Aperture has elongate outline and is situated on the apertural face of the last-formed chamber; the aperture's elongation axis corresponds to that of the chamber elongation. Peripheral structure consists of one delicate imperforate rim, which is more rarely preserved, and borders the aperture on one side; it extends towards the chamber interior as a short toothplate. Test wall is calcitic, hyaline and perforate. Primary ornamentation is absent in the representatives of this species. The secondary ornamentation consists primarily of vestigial CBE that are aligned and sometimes fused to form two well-developed ridges that run parallel to and on each side of the growth axis on the test lateral sides. Rare isolated vestigial CBE can occur in the posterior third of the test lateral sides and especially towards the periphery. Peripheral keel is another feature pertaining of the secondary ornamentation. Test wall is simple on most of the chambers, but incipiently reticulately-ridged wall with wavy surface can occur on the apertural face or portions of it of some specimens. Pores are circular, with a measured maximum diameter of 0.0006-0.0027 mm; pores with an irregular or elongate outline are extremely rare. Chamber backward extensions are imperforate.

Remarks. *Bolivinoides draco* is the only species of the genus that evolved a true keel, which in most specimens has a continuous appearance; it is also characterized by the two well-developed ridges with continuous aspect that are parallel to and on each side of the growth axis.

Stratigraphical range: Maastrichtian.

Geographical distribution. EU (Austria, Belgium, Czech Republic, Denmark, France, Germany, Italy, Poland, Slovakia, The Netherlands), Africa (Egypt, Libya, Tanzania, Tunisia), Asia (Azerbaijan, India, Iran, Kazakhstan, Russia), Australia, Caribbean Region (Cuba, Trinidad), Mexico, Middle East (Iraq, Israel), South America (Brazil, Colombia), UK (England), Ukraine, USA (California, New Jersey), East Indian Ocean, North Atlantic Ocean, and South Atlantic Ocean.

Bolivinoides regularis Reiss, 1954
(Figs. 39 and 40)

Synonymy

Bolivinoides delicatula regularis REISS 1954, p. 157-158, pl. 31, figs 1-4.

Bolivinoides curta REISS 1954, p. 158, pl. 30, figs 15-16.

Bolivinoides decorata delicatula Cushman. Pożaryska 1954, p. 255, fig. 7.

Bolivinoides regularis Reiss. Bieda 1958, p. 36, fig. 9. Hanzliková 1972, p. 83, pl. 19, fig. 15. Neagu 1979, pl. 53, fig. 40. Liszkowa and Morgiel 1985, pl. 2, fig. 4. Felder and Bless 1993, fig. 6: O-P.

Bolivinoides delicatulus curtus Reiss. Hiltermann 1963, pl. 3, figs 4, 8-9. Beckmann and Koch 1964, p. 43, pl. 6, figs 1-5, 7-10. Bolli et al. 1994, p. fig. 34: 36.

Bolivinoides delicatulus regularis Reiss. Hiltermann 1963, pl. 3, figs 7, 11-12, 20. McGugan 1974, pl. 1, fig. 1. Koch 1977, p. 58, pl. 13, figs 4-5.

Bolivinoides delicatulus n. subsp. aff. *delicatulus*. Beckmann and Koch 1964, p. 42, pl. 5, figs 8-9, 13-14.

Bolivinoides delicatulus (Cushman). Cati 1964, p. 238, pl. 38, fig. 21.

Bolivinoides sidestrandensis Barr. Petters 1977, p. 1034, pl. 1, fig. 5. Gawor-Biedowa 1992, p. 107, pl. 20, fig. 1. Frenzel 2000, p. 144, pl. 25, fig. 1. Dubicka and Peryt 2016, p. 82, fig. 4: 7-10. Dubicka in Walaszczyk et al. 2016, text-fig. 5: 12.

Bolivinoides curtus Reiss. Anan and Hamdan 1992, p. 205,

Fig. 40. Detail micrographs of two specimens of *Bolivinoides regularis* Reiss, 1954 from lower Maastrichtian of the surroundings of Lüneburg (Germany) showing the weakly developed secondary ornamentation on the test lateral sides (1, 2a) and primary ornamentation consisting of rare scattered pustules over the chambers of the earlier portion of the test (2b); both specimens are from the McGugan Collection, University of Calgary.

text-fig. 5. El-Nady 2006, p. 674, pl. 1, fig. 4. Anan 2011b, p. 134, pl. 1, fig. 1.

Bolivinoides peterssoni Brotzen. Peryt 2000, pl. 1, fig. 3.

Bolivinoides compressus Reiss. El-Nady 2006, p. 674, pl. 1, fig. 3.

Transitional forms between *Bolivinoides decoratus* to *B. sidestrandensis*. Dubicka and Peryt 2016, fig. 5-6.

Diagnosis. Species of *Bolivinoides* with narrow CBE separated by narrow inter-CBE spaces and primary ornamentation consisting of scattered pustules developed over the chambers of the posterior portion of the test.

Description. Test consists of the proloculus followed by chambers alternately added with respect to the test growth axis resulting in a biserial arrangement. Chambers are low and elongate, with the elongation axis forming an acute angle with that of the test growth along a posterior-anterior direction. There is a gradual increase in the chamber size through ontogeny, which results in a test flaring outline in the direction of growth. The last-formed four to seven pairs of chambers present well-developed and narrow backward extensions, which attach to the previous chamber of the row; the backward extensions are separated by inter-CBE spaces of around the same width with the adjacent CBE. Each chamber has different types of backward extensions, as follows: one apertural CBE, which is the largest in most of the studied specimens, one peripheral CB that is the smallest, and eight to ten median CBE symmetrically arranged on the two lateral sides; therefore, there are four to five median CBE on one lateral side of one chamber. There is a distinct decrease of the median CBE of one chamber on a direction from the aperture to periphery. Test is compressed in edge view, with nearly parallel sides in the adult stage, due to the very low increase in chamber thickness through ontogeny. Periphery is subrounded to subangular and simple, without peripheral structures. Aperture is elongate and situated on the apertural face of the last-formed chamber; aperture elongation axis matches that of the chamber elongation. Periapertural structure consists of one delicate and imperforate rim, which is rarely preserved and border the aperture on one side; the periapertural rim extends within the chamber interior as a short toothplate. Test wall is calcitic, hyaline, simple and perforate. Primary ornamentation consists of small-sized

Fig. 39. Three specimens of *Bolivinoides regularis* Reiss, 1954 from lower Maastrichtian of the surroundings of Lüneburg (Germany); all the specimens are from the McGugan Collection, University of Calgary.

scattered dome-like pustules that occur sporadically over the earlier portion of the test lateral sides; the measured diameter of these structures is of 0.0049–0.0159 mm. The secondary ornamentation is poorly developed and consists of vestigial CBE over the posterior third of the test lateral sides of some specimens. The CBE present a distinct converging pattern in the posterior half of the test lateral sides and in some specimens form a weak longitudinal ridge-like structure that corresponds to the test growth axis. Most of the pores are circular, rarely with irregular outline, and with a measured diameter or maximum dimension respectively of 0.0003–0.0007 mm.

Remarks. *Bolivinoides regularis* differs from *B. delicatula* mainly by the reduced primary ornamentation: the former has smooth chamber surface with scattered dome-like pustules occurring over the posterior third of the test lateral sides of some specimens, whereas the latter presents a primary ornamentation consisting of smaller scattered pustules and short longitudinal costae; in addition, in the former species the CBE are narrower and separated by wider inter-CBE spaces. This species differs from *B. sidestrandensis* mainly by having longer and slender tests, and more prominent CBE. Reiss (1954) included the specimens of this species within two taxonomic units: *Bolivinoides delicatula regularis* and *B. curta*; these are considered synonyms representing different test varieties of the same species. The former is more elongate and most likely includes the gerontic specimens, whereas the latter are slightly larger adult specimens.

Stratigraphical range: Maastrichtian–Paleocene.

Geographical distribution. Middle East (Israel), Africa (Egypt), Caribbean Region (Gulf of Mexico, Trinidad), EU (Belgium, Czech Republic, Germany, Italy, Poland, The Netherlands), Mexico, UK (Northern Ireland), and USA (Alabama, New Jersey).

Bolivinoides doreeni Finlay, 1940
(Figs. 41 and 42)

Synonymy

Bolivinoides doreeni FINLAY 1940, p. 452, pl. 63, figs 56–58.

Bolivinoides decorata (Jones). Sellier de Civrieux 1952, p. 272, pl. 6, figs 19–20.

Bolivinoides draco doreeni Finlay. Reiss 1954, p. 155, pl. 29, figs 4–7. Said and Kenawy 1956, p. 140, pl. 3, fig. 42. Scheibnerová 1968, p. 52, pl. 3, figs 1–5. Strong and Edwards 1979, fig. 6. Khalil 1998, fig. 8: 3. Anan 2017, p. 4, pl. 1, fig. 3.

Bolivinoides draco dracoformis VASSILENKO 1961, p. 190, pl. 40, fig. 10, pl. 41, fig. 3.

Transitional specimen *Bolivinoides draco miliaris*–*B. draco draco*. Hiltermann 1963, pl. 4, fig. 20.

Bolivinoides decoratus giganteus Hiltermann and Koch. Pringgoprawiro 1965, p. 34, pl. 7, fig. 4.

Bolivinoides draco draco (Marsson). Koch 1977, p. 57, pl. 12, figs 2–3. Schönfeld 1990, p. 78, pl. 3, figs 1–2. Khalil 1998, fig. 8: 1, 4.

Bolivinoides draco (Marsson). Villain 1977, p. 69, pl. 5, fig. 13. Van Morkhoven et al. 1986, p. 378, pl. 124, figs 1–2.

Fig. 41. Two specimens of *Bolivinoides doreeni* Finlay, 1940 from the lower Maastrichtian of the surroundings of Lüneburg (Germany); both specimens are from the McGugan Collection, University of Calgary.

Bolivinoides doreeni Finlay. Anan 2011b, p. 136, pl. 1, fig. 4. *Bolivinoides giganteus* Hiltermann and Koch. Dubicka and Peryt 2016, p. 84, fig. 7: 3–4, 6–9.

Diagnosis. Species of *Bolivinoides* with two parallel and discontinuous ridges in the central portion of the lateral sides of the test.

Description. Test consists of the proloculus, which is followed by chambers alternately added with respect to the test growth axis forming a biserial arrangement. Chambers are low and their elongation axis forms an acute angle with that of the test growth along a posterior–anterior direction. Chambers increase in size through ontogeny resulting in a test flaring outline in the direction of growth. The last-formed four to five pairs of chambers present well-developed and narrow backward extensions that attach to the previous chamber

Fig. 42. Detail micrographs of one specimen of *Bolivinoides doreeni* Finlay, 1940 from the lower Maastrichtian of the surroundings of Lüneburg (Germany) showing pore distribution over the test last-formed chamber (1a) and pore shape (1b); specimen from the McGugan Collection, University of Calgary.

of the row and are separated by wide inter-CBE spaces; frequently the inter-CBE spaces are slightly wider than the adjacent CBE. There are different types of CBE, as follows: one apertural CBE, which is the largest, one peripheral CBE that is the smallest, and ten to twelve median CBE that are symmetrically distributed on the two lateral sides of one chamber; therefore, there are five to six median CBE on one lateral side of one chamber. There is a slow decrease in size of the median CBE of one chamber along a direction from the aperture towards periphery. Sutures are straight or slightly curved anteriorly. Test is compressed in edge view, with a slow increase of chamber thickness throughout ontogeny. Periphery is acute and with one keel resulted from the peripheral CBE of the successively added that fuse; the peripheral keel can also be observed in lateral view as a rim-like structure that is better developed in the posterior side of the test. Aperture is elongate and situated on the apertural side of the last-formed chamber; the aperture's axis of elongation matches that of chamber elongation. Periapertural structure consists of a delicate and imperforate rim, which is rarely preserved and borders the aperture on one side; the periapertural rim extends towards the chamber interior as a short toothplate. Test wall is calcitic, hyaline and perforate. Primary ornamentation is completely absent in this species. Secondary ornamentation consists of vestigial CBE and occur especially in the posterior third of the test's two lateral sides. Two parallel ridges situated on the two sides of the test growth axis occur on the lateral sides; the ridges have a discontinuous aspect. The peripheral keel is another feature of secondary ornamentation. Test wall is simple on most of the test chambers and occasionally incipiently reticulately-ridged on the apertural side of the last-formed chambers. Most of the pores are circular and rarely present an irregular outline; the measured diameter or maximum dimension respectively is of 0.0008-0.0033 mm. In most specimens the CBE are imperforate.

Remarks. *Bolivinoides doreeni* differs from *B. draco* mainly by having less developed secondary ornamentation: the two longitudinal ridges situated on each of the two test lateral sides have a discontinuous rather than continuous appearance, and the peripheral keel is weaker. There are more median CBE on one lateral side of one chamber in *B. doreeni* (five to six) when compared to *B. draco* (four to five). In addition, the tests included in the former species are much larger when compared to those included in the latter. This species differs from *B. gigantea* mainly by having more prominent longitudinal ridges on each of the two test lateral sides and peripheral keel.

Stratigraphical range: Maastrichtian.

Geographical distribution. New Zealand, Africa (Egypt), Asia (Kazakhstan), EU (Austria, France, Germany, Poland, Slovakia), Gulf of Mexico, Middle East (Israel), South America (Venezuela), East Indian Ocean, and North Atlantic Ocean.

Bolivinoides vistulae Pożaryska, 1954 – emended
(Fig. 43)

Synonymy

Bolivinoides vistulae POŻARYSKA 1954, p. 253, fig. 2.
Bolivinoides vistulae Pożaryska. Pożaryska 1965, p. 101, pl. 14, fig. 1. Dubicka and Peryt 2016, p. 88, fig. 10: 6-10.

Fig. 43. Two specimens of *Bolivinoides vistulae* Pożaryska, 1954 from the upper Maastrichtian of the North Atlantic Ocean (Orphan Knoll). 1: specimen from Sample 12-111A-11-2, 115-116 cm. 2: specimen from Sample 12-111A-11-4, 116-117 cm.

Michalski et al. 2016, fig. 9: J-L, fig. 10: J, fig. 11: F. Dubicka in Walaszczyk et al. 2016, text-fig. 5: 14.

Bolivinoides draco draco (Marsson). Sliter 1968, p. 88, pl. 12, fig. 17.

Bolivinoides dentatus GAWOR-BIEDOWA 1992, p. 100, pl. 18, figs 8-9.

Bolivinoides paleocenica (Brotzen). Khalil 1998, fig. 8: 7.

Bolivinoides delicatus Cushman. Hofmann et al. 1999, pl. 1, fig. 11.

Bolivinoides paleocenicus (Brotzen). El-Nady 2006, p. 679, pl. 1, fig. 19.

Bolivinoides decoratus (Jones). Anan 2011b, p. 135, pl. 1, fig. 2.

Bolivinoides sp. Remin et al. 2012, fig. 3: C.

Diagnosis. Species of *Bolivinoides* with a wide peripheral rim and dentate periphery in the adult portion of the test.

Description. Test consists of the proloculus followed by chambers alternately added with respect to the test growth axis resulting in a biserial arrangement. Chambers are low and elongate, with the elongation axis forming an acute angle with that of the test growth along a posterior-anterior direction. Chambers increase in size as added throughout ontogeny, resulting in a test outline with flaring aspect in the direction of growth as see in lateral view. The last-formed five to six pairs of chambers present backward extensions that attach to the previous chamber of the row; the backward extensions are wide and separated by distinct inter-CBE spaces that increase in width through ontogeny. Each chamber presents different types of backward extensions, as follows: one apertural CBE, which often is the largest, one peripheral CBE that in gerontic specimens is distinctly detached from the periphery, and six to eight median CBE symmetrically distributed on the two lateral sides of one chamber; therefore, there are three to four CBE on each lateral side of one chamber. There is a decrease in the median CBE of one chamber on a direction from aperture towards periphery, and the M1 CBE is in some specimens the largest of the backward extensions of one chamber. Sutures

are straight to gently curved anteriorly, and often obscured by the backward extensions of the subsequently added chambers of the row. Test is compressed in edge view, with slightly divergent sides due to the slow increase in chamber thickness through ontogeny. Periphery is subrounded and simple, without peripheral structures; in lateral view the periphery can have a weakly dentate aspect due to the detached peripheral CBE. Aperture has an elongate outline and is situated on the apertural side of the last-formed chamber; aperture's elongation axis corresponds to that of the chamber. Periapertural structure consists of one imperforate rim, which borders the aperture on one side and extends towards the chamber interior as a short toothplate. Test wall is calcitic, hyaline and perforate. There is no primary ornamentation in this species and therefore, the chamber surface has a smooth appearance. Secondary ornamentation consist of rare vestigial CBE that occur in the posterior portion of the test lateral sides; notably, these structures that can have a circular or elongate outline, are often difficult to recognize. Pores are have a circular outline, more rarely irregular, and with a measured diameter or maximum dimension respectively of 0.0004-0.0009 mm.

Remarks. In the original description, Pożaryska (1954) noted that this species resembles *B. peterssoni*, of which it differs by having more prominent CBE on the test lateral side; these morphological differences were practically accepted in unchanged form in the subsequent works where *B. vistulae* was described (Pożaryska, 1965; Dubicka and Peryt, 2016). The nature of the emendation proposed herein is that the two species differ by the detached peripheral CBE in *B. vistulae*, one feature that is not encountered in *B. peterssoni*. In this context we can draw the conclusion that the ancestor-descendant relationship between *B. peterssoni* and *B. vistulae* inferred by Dubicka and Peryt (2016) is unclear and, according to their definition of the two species, can be confused for ontogenetic differences within the same species; such ambiguity is avoided through the emendation of *B. vistulae* as proposed herein.

Stratigraphical range: Upper Maastrichtian-lower Paleocene.

Geographical distribution. EU (Austria, Poland), Africa (Egypt), USA (California), and North Atlantic Ocean.

Genus *Trachelinella* Montanaro Gallitelli, 1956

Type species: *Bolivina watersi* Cushman, 1927a.

Synonymy

Trachelinella MONTANARO GALLITELLI 1956, p. 38.

Trachelinella Montanaro Gallitelli. Montanaro Gallitelli 1957, p. 150.

Trachelinella Montanaro Gallitelli. Loeblich and Tappan 1964, p. 736.

Trachelinella Montanaro Gallitelli. Loeblich and Tappan 1987, p. 501.

Trachelinella Montanaro Gallitelli. Georgescu et al. 2011, p. 122. Emended.

Diagnosis. Bolivinoidids with the aperture in terminal position and produced on a short neck and test surface ornamented with pore mounds that can be scattered or fused.

Description. Test is biserial throughout consisting of the proloculus followed by chambers alternately added with respect to the test growth axis. Earlier chambers are subglobular and

the last-formed ones elongate and with backward extensions; the plane of chamber addition can present a rotation pattern around the test growth axis. There is one peripheral CBE and one to two median CBE on each lateral side of one chamber; some of the chambers of the test juvenile portion do not have backward extensions. Sutures are depressed, curved anteriorly and often obscured by the backward extensions and/or ornamentation structures of the successively added chamber of the row. Test is laterally compressed, with subangular to angular periphery and simple, without peripheral structures. Aperture is terminal and with elongate outline; it is produced on a short neck and surrounded by an imperforate rim. Test wall is calcitic, hyaline, simple and perforate. Primary ornamentation consists of well-developed pore mounds with circular outline that can fuse and form ornamented platforms. Pores present a circular or more rarely elliptical shape.

Remarks. *Trachelinella* differs from *Bolivinoides* mainly by having the aperture in terminal position instead at the base or on the apertural face of the last-formed chamber and periapertural structure consisting of a rim that completely surrounds the aperture rather than an imperforate rim that borders the aperture on one side. In addition, the backward extensions of *Trachelinella* have in general subtriangular shape, occasionally present an irregular appearance, whereas in *Bolivinoides* they are in general more numerous and posteriorly rounded.

Species included

Trachelinella watersi (Cushman, 1927a)

Stratigraphical range: Campanian-Maastrichtian.

Trachelinella watersi (Cushman, 1927a)

(Figs. 44 and 45)

Synonymy

Bolivina watersi CUSHMAN 1927a, p. 88, pl. 12, fig. 6.

Bolivina watersi Cushman. Cushman 1946, p. 128, pl. 53, fig. 18.

Bolivinoides watersi (Cushman). Reiss 1954, p. 156, pl. 29, fig. 8.

Trachelinella watersi (Cushman). Montanaro Gallitelli 1956, pl. 7, figs 8-10. Montanaro Gallitelli 1957, p. 150, pl. 34, fig. 13. Perlmutter and Todd 1965, p. 15, pl. 2, fig. 12. Georgescu et al. 2011, p. 122, pl. 4, figs 1-10.

Bolivina praecrenulata GAWOR-BIEDOWA 1992, p. 96, pl. 17, fig. 6.

Bolivinoides granulatus Hofker. Dubicka and Peryt 2016, p. 86, fig. 9: 7.

Diagnosis. Species with biserial throughout test, aperture produced on a short neck and ornamentation consisting of pore mounds and pore-mounded structures.

Description. Test consists of the proloculus followed by chambers alternately added with respect to the test growth axis resulting in a biserial arrangement. Chambers are elongate, with the main axis tilted and forming an acute angle with the test growth axis along a posterior to anterior direction. Chambers increase in size throughout ontogeny at a constant rate and the plane in which they are added can rotate around the test growth axis. The last-formed seven to ten pairs of chambers present backward extensions. Chambers backward extensions are of different types, as follows: one peripheral CBE and two to four median CBE symmetrically distributed

Fig. 44. Three specimens of *Trachelinella watersi* (Cushman, 1927a) from the upper Maastrichtian of Texas. 1: specimen from Well Mullinax-3, Sample 42, 4-3, 7.15 m. 2: specimen from Well Mullinax-3, Sample 42, 4-3, 7.15 m previously illustrated by Georgescu et al. (2011, pl. 4, figs 8-9). 3: specimen from Well Mullinax-3, Sample 42, 4-3, 7.15 m.

on the two lateral sides of a chamber; therefore, there are one to two median CBE on a lateral side of one chamber. There is a distinct trend to increase the number of median CBE from one to two in the course of ontogenetic development. Test is compressed in edge view, with the chambers presenting a relatively low rate of increase in thickness. Periphery is subangular to angular and simple, without periapertural structures. Aperture has an elongate shape, is situated in terminal position and is produced on a short neck. Periapertural structure consists of one imperforate rim that completely surrounds the aperture; there are no indications on the presence of a (short) toothplate in this species. Test wall is calcitic, hyaline, simple and perforate. Primary ornamentation consists of well-developed circular pore mounds with a

measured diameter of 0.0029-0.0097 mm; most of the pore mounds occur as isolated and scattered structures, but they can fuse and form pore mounded ornamentation platforms especially towards the posterior portion of the chambers. Most of the pores present a circular outline, more rarely elliptical; the measured diameter is of 0.0012-0.0020 mm, whereas in the case of those pores with elliptical outline the measured longer axis is of 0.0027-0.0029 mm.

Remarks. The evolutionary occurrence of this species is not known with precision, which contrasts with the upper Campanian-Maastrichtian portion of its stratigraphical range that is well documented. Reiss (1954, pl. 29, fig. 8) figured one specimen from the lower Campanian of Israel; this specimen has a smooth wall appearance, which is completely different from that known from the specimens collected from the upper Campanian-Maastrichtian and illustrated with the aid of the SEM. Notably, in all the reports of *T. watersi* in which was used the same drawing technique as that of Reiss (1954) the chamber surface appears smooth (e.g., Cushman, 1927a; Cushman, 1946, etc); in contrast, the illustrations with the aid of the SEM reveal consistently the primary ornamentation consisting of pore mounds (Gawor-Biedowa, 1992; Georgescu et al., 2011). The conclusion that can be drawn from these data is that the re-examination of the Israeli material illustrated by Reiss (1954), this time with the aid of an SEM that was not in the current practice in the early 1950s, would only help in clarifying the stratigraphical position of the evolutionary occurrence of *T. watersi*. The specimen figured by Dubicka and Peryt (2016, fig. 9: 7) and assigned to *Bolivinoidea granulatus* by these authors, is most interesting for it has mixed features of *B. culverensis* (gross test architecture) and *T. watersi* (test ornamentation). It is hard to tell based only on this specimen if there is distinct species between *B. culverensis* and *T. watersi*; more material is necessary in order to clarify this aspect of evolutionary micropaleontology.

Stratigraphical range: Campanian-Maastrichtian.

Geographical distribution. USA (Mississippi, New York, Texas), EU (Poland), and Middle East (Israel).

Genus *Swiecickina* – n. gen.

Type species: *Bolivina paleocenica* Brotzen, 1948.

Name origin. Genus named after Dr. Anthony Swiecicki for his outstanding contributions to the biostratigraphy of Late Cretaceous foraminifera; to his name the suffix *-na* is added.

Diagnosis. Bolivinoidids with strongly compressed tests and ornamentation consisting of narrow ridges that can be distributed without a preferential pattern, anastomosed in the central portion of lateral sides, or extended over the chamber backward extensions.

Description. Test is biserial and consists of the proloculus followed by chambers alternately with respect to the test growth axis. Earlier chambers are subglobular and the last-formed ones elongate and with backward extensions. The chamber backward extensions can be weak to well-developed. There are two, more rarely four median CBE, which are symmetrically

Fig. 45. Detail micrographs of three specimens of *Trachelinella watersi* (Cushman, 1927a) from the upper Maastrichtian of Texas showing the ornamentation consisting of pore mounds on the test lateral sides (1, 3) and over the peripheral region (2). 2: specimen from Well Mullinax-3, Sample 42, 4-3, 7.15 m previously illustrated by Georgescu et al. (2011, pl. 4, fig. 10).

arranged on the two lateral side of the test; the peripheral CBE can be detached from the previous chamber of the row, resulting in a dentate aspect of the periphery. Aperture is elongate and situated on the apertural side of the last-formed chamber. Peripheral structure consists of one imperforate rim that borders the aperture on one side and extends in the umbilicus as a short toothplate. Test is laterally compressed, with parallel lateral sides in edge view; chamber surface is flat on the lateral sides of the test. Periphery is subangular and simple, without peripheral structures. Test wall is calcitic, hyaline, simple and perforate. Primary ornamentation consists of scattered pustules that can fuse and form narrow ridges on the lateral sides that can present an irregular arrangement, can anastomose towards the central part or begin from above and follow the direction of the CBE in the adult portion of the test.

Remarks. *Swiecickina* accommodates one lineage that shows the development of a dentate periphery in the adult portion of the test, and narrow ridges on the test lateral sides that in the early stage of the lineage are irregularly arranged and then line the sutures towards the central part of the lateral sides, not reaching the peripheral region. This genus differs from *Bolivinoides* mainly by having a flat chamber surface on the lateral side, dentate periphery in the adult portion of the test, less developed CBE, and ornamentation consisting of ridges that occurs on the test lateral sides. *Swiecickina* differs from *Aragonia* Finlay, 1939 mainly by occurrence of the CBE, which are absent in the latter genus.

Species included (in stratigraphical order)

Swiecickina clavatus (Plotnikova, 1967)

S. senonicus (Dain, 1961)

S. paleocenica (Brotzen, 1948)

Stratigraphical range: Uppermost Campanian-Paleocene.

Swiecickina clavatus (Plotnikova, 1967)

(Fig. 46)

Synonymy

Bolivinoides clavatus PLOTNIKOVA 1967, p. 96, pl. 15, fig. 6.

Bolivinoides clavatus Plotnikova. Kaptarenko-Chernousova et al. 1979, p. 145, pl. 55, fig. 2. Gawor-Biedowa 1992, p. 98, pl. 19, figs 10-11. Frenzel 2000, p. 134, pl. 24, figs. 3-6.

Diagnosis. Species of *Swiecickina* with the primary ornamentation consisting of scattered dome-like pustules that can fuse forming short ridges, which do not present a preferential ornamentation.

Description. Test consists of the proloculus followed by chambers alternately added with respect to the test growth chamber resulting in a biserial arrangement. Chambers increase gradually in size through ontogeny; the rate of chamber increase in size can vary but in general the adult portion consists of chambers with a low rate of width increase. The last-formed one or two pairs of chambers present two short median backward extensions with low relief; the two CBE are symmetrically distributed on the two sides of the chamber and therefore, there is one CBE on one lateral side of one chamber. A peripheral CBE, which is detached from the periphery, can occur in the last-formed chamber. Sutures are depressed, mostly straight in the earlier portion of the test and gently curved towards the direction of growth in the adult part. Test is compressed in edge view, with nearly parallel lateral sides due to the lack of increase of chamber

Fig. 46. One specimen of *Swiecickina clavatus* (Plotnikova, 1967) from the C.G. Ehrenberg Collection (Museum of Natural History, Berlin); specimen collected from the bulk sample 2344a (proximity of the Campanian/Maastrichtian), Moen Island (Denmark).

width through ontogeny; chamber surface on the lateral sides is flat. Periphery is subrounded and simple, without peripheral structures. Aperture has an elongate outline and is bordered by an asymmetrical imperforate rim that extends towards the chamber interior as a short toothplate. Test wall is calcitic, hyaline, dimple and perforate. Primary ornamentation consists of scattered dome-like pustules that can fuse and form ridges with irregular arrangement over the test lateral sides. Pores are mostly circular and simple, with a measured diameter of 0.0003-0.0006 mm.

Remarks. *Swiecickina clavatus* differs from any bolivinoidid species by the primary ornamentation consisting of scattered dome-like pustules that can fuse and form ridges, which present an irregular arrangement over the test lateral side.

Stratigraphical range: Upper upper Campanian-lower Maastrichtian.

Geographical distribution. Ukraine and EU (Germany, Poland).

Swiecickina senonicus (Dain, 1961)

(Figs. 47 and 48)

Synonymy

Bolivinoides senonicus DAIN 1961, p. 30, pl. 2, figs 8-9, text-fig. 2: 7.

Bolivinoides senonicus Dain. Baranovskaya and Bulinnikova 1964, p. 295, pl. 64, figs 10-12. Gorbenko 1974, p. 53, pl. 9, fig. 3. Kaptarenko-Chernousova et al. 1979, p. 147, pl. 55, fig. 9.

Aragonia sp. 1. Van Morkhoven et al. 1986, p. 374, pl. 122, figs 1-3.

Diagnosis. Species of *Swiecickina* with dentate periphery in the adult portion of the test and anastomosing ridges in the central portions of the test lateral sides.

Description. Test consists of the proloculus followed by chambers alternately added with respect to the test growth axis resulting in a biserial arrangement. Chambers are elongate and relatively low, with the elongation axis forming an acute angle with the test growth axis along a posterior to anterior direction. Chambers increase in size through ontogeny at various rates; there is a slow rate of size increase in the early portion of the test, and a rapid one in the adult portion resulting in a flaring test in the direction of growth. The last-formed one or

Fig. 47. Three specimens of *Swieickina senonicus* (Dain, 1961) from the lower Maastrichtian sediments of Yucatan outer shelf (DSDP Site 95). All the specimens are from Sample 10-95-13-1, 101.5-103 cm.

Fig. 48. Detail micrographs of three specimens of *Swieickina senonicus* (Dain, 1961) from the lower Maastrichtian sediments of Yucatan outer shelf (DSDP Site 95) showing the anastomosing ridges in the central part of the test lateral side over the central zigzag suture. All the specimens are from Sample 10-95-13-1, 101.5-103 cm.

two pairs of chambers present short and low-relief backward extensions. There are different types of backward extensions, as follows: one peripheral CBE that is frequently detached in the adult portion of the test conferring the periphery a well-marked dentate aspect, and two to more rarely four median CBE, which are symmetrically arranged on the two lateral sides of one chamber; therefore, there are one to more rarely two CBE on one lateral side of one chamber. Sutures are straight, rarely gently curved in the direction of growth. Test is compressed in edge view, with the two sides parallel due to the lack of chamber increase in width through ontogeny, especially in the adult portion of the test. Periphery is subrounded and simple, without peripheral structures. Aperture is situated on the apertural face of the last-formed chamber and has an elongate outline, with the elongation axis matching that of the respective chamber elongation. Peripheral structures consist of one asymmetrical imperforate rim, which extends towards the chamber interior as a short toothplate. Test wall

is calcitic, hyaline, simple and perforate. Primary ornamentation consists of ridges that anastomose in the central portions of the two lateral sides, extending from and following the direction of the CBE; there are rare specimens where the ridges occur on more than two pairs of chambers. Pores are mostly circular and simple, with a measured diameter of 0.0004-0.0007 mm.

Remarks. *Swieickina senonicus* differs from *S. clavatus* by having distinct dentate periphery in the adult portion of the test, more pronounced CBE and more prominent primary ornamentation, which consists of anastomosing ridges that can begin from the CBE and extend to the central portion of the lateral sides instead of ridges with irregular arrangement. This is the earliest species of the *Swieickina* lineage in which the development of the mixed features between *Bolivinoidea* and *Aragonia* is apparent.

Stratigraphical range: Lower to lower upper Maastrichtian.

Geographical distribution. Asia (Russia), Caribbean Region (Gulf of Mexico, Yucatan outer shelf), and Ukraine.

Swieickina paleocenica (Brotzen, 1948)
(Fig. 49)

Synonymy

Bolivina paleocenica BROTZEN 1948, p. 66, pl. 9, fig. 5.

Bolivinoidea paleocenica (Brotzen). Hiltermann 1952, p. 61, text-figs 4-5. Hofker 1952, fig. 4. Wicher 1953, fig. 1: 5. Reiss 1954, p. 157, pl. 30, figs 9-11. Hofker 1956a, p. 70, fig. 76. Hofker 1956b, fig. 23. Hofker 1957, p. 253, fig. 305 (c, e). Bieda 1958, p. 42, fig. 13.

Hofker 1960b, fig. 10. Hiltermann and Koch 1962, p. 317, pl. 50, fig. 15. Hofker 1962, text-fig. 15: B-C. Hofker 1966, pl. 10, fig. 75. Neagu 1979, pl. 53, fig. 41.

Bolivinoidea paleocenica (Brotzen). Hiltermann 1963, pl. 3, fig. 3. Medizza 1967, p. 26, fig. 2: 2-3. Barr 1970, p. 650, pl. 99, fig. 7. Szczuchura and Pożaryska 1974, p. 49, pl. 4, fig. 9. Govindan 1977, p. 470, pl. 1, figs 9-10, text-fig. 10: i-k. Koch 1977, p. 60, pl. 13, figs 1-3. Swieicki 1980, p. 181, pl. 6, figs 12-13. Yasuda 1986, p. 81, pl. 11, fig. 1. Hart et al. 1989, p. 324, pl. 7.5, fig. 4. Schönfeld 1990, p. 83, pl. 3, figs 7-8. Gawor-Biedowa 1992, p. 104, pl. 20, fig. 3. Hewaidy and El Ashwah 1993, pl. 4, fig. 16. Widmark 1997, p. 34, fig. 13: H. Frenzel 2000, p. 141, pl. 24, figs 7-9, text-fig. 76. Dubicka and Peryt 2016, p. 89, fig. 10: 11.

Bolivinoidea cf. paleocenica (Brotzen). De Klasz and de Klasz 1993, p. 314, pl. 1, fig. 11.

Bolivinoidea delicatulus Cushman. Alegret and Thomas 2007, pl. 1, fig. 3.

Fig. 49. Specimen of *Swiecickina paleocenica* (Brotzen, 1948) from the upper Maastrichtian sediments of the North Atlantic Ocean (Orphan Knoll), Sample 12-111A-11-6, 56-57 cm.

Diagnosis. Species of *Swiecickina* with dentate periphery in the adult portion of the test and well-developed ridges over CBE.

Description. Test consists of the proloculus followed by chambers alternately added with respect to the test growth axis resulting in a biserial arrangement. Chambers are low and elongate, with the elongation axis forming an acute angle with that of the test growth along a posterior-anterior direction. Chambers increase in size as added throughout the ontogeny resulting in a flaring test outline in the direction of growth. The last four to more rarely five pairs of chambers present short backward extensions, which are of different types, as follows: one median CBE, rarely two, and one peripheral CBE; the peripheral CBE is detached from the periphery in the adult portion of the test resulting in a dentate periphery. Sutures are depressed and straight or gently curved in the direction of growth. Test is compressed in edge view, with the two lateral sides nearly parallel due to the absence or slow rate of chamber increase in thickness. Periphery is subrounded and simple, without peripheral structures. Aperture is elongate and situated on the apertural face of the last-formed chamber. Periapertural structure consists of an asymmetrical imperforate rim, which is rarely preserved, that extends towards the chamber interior as a short toothplate. Test is calcitic, hyaline, simple and perforate. Primary ornamentation is more prominent in the central portion of the lateral sides and over the last-formed chambers of the test; it consists of short ridges in the continuation of the CBE that have the distinct tendency to anastomose on the lateral sides along the test growth axis. Pores are simple, with a circular and more rarely irregular outline, and a measured diameter or maximum dimension respectively of 0.0003-0.0008 mm.

Remarks. *Swiecickina paleocenica* differs from *S. senonicus* mainly by having more prominent primary ornamentation and better defined ridges over the CBE; it differs from *S. clavatus* mainly by having more prominent ornamentation in which some ridges present a preferential ornamentation that follows the elongation direction of the CBE, and a more prominent dentate periphery in the adult portion of the test.

Stratigraphical range: Maastrichtian-Paleocene.

Geographical distribution. EU (Belgium, Germany, Italy, Poland, Sweden, The Netherlands), Africa (Egypt, Libya), Asia (India, Japan), Mexico, Middle East (Israel), UK (England), North Atlantic Ocean, and South Atlantic Ocean.

STRATIGRAPHICAL DISTRIBUTION AND BIOSTRATIGRAPHY

The development of the new classification framework for the Santonian-Maastrichtian representatives of the family Bolivinoididae led to a critical review of the stratigraphical ranges of all the valid species (Fig. 50). This new classification presents a major advantage when compared to the previously developed ones: it includes data on test wall ultrastructure and high-detail test morphological features, which were only occasionally used in the past. Taking in consideration this advance in understanding the taxonomy and classification of the bolivinoidid group, and despite some incertitude that still persists it appears possible to make another step forward, namely towards refining the biostratigraphical framework based on the representatives of this group.

A historical perspective on methodology

The classical perspective on the use in biostratigraphy of the species of *Bolivinoides* was developed in the fifties and persisted over the next decades. According to it the stratigraphical ranges of the *Bolivinoides* species were plotted on the succession of stages and used in the definition of assemblage zones (Hiltermann, 1952, 1956; Hiltermann and Koch, 1955, 1956, 1960, 1962; Beckmann and Koch, 1964; Koch, 1977; Schönfeld, 1990, etc.). The potential and actual importance of some species of this genus in biostratigraphy was stressed occasionally, but no definition of biozones was provided within this stream of studies.

One significant step forward happened when Van Hinte (1967) provided a biostratigraphical framework based on five species of *Bolivinoides* for the Santonian-Maastrichtian stratigraphical interval. In stratigraphical order the five species are *B. strigillatus*, *B. hiltermanni*, *B. decoratus*, *B. miliaris* and *B. draco*. This author noted the incipient stage of development of a *Bolivinoides*-based biostratigraphy and showed that new data are necessary in order to validate and further improve the resolution of this biostratigraphical scheme. In a subsequent contribution, Van Hinte (1976, p. 277) considered all the five biozones as taxon range biozones. The pioneering work of Van Hinte (1967) was followed shortly by other authors. For example Ansary and Tewfik (1969) recognized the two Maastrichtian *Bolivinoides* biozones in Egypt, and the whole succession of five biozones by Barr (1970) in Libya and Petters (1976, 1977) in the New Jersey coastal plain subsurface. Biostratigraphical resolution increased gradually in these studies and as a result, other studies that focused on the Late Cretaceous biostratigraphy

Species stratigraphic ranges

Fig. 50. Stratigraphical ranges of the species of Bolivinoididae; B-Bolivinoides, T-Trachelinella, S-Swiecickina. Planktic foraminiferal zonation; grey background-heterohelicids, white background-coiled taxa; B-Bucherina, Ga-Globotruncana, Gu-Gublerina, Hx-Heterohelix, Pr-Praeglobotruncana, Rd-Radotruncana, S-Sigalia, V-Ventilabrella. Echinoderm/coleoid zonation; dark grey background-coileoid taxa, light grey background-echinoids, white background-crinoid taxa. B-Belemnella, Bl-Belemnella, G-Gonioteuthis, O-Offaster, M-Micraster, Mp-Marsupites, U-Uintacrinus.

started taking in consideration the *Bolivinoidea*-based framework (e.g., Robaszynski et al., 1984).

Biostratigraphical studies based on the species of *Bolivinoidea* diversified in the last decades. On one hand, there is a certain preoccupation in recognizing with precision the evolution and extinction of these taxa, especially in regions with a long tradition in these studies (e.g., Hart et al., 1989; Hampton et al., 2007; Bailey et al., 2009). Another trend is to refine the existing framework of Van Hinte (1967) by defining new biozones, based especially on local occurrences; this category is herein exemplified by the study of Jaff et al. (2014). Biozones based on local and regional occurrences have a low biostratigraphical resolution, which is excellently illustrated in this study by the succession of upper Campanian biozones (*Bolivinoidea decoratus* subzone at the base and *B. laevigatus* subzone above it) (Jaff et al., 2014, p. 165, 168-169), which is an anti-evolutionary succession (*B. laevigatus* is the ancestor of *B. decoratus* and not vice versa). A collapse in biostratigraphical resolution in the use of the representatives of the genus *Bolivinoidea* in Late Cretaceous biostratigraphy is recorded in the study of Dubicka in Walaszczyk et al. (2016); besides the low taxonomic resolution given by the misidentification of one index species (*B. culverensis*) and use of a vaguely defined species without revision (*B. vistulae*), most of the *Bolivinoidea*-based biozones are of partial taxon range type, with the species of this genus providing no bioevent corresponding to either lower or upper boundary (*B. decoratus*, *B. miliaris*, and *B. intermedius*).

This brief review allows us to observe that the evolutionary studies are paramount in conferring stability to a biostratigraphical framework, and explains why the successful and widely accepted biozonation of Van Hinte (1967) could have been produced only after the high-resolution evolutionary scheme of Barr (1966a) was published. It needs to be mentioned at this time that the classification in the present work and the stratigraphical ranges of the bolivinoidid species are given after the realization of such an evolutionary framework for the entire group (Fig. 51); it will be presented shortly as a distinct article, being given the length and complexity of this topic's presentation.

Methodology

A new biostratigraphical framework is developed herein and it is based on a new methodology, which appears necessary given the impediments encountered up to date in using bolivinoidid taxa in biostratigraphy. Experience and records in a large number of sections and regions show that in general that a small number of bolivinoidid species are recorded at one location. Moreover, these species are recorded in abundant assemblages in onshore sections rather than offshore drillings, where the samples are considerably smaller and therefore, sample bias is significant. In addition, the onshore sections encompass frequently shorter stratigraphical intervals and often appear difficult if not impossible to reconstruct with accuracy the stratigraphical ranges of certain species based on occurrences from one single region. Acknowledging these constraints given by the specifics of bolivinoidid fossil record is not enough, for there is another impediment given by the nature of studies realized in the past. According to the existing data there are regions in which many bolivinoidid

taxa occur and can be used in evolutionary reconstructions and for biostratigraphical purposes; classical examples can be considered England (Barr, 1966a), Libya (Barr, 1970) and New Jersey (Petters, 1967). But the experience shows the first and up till now only successful framework, which is that produced by Van Hinte (1967), was realized only by expanding the zone of provenance of the studied material. This pattern is used in the development of the new methodology, which is proposed herein.

At first it was realized a large-scale documentation of the occurrences of bolivinoidid taxa worldwide. The number of works (articles and books) in which bolivinoidid species were reported and are herein included in the synonymy lists and references, is more than 4.5 times higher than that the most documented article existing in the literature up to date. The wealth of data provided by these works were fully considered and integrated with the new data on the test morphology including ultrastructures produced in the course of the present study; all these data were further used to define the stratigraphical range of each of the species considered valid. By this it was sought to define the stratigraphical ranges of the bolivinoidid species at the scale of their entire areal of distribution. The new biostratigraphical framework is based on these ranges.

Biostratigraphical framework

A biostratigraphical framework consisting of eleven biozones is developed for the upper middle Santonian-Maastrichtian stratigraphical interval (Fig. 52). All the biozones are of interval biozone type and the bioevents that mark their boundaries are first occurrences (FO) of bolivinoidid taxa; the only exception to this rule is the upper boundary of the Maastrichtian, which is given by the extinction of the planktic foraminiferal species *Ventilabrella hariaensis* (Nederbragt, 1991). The eleven biozones are defined in stratigraphical order.

Bolivinoidea antiquum Biozone

- Age. Late middle Santonian-earliest late Santonian.
- Definition. Stratigraphical interval between the FO of the index species and the FO of *Bolivinoidea strigillata*.

Bolivinoidea strigillata Biozone

- Age. Late Santonian.
- Definition. Stratigraphical interval between the FO of the index species and the FO of *Bolivinoidea culverensis*.

Bolivinoidea culverensis Biozone

- Age. Latest Santonian-early early Campanian.
- Definition. Stratigraphical interval between the FO of the index species and the FO of *Bolivinoidea pustulata*.

Bolivinoidea pustulata Biozone

- Age. Late early Campanian.
- Definition. Stratigraphical interval between the FO of the index species and the FO of *Bolivinoidea decoratus*.

Bolivinoidea decoratus Biozone

- Age. Latest early Campanian-middle Campanian.
- Definition. Stratigraphical interval between the FO of the index species and the FO of *Bolivinoidea delicatula*.

Fig. 51. Evolutionary relationships amongst the Upper Cretaceous bolivinoidid species. Keys for abbreviations and backgrounds as in the previous figure.

Fig. 52. Biostratigraphical framework based on the representatives of the family Bolivinoididae and illustrated index species. Grey-not zoned.

***Bolivinoides delicatula* Biozone**

- Age. Latest middle Campanian-late Campanian.
- Definition. Stratigraphical interval between the FO of the index species and the FO of *Bolivinoides miliaris*.

***Bolivinoides miliaris* Biozone**

- Age. Late Campanian.
- Definition. Stratigraphical interval between the FO of the index species and the FO of *Bolivinoides australis*.

***Bolivinoides australis* Biozone**

- Age. Latest Campanian-earliest Maastrichtian.
- Definition. Stratigraphical interval between the FO of the index species and the FO of *Bolivinoides draco*.

***Bolivinoides draco* Biozone**

- Age. Early Maastrichtian.
- Definition. Stratigraphical interval between the FO of the index species and the FO of *Swiecickina paleocenica*.

***Swiecickina paleocenica* Biozone**

- Age. Late early Maastrichtian-early late Maastrichtian.
- Definition. Stratigraphical interval between the FO of the index species and the FO of *Bolivinoides vistulae*.

***Bolivinoides vistulae* Biozone**

- Age. Late Maastrichtian.
- Definition. Stratigraphical interval between the FO of the index species to the Cretaceous/Paleogene boundary, which is marked by the extinction of *Ventilabrella hariaensis* (Nederbragt, 1991).
- Remarks. The homonym biozone defined by Dubicka in Walaszczyk et al. (2016, p. 328) is based on an imprecise description of the index species that impedes in the identification of the biozone lower boundary. The taxonomic status of the species *B. vistulae* is stabilized through the emendation proposed in this work, but the validity of the biozone described before a precise and usable definition of the index species Dubicka and Peryt (2016) cannot be accepted.

CONCLUSIONS

A revision of the benthic foraminiferal family Bolivinoididae is proposed to accommodate the new data on test ultrastructure, ornamentation and porosity and re-evaluated features pertaining of the gross test architecture. This family is revised to include calcareous benthic foraminifera with the chambers of the last pairs with backward extensions and different morphologies of the aperture and periapertural structures. Three genera are included within family Bolivinoididae: *Bolivinoides* Cushman, 1927, *Trachelinella* Montanaro Gallitelli, 1956 and *Swiecickina* - new genus. A total of 24 species are included within these three genera.

Bolivinoides evolved in the middle Santonian and became extinct probably in the upper part of the Paleocene; occurrences of certain species of this genus within rocks of early Eocene age appear questionable and require further documentation. *Bolivinoides antiquum* - n. sp. is the oldest species of this genus and has the recorded evolutionary occurrence in the middle Santonian. The re-evaluation of the morphological features

pertaining of both test gross architecture and ultrastructural features leads to a highly accurate definition of the species. A terminology for the chamber backward extensions is proposed for the first time: apertural CBE, median CBE and peripheral CBE. Ornamentation structures are separated into primary and secondary ornamentation. Primary ornamentation consists of structures developed directly from the test wall. It can be absent resulting in a smooth chamber surface or consist of pustules, circular or elongate pore mounds and fine costae; notably, costate ornamentation is for the first time reported in the representatives of the genus *Bolivinoides* (e.g., *B. delicatula*). Secondary ornamentation consists of structures resulted from modifications of the chamber backward extensions and include peripheral keel (e.g., *B. draco*), ridges parallel to the growth axis (e.g., *B. draco*, *B. doreeni*) and peripheral rim (*B. peterssoni*). Dentate periphery is reported for the first time, in incipient stage at *B. peterssoni* and well-developed in the case of *B. vistulae*, which is a paramount feature in the revision of the latter.

Trachelinella of the Campanian-Maastrichtian stratigraphical interval is transferred to Bolivinoididae from the family Loxostomidae Loeblich and Tappan, 1962 based on the occurrence of chamber backward extensions. The ornamentation consisting of pore mounds that can fuse and form wide ornamentation structures is paramount in the definition of this genus as revised by Georgescu et al. (2011). The terminal aperture produced at the end of a short neck is another feature that differentiates *Trachelinella* from *Bolivinoides* where the aperture occurs on or at the base of the apertural face. *Trachelinella* is a monospecific genus consists of *T. watersi*.

Swiecickina is a new genus that includes three taxa that were originally described and subsequently assigned to *Bolivinoides*. The tests of the three species included within *Swiecickina* have a dentate periphery and are laterally compressed resulting in parallel or nearly parallel lateral sides an edge view and ornamentation consisting of ridges; the ridges can be irregularly distributed over the laterals sides, anastomosed in the central portion of the test lateral sides, and developed over the chamber backward extensions and conferring them a keeled appearance. *Swiecickina* is recorded from the uppermost Campanian-Paleocene stratigraphical interval.

A biostratigraphical framework based on species of the family Bolivinoididae is proposed for the middle Santonian-Maastrichtian stratigraphical interval. It consists of eleven biozones, ten having index species from the genus *Bolivinoides* and one from *Swiecickina*. The resolution of this new framework shows a considerable increase in resolution when compared to that previously developed by Van Hinte (1967) and Barr (1970) that consists of five biozones for the same stratigraphical interval. This marks a decrease in the average time interval spanned by one biozone from ~3.9 million years to slightly over 1.7 million years.

Acknowledgments. I would like to thank all the colleagues that helped me during this study. I thank especially Dr. D. Lazarus (Museum of Natural History, Berlin) for granting the permission to study the C.G. Ehrenberg Collection, Dr. C.M. Henderson (University of Calgary) who made available for study the McGugan Collection (University of Calgary)

and Dr. R.K. Olsson (Rutgers, the State University of New Jersey) who allowed the study of the S.W. Petters Collection during 1998-1999 when I was Senior Fulbright Fellow at this university. The headquarters of the Deep Sea Drilling Project/Ocean Drilling Program are thanked for answering positively to all of my many sample requests; this material forms the backbone of the present study. Special thanks are for Dr. J. Zhao (University of Calgary Libraries) for providing most of the references used in this article; without her help such level of documentation could not have been achieved. My gratitude is directed towards Schweizerbart Science Publishers, Elsevier Publishing Company and Cushman Foundation for granting the permission to re-illustrate some previously published specimens. The manuscript was reviewed and improved by two anonymous reviewers. Last but not least, Dr. B.P. Onac, editor of *Studia Universitatis Babeş-Bolyai, Geologia* is thanked for accepting the manuscript for publication.

REFERENCES

- Abdelghany, O., 2003, Late Campanian-Maastrichtian foraminifera from the Simsim Formation of the western side of the Northern Oman Mountains. *Cretaceous Research*, 24 (4): 391-405.
[https://doi.org/10.1016/S0195-6671\(03\)00051-X](https://doi.org/10.1016/S0195-6671(03)00051-X)
- Akimetz, V.S., 1961. Stratigraphy and foraminifera of the Upper Cretaceous deposits of Belarus. *Paleontologiya i Stratigrafiya BSSR*, 3: 3-245 (In Russian).
- Al-Shaibani, S.K., 1971, An Upper Maastrichtian foraminiferal fauna from Dörfles, Lower Austria. *Geologisches Jahrbuch*, 17: 105-119.
- Alegret, L., Ortiz, S. & Molina, E., 2009, Extinction and recovery of benthic foraminifera across the Paleocene-Eocene thermal maximum at the Alamedilla section (southern Spain). *Palaeogeography, Palaeoclimatology, Palaeoecology*, 279: 186-200.
<https://doi.org/10.1016/j.palaeo.2009.05.009>
- Alegret, L., Thomas, E., 2001, Upper Cretaceous and lower Paleogene benthic foraminifera from northeastern Mexico. *Micropaleontology*, 47: 269-316.
<https://doi.org/10.2113/47.4.269>
- Alegret, L., Thomas, E., 2007, Deep-Sea environments across the Cretaceous/Paleogene boundary in the eastern South Atlantic Ocean (ODP Leg 208, Walvis Ridge). *Marine Micropaleontology*, 64: 1-17.
<https://doi.org/10.1016/j.marmicro.2006.12.003>
- Alegret Badiola, L., 2008, Evolución paleoambiental durante el tránsito Cretácico/Paleógeno basada en foraminíferos bentónicos. *Revista Española de Paleontología*, 23: 15-30.
<https://recyt.fecyt.es/index.php/REP/article/view/8149>
- Aliiev, M.M., Aliyulla, H. & Halilov, A.G., 1988, *Cretaceous Fauna of Azerbaijan*. Akademya Nauk Azerbajanskoy SSR, Institut Geologii im. I.M. Gubkina. Elm, Baku, 447 p. (In Russian).
- Almogi-Labin, A., Flexer, A., Honigstein, A., Rosenfeld, A. & Rosenthal, E., 1990, Biostratigraphy and tectonically controlled sedimentation of the Maastrichtian in Israel and adjacent countries. *Revista Española de Paleontología*, 5: 41-52.
- Anan, H.S., 2011a, Additional to the Maastrichtian foraminifera of the Middle East. *Revue de Paléobiologie*, 30: 295-311.
- Anan, H.S., 2011b, Paleontology and stratigraphic significance of the Maastrichtian-Paleocene genus *Bolivinoidea* in some tethyan regions. *Egyptian Journal of Paleontology*, 11: 133-144.
- Anan, H.S., 2017, Evolutionary lineage of the Maastrichtian *Bolivinoidea draco* group (benthic foraminifera) in Abu Zenima section, west central Sinai, Egypt. *Arabian Journal of Geosciences*, 10 (19): 431.
<https://doi.org/10.1007/s12517-017-3219-2>
- Anan, H.S., Hamdan, A.R.A., 1992, *Bolivinoidea curtus* Reiss from the Paleocene of Jabal Malaqet, East of Al Ain, West of the northern Oman Mountains, United Arab Emirates. *Journal of the Faculty of Science of the U.A.E. University*, 4: 200-211.
- Ansary, S.E., Andrawis, S.F. & Fahmy, S.E., 1962, Biostratigraphic studies of the Upper Cretaceous sections in the G.P.C. wells in the Eastern Desert and Sinai. *Fourth Arab Petroleum Congress, Beirut*, paper 24 (B-3), 339-353.
- Ansary, S.E., Tewfik, N.M., 1969, Biostratigraphy and time stratigraphy of subsurface Upper Cretaceous of Ezz El Orban area, Gulf of Suez region, U.A.R. In *Proceedings of the Third African Micropaleontological Colloquium, Cairo* (Saird, R., Ed.). Cairo, p. 95-106.
- Applin, P.R., Applin, E.R., 1944, Regional subsurface stratigraphy and structure of Florida and southern Georgia. *Bulletin of the American Association of Petroleum Geologists*, 28: 1673-1753.
<http://archives.datapages.com/data/bulletns/1944-48/data/pg/0028/0012/1650/1673.htm>
- Ayyad, S.N., Abed, M.N. & Abu Zied, R.H., 1997, Biostratigraphy of the Upper Cretaceous rocks in Gebel Arif El-Naga, northeastern Sinai, Egypt, based on benthonic foraminifera. *Cretaceous Research*, 18: 141-159.
<https://doi.org/10.1006/cres.1996.0060>
- Bailey, H.W., Hart, M.B. & Swiecicki, A., 2009, Evolutionary lineages of benthic foraminifera in the chalk seas of N.W. Europe and their application to problem solving. In *Geologic Problem Solving with Microfossils: A Volume in Honor of Garry D. Jones* (Demchuk, T.D., Gary, A.C., Eds.). *Society of Economic Paleontologists and Mineralogists (Society of Sedimentary Geology), Special Publication*, 93: 233-249.
- Baranovskaya, O.F., Bulinnikova, S.P., 1964, Genus *Bolivinoidea*. In *Foraminifera of the Cretaceous and Paleogene deposits of the West-Siberian Lowlands* (Subbotina, N.N., Ed.). *Trudy VNIGRI*, 234: 1-454 (In Russian).
- Barr, F.T., 1966a, The foraminiferal genus *Bolivinoidea* from the Upper Cretaceous of the British Isles. *Palaeontology*, 9: 220-243.
https://www.palass.org/publications/palaeontology-journal/archive/9/2/article_pp220-243
- Barr, F.T., 1966b, Upper Cretaceous foraminifera from the Ballydeenlea Chalk, County Kerry, Ireland. *Palaeontology*, 9: 492-510.
https://www.palass.org/publications/palaeontology-journal/archive/9/3/article_pp492-510

- Barr, F.T., 1967, *Bolivinoides culverensis*, new name for the Campanian foraminifer *B. hiltermanni* Barr. *Contributions from the Cushman Foundation for Foraminiferal Research*, 18: 136.
- Barr, F.T., 1970, The foraminiferal genus *Bolivinoides* from the Upper Cretaceous of Libya. *Journal of Paleontology*, 44: 642-654.
<http://www.jstor.org/stable/1302658>
- Barr, F.T., Cordey, W.G., 1964. Some Upper Cretaceous foraminifera from the Chapman Collection. *Journal of Paleontology*, 38: 306-310.
<http://www.jstor.org/stable/1301550>
- Bate, R.H., Bayliss, D.D., 1969, An outline account of the Cretaceous and Tertiary foraminifera and of the Cretaceous ostracods of Tanzania. *Proceedings of the Third African Micropaleontology Colloquium*, Cairo, p. 113-164.
- Belford, D.J., 1960, Upper Cretaceous foraminifera from the Toolonga Calcilitite and Gingin Chalk, Western Australia. *Bulletin of the Bureau of Mineral Resources, Geology and Geophysics*, 57: 1-118.
- Beckmann, J.P., 1960, Distribution of benthonic foraminifera at the Cretaceous-Tertiary boundary of Trinidad (West Indies). *Reports of the 21st International Geological Congress, Norden 1960*, 5: 57-69.
- Beckmann, J.P., 1978, Late Cretaceous smaller benthic foraminifera from Sites 363 and 364 DSDP Leg 40, Southeast Atlantic Ocean. In *Initial Reports of the Deep Sea Drilling Project* (Bolli, H.M., Ryan, W.B.F., et al., Eds.). United States Government Printing Office, Washington, D.C., 40: 759-781.
<https://doi.org/10.2973/dsdp.proc.40.118.1978>
- Beckmann, J.P., Koch, W., 1964, Vergleiche von *Bolivinoides*, *Aragonia* und *Tappanina* (Foraminifera) aus Trinidad (Westindien) und Mitteleuropa. *Geologisches Jahrbuch*, 83: 31-64.
- Beniamovsky, V.N., Alekseev, A.S., Ovechkina, M.N., Vishnevskaya, V.S., Podgaetskii, A.V. & Pronin, V.G., 2012, Upper Campanian-lower Maastrichtian sections of the northwestern Rostov region. Article 1. Description, paleontological assemblages, and lithostratigraphy. *Stratigraphy and Geological Correlation*, 20: 346-379.
<https://doi.org/10.1134/S0869593812040028>
- Beniamovsky, V.N., Kopaevich, L.F., 2016, The Alan-Kyr Coniacian-Campanian section (Crimean Mountains): biostratigraphy and paleobiogeography aspects. *Moscow University Geology Bulletin*, 71 (3): 217-233.
<https://doi.org/10.3103/S0145875216030042>
- Bettenstaedt, F., Wicher, C.A., 1955, Stratigraphic correlation of Upper Cretaceous and Lower Cretaceous in the Tethys and boreal by the aid of microfossils. *Proceedings of the Fourth World Petroleum Congress, Geology and Geophysics*, I/D: 493-545.
- Bieda, E., 1958, Otwornice przewodnie I wiek kredy piszącej Mielnika. *Biuletyn Instytutu Geologicznego*, 121: 17-89. (In Polish).
- Bless, M.J.M., Felder, P.J. & Jagt, J.W.M., 1990, Repeated Tethyan influences in the early Campanian to middle late Maastrichtian successions of Folx-les-Caves and Orp-le-Petit (eastern Brabant Massif, Belgium). *Annales de la Société Géologique de Belgique*, 113: 179-197. (Published in 1991).
<http://popups.uliege.be/0037-9395/index.php?id=1341>
- Bolli, H.M., Beckmann, J.-P. & Saunders, J.B., 1994, *Benthic foraminiferal biostratigraphy of the south Caribbean region*. Cambridge University Press, Cambridge, 408 p.
<https://doi.org/10.1017/CBO9780511564406>
- Boltovskoy, E., Watanabe, S., Totah, V. & Vera Ocampo, J., 1995, Benthic foraminifera from DSDP Site 516 (upper Maastrichtian-Quaternary, South Atlantic). *Revista Española de Micropaleontología*, 27: 111-140.
- Brotzen, F., 1945, De geologiska resultaten från borrhningarna vid Höllviken. Preliminär rapport. Del I: Kritan. *Sveriges Geologiska Undersökning*, 465 (38): 1-64.
- Brotzen, F., 1948, The Swedish Paleocene and its foraminiferal fauna. *Sveriges Geologiska Undersökning*, 493 (42): 1-140.
- Bukowy, S., Geroch, S., 1956, On the age of exotic conglomerates at Kruhel Wielki near Przemyśl (Carpathians). *Annales de la Société Géologique de Pologne*, 26: 297-329. (In Polish) (Published in 1957).
- Busulini, A., Dieni, I., Massari, F., Pejović & Wiedmann, J., 1984. Nouvelles données sur le Crétacé Supérieur de la Sardigne Orientale. *Cretaceous Research*, 5: 243-258.
[https://doi.org/10.1016/S0195-6671\(84\)80021-X](https://doi.org/10.1016/S0195-6671(84)80021-X)
- Butt, A.A., 1986, Upper Cretaceous biostratigraphy in the Sirte Basin, northern Libya. *Revue de Paléobiologie*, 5: 175-191.
- Bykova, N.K., 1959, Questions towards the laws of phylogenetic development of foraminifera in the conditions of periodical environmental changes. *Voprosy Paleontologii i Biostratigrafii, Trudy Vsesoyuznogo Paleontologicheskogo Obshchestva*, 2:63-75. (In Russian).
- Campbell, R.J., Howe, R.W. & Rexilius, J.P., 2004, Middle Campanian-lowermost Maastrichtian nannofossil and foraminiferal biostratigraphy of the northwestern Australian margin. *Cretaceous Research*, 25: 827-864.
<https://doi.org/10.1016/j.cretres.2004.08.003>
- Carsey, D.O., 1926, Foraminifera of the Cretaceous of central Texas. *Texas University Bulletin*, 2612: 3-56.
<https://repositories.lib.utexas.edu/handle/2152/23442>
- Cati, F., 1964, Una microfauna campaniana dei Monti Berici (Vicenza). *Giornale di Geologia*, 32: 199-271.
- Chapman, F., 1892, Microzoa from the Phosphatic Chalk of Taplow. *Quarterly Journal of the Geological Society of London*, 48: 514-518.
<https://doi.org/10.1144/GSL.JGS.1892.048.01-04.38>
- Cole, W.S., 1938, Stratigraphy and micropaleontology of two deep wells in Florida. *Bulletin of the Florida State Geological Survey*, 16: 1-48.
- Cushman, J.A., 1926a, Some foraminifera from the Mendez Shale of eastern Mexico. *Contributions from the Cushman Laboratory for Foraminiferal Research*, 2: 16-28.
- Cushman, J.A., 1926b, The foraminifera of the Velasco Shale of the Tampico Embayment. *American Association of Petroleum Geologists Bulletin*, 10: 581-612.
- Cushman, J.A., 1927a, American Upper Cretaceous species of *Bolivina* and related species. *Contributions from the Cushman Laboratory for Foraminiferal Research*, 2: 85-91.

- Cushman, J.A., 1927b. An outline of a re-classification of the foraminifera. *Contributions from the Cushman Laboratory for Foraminiferal Research*, 3: 1-105.
- Cushman, J.A., 1927c. Some characteristic Mexican fossil foraminifera. *Journal of Paleontology*, 1: 147-172.
- Cushman, J.A., 1931. The foraminifera of the Saratoga Chalk. *Journal of Paleontology*, 5: 297-315.
- Cushman, J.A., 1932. The foraminifera of the Annona Chalk. *Journal of Paleontology*, 6: 330-345.
- Cushman, J.A., 1933. An illustrated key to the genera of the Foraminifera. *Cushman Laboratory for Foraminiferal Research, Special Publication*, 5: 1-40.
- Cushman, J.A., 1937. A few new species of American Cretaceous foraminifera. *Contributions from the Cushman Laboratory for Foraminiferal Research*, 13: 100-105.
- Cushman, J.A., 1944. The foraminiferal fauna of the type locality of the Pecan Gap Chalk. *Contributions from the Cushman Laboratory for Foraminiferal Research*, 20: 1-16.
- Cushman, J.A., 1946. Upper Cretaceous foraminifera of the Gulf Coastal region of the United States and adjacent areas. *United States Geological Survey Professional Paper*, 206: 1-160.
- Cushman, J.A., 1948. Foraminifera from the Hammond Well. In: *The Stratigraphy, Paleontology, and Sedimentology of Three Deep test Wells on the Eastern Shore of Maryland*. Baltimore: Department of Geology, Mines, and Water Resources, 2: 213-267.
- Cushman, J.A., 1950. *Foraminifera. Their Classification and Economic Use*, (4th ed.). Harvard University Press, Cambridge, Massachusetts, 605 p.
- Cushman, J.A., Jarvis, P.W., 1928. Cretaceous foraminifera from Trinidad. *Contributions from the Cushman Laboratory for Foraminiferal Research*, 4: 85-103.
- Cushman, J.A. and Jarvis, P.W., 1932. Upper Cretaceous foraminifera from Trinidad. *Proceedings of the United States National Museum*, 80: 1-60.
- Cushman, J.A., Renz, H.H., 1946. The foraminiferal fauna of the Lizard Springs Formation of Trinidad, British West Islands. *Cushman Laboratory for Foraminiferal Research, Special Publication*, 18: 1-48.
- Dailey, D.H., 1983. Late Cretaceous and Paleocene benthic foraminifera from Deep Sea Drilling Project Site 516, Rio Grande Rise, western South Atlantic Ocean. In *Initial Reports of the Deep Sea Drilling Project* (Barker, P.F., Johnson, D.A. et al., Eds.). United States Government Printing Office, Washington, D.C., 72: 757-782. <https://doi.org/10.2973/dsdp.proc.72.134.1983>
- Dain, L.G., 1934. Foraminifera of the Upper Jurassic and Cretaceous deposits of the Djasky-Bai of the Temir region. *Trudy Neftyanogo Geologo-Razvednochnogo Instituta*, 43: 1-62 (In Russian).
- Dain, L.G., 1961. Some species of foraminifera from the Upper Cretaceous of the Shumihinsk District of the Chelyabinsk region. *Trudy VNIGRI (170), Microfauna SSSR*, 12: 4-37 (In Russian).
- Dampel, N., 1934. Foraminifera from the Upper Cretaceous of the Karaton Field (Emba Region). *Trudy Neftyanogo Geologo-Razvednochnogo Instituta*, 50: 3-33 (In Russian).
- Dubicka, Z., 2015. Benthic foraminiferal biostratigraphy of the lower and middle Campanian of the Polish Lowlands and its application for interregional correlation. *Cretaceous Research*, 56: 491-503. <https://doi.org/10.1016/j.cretres.2015.06.012>
- Dubicka, Z., Jurkowska, A., Thibault, N., Razmjooei, M.J., Wójcik, K., Gorzelak, P. & Felisiak, I., 2017. An integrated stratigraphic study across the Santonian-Campanian boundary at Bocieniec, southern Poland: a new boundary stratotype candidate. *Cretaceous Research*, 80: 61-85. <https://doi.org/10.1016/j.cretres.2017.07.012>
- Dubicka, Z., Peryt, D., 2011. Integrated biostratigraphy of upper Maastrichtian chalk at Chelm (SE Poland). *Annales Societatis Geologorum Poloniae*, 81: 185-197. http://www.asgp.pl/81_2_185_197
- Dubicka, Z., Peryt, D., 2016. *Bolivinoidea* (benthic foraminifera) from the Upper Cretaceous of Poland and western Ukraine: taxonomy, evolutionary changes and stratigraphic significance. *Journal of Foraminiferal Research*, 46: 75-94. <https://doi.org/10.2113/gsjfr.46.1.75>
- Edgell, H.S., 1954. The stratigraphical value of *Bolivinoidea* in the Upper Cretaceous of northwest Australia. *Contributions from the Cushman Foundation for Foraminiferal Research*, 5: 68-76.
- Egger, J.G., 1899. Foraminiferen und Ostrakoden aus den Kreidemergeln der Oberbayerischen Alpen. *Abhandlungen der Mathematisch-physikalischen Classe der Königlich Bayerischen Akademie der Wissenschaften*, 21: 3-230. (published in 1902).
- El Ashwah, A.A.E., 2005. Stratigraphical studies on the Late Cretaceous rocks in Shushan well no. 1, north Western Desert, Egypt. *Egyptian Journal of Paleontology*, 5: 107-118.
- El-Dawy, M.H., 2001. Paleocene benthic foraminiferal biostratigraphy and paleobathymetry in the sections between El Sheikh Fadl and Ras Gharib, Eastern Desert, Egypt. *Micropaleontology*, 47: 23-46. <https://doi.org/10.2113/47.1.23>
- El-Nady, H., 2006. Contribution to the stratigraphic significance of the genus *Bolivinoidea* and their paleoecology across the Campanian/Maastrichtian boundary in the Gabal El-Mouriefik, eastern Sinai, Egypt. *Revue de Paléobiologie*, 25: 671-692.
- Eternod Olvera, Y., 1959. Foraminiferos del Cretacico superior de la Quenca de Tampico-Tuxpan, Mexico. *Boletin de la Asociación Mexicana de Geólogos Petroleros*, 11: 63-134.
- Felder, P.J.S., Bless, M.J.M., 1993. The Vijlen Chalk (early early to early late Maastrichtian) in its type area around Vijlen and Mameis (southern Limburg, The Netherlands). *Annales de la Société Géologique de Belgique*, 116: 61-85.
- Finlay, H.J., 1939. New Zealand Foraminifera: key species in stratigraphy - No. 3. *Transactions and Proceedings of the Royal Society of New Zealand*, 69: 309-329.
- Finlay, H.J., 1940. New Zealand Foraminifera: key species in stratigraphy - No. 4. *Transactions and Proceedings of the Royal Society of New Zealand*, 69: 448-472.
- Franke, A., 1928. Die Foraminiferen der Oberen Kreide Nord- und Mitteldeutschlands. *Abhandlungen der Preußischen Geologischen Landesanstalt*, 111: 1-207.

- Frenzel, P., 2000. Die benthischen Foraminiferen der Rügener Schreiekreide (Unter-Maastricht, NE-Deutschland). *Neue Paläontologische Abhandlungen*, 3: 1-361.
- Frizzell, D.L., 1954, Handbook of Cretaceous foraminifera of Texas. *Bureau of Economic Geology, University of Texas, Report of Investigations*, 22: 1-232.
- Galloway, J.J., 1933, *A manual of foraminifera*. The Principia Press, Bloomington, 483 p.
- Gasiński, M.A., 1993, Foraminiferids of the San Nicola section, Cilento Flysch (southern Apennines). *Geologica Romana*, 29: 181-185.
- Gawor-Biedowa, E., 1984, Foraminiferal zonation of the Upper Cretaceous deposits in Poland (except for the Carpathians and Sudeten). In *Benthos '83, 2nd Symposium on Benthic Foraminifera* (Oertli, H.J., Ed.), Pau et Bordeaux, p. 213-223.
- Gawor-Biedowa, E., 1992, Campanian and Maastrichtian foraminifera from the Lublin Upland, Eastern Poland. *Palaeontologia Polonica*, 52: 3-187.
<http://www.palaeontologia.pan.pl/>
- Georgescu, M.D., 2006, Santonian-Campanian planktonic foraminifera in the New Jersey coastal plain and their distribution related to the relative sea-level cycles. *Canadian Journal of Earth Sciences*, 43: 101-120.
<https://doi.org/10.1139/e05-090>
- Georgescu, M.D., 2009, On the origins of superfamily Heterohelicoidea Cushman, 1927 and the polyphyletic nature of planktic foraminifera. *Revista Española de Micropaleontología*, 41: 107-144.
- Georgescu, M.D., 2010a, Origin, taxonomic revision and evolutionary classification of the late Coniacian-early Campanian (Late Cretaceous) planktic foraminifera with multichamber growth in the adult stage. *Revista Española de Micropaleontología*, 42: 59-118.
- Georgescu, M.D., 2010b, Evolutionary classification of the Upper Cretaceous (Turonian-lower Campanian) planktic foraminifera with incipient meridional ornamentation. *Journal of Micropaleontology*, 29: 149-161.
<https://doi.org/10.1144/0262-821X10-004>
- Georgescu, M.D., 2012, Restudy of the type specimens of *Phanerostomum asperum* Ehrenberg 1854 (Late Cretaceous, Foraminiferida). *Micropaleontology*, 58: 291-303.
<http://www.jstor.org/stable/23250635>
- Georgescu, M.D., 2013, Revised evolutionary systematics of the Cretaceous planktic foraminifera described by C.G. Ehrenberg. *Micropaleontology*, 59: 1-49.
<http://www.jstor.org/stable/24413314>
- Georgescu, M.D., 2014a, Evolution of central perforate plate in the new Condensed Lineage *Eicheriella*. In *Evolutionary Classification and English-based Nomenclature in Cretaceous Planktic Foraminifera* (Georgescu, M.D., Henderson, C.M., Eds.). Nova Science Publishers, New York, p. 93-100.
- Georgescu, M.D., 2014b, New Late Cretaceous (Santonian-Maastrichtian) heterohelicoid planktic foraminifera from the Pacific and Indian Oceans and their biostratigraphic and evolutionary significance. In *Evolutionary Classification and English-based Nomenclature in Cretaceous Planktic Foraminifera* (Georgescu, M.D., Henderson, C.M., Eds.). Nova Science Publishers, New York, p. 39-71.
- Georgescu, M.D., 2017, Upper Cretaceous planktic foraminiferal biostratigraphy. *UBB Geologia*, 61: 5-20.
<https://doi.org/10.5038/1937-8602.61.1.1297>
- Georgescu, M.D., Arz, J.A., Macauley, R.V., Kukulski, R.B., Arenillas, I. & Pérez-Rodríguez, I., 2011, Late cretaceous (Santonian-Maastrichtian) serial foraminifera with pore mounds or pore mound-based ornamentation structures. *Revista Española de Micropaleontología*, 43: 109-139.
- Glazumova, A.E., Balakhmatova, V.T., Lipman, R.H., Romanova, V.I. & Hohlova, I.A., 1960, Stratigraphy and fauna of the Cretaceous deposits of Siberia. *Trudy VSEGEI*, 29: 1-347.
- Goel, R.K., 1965. Contribution à l'étude des foraminifères du Crétacé supérieur de la Basse-Seine. *Bulletin du Bureau de Recherches Géologiques et Minières*, 1965/5: 49-157.
- Gorbenko, V.F., 1974, Phylum Protozoa, class Sarcodina, subclass Foraminifera. In *Atlas of the Cretaceous fauna from Donbass*. Nedra, Moscow, 26-61. (In Russian).
- Govindan, A., 1977, Upper Cretaceous and Lower tertiary foraminiferal genus Bolivinoidea from the Cauvery Basin, South India and its palaeobiogeographical significance. *Journal of the Geological Society of India*, 18: 459-476.
- Graham, J.J., Church, C.C., 1963, Campanian foraminifera from the Stanford University Campus, California. *Stanford University Publications, Geological History*, 8: 1-106.
- Hagn, H., 1953, Die Foraminiferen der Pinswanger Schichten (Unteres Obercampan). *Palaeontographica*, 104: 1-119.
- Hampton, M.J., Bailey, H.W., Gallagher, L.T., Mortimore, R.N. & Wood, C.J., 2007, The biostratigraphy of Seaford Head, Sussex, southern England; an international reference section for the basal boundaries for the Santonian and Campanian Stages in chalk facies. *Cretaceous Research*, 28: 46-60. <https://doi.org/10.1016/j.cretres.2006.05.025>
- Hanzliková, E., 1969, The foraminifera of the Frýdek Formation (Senonian). *Sborník Geologických Věd, Paleontologie*, 11: 7-79.
- Hanzliková, E., 1972, Carpathian Upper Cretaceous Foraminiferida of Moravia (Turonian-Maastrichtian). *Rozprawy Ústředního Ústavu Geologického*, 39: 1-160.
- Hart, M.B., Bailey, H.W., Crittenden, S., Fletcher, B.N., Price, R.J. & Swiecicki, A., 1989. Cretaceous. In *Stratigraphical Atlas of Fossil Foraminifera*, (2nd ed.) (Jenkins, D.G., Murray, J.W., Eds.). Ellis Horwood Limited, Chichester, p. 273-371.
- Hay, W.H., DeConto, R.M., Wold, C.N., Wilson, K.M., Voigt, S., Schulz, M., Wold, A.R., Dullo, W., Ronov, A.B., Balukhovskiy, A.N. & Söding, E., 1999, Alternative global Cretaceous paleogeography. In *Evolution of the Cretaceous Ocean-Climate System* (Barrera, E., Johnson, C.C., Eds.). *The Geological Society of America Special Publication*, 332: 1-47. <https://doi.org/10.1130/0-8137-2332-9.1>
- Heller, I., 1975, Microbiostratigraphy of the Cretaceous deposits in the southern part of Łódź Synclinorium (Central Poland). *Rocznik Polskiego Towarzystwa Geologicznego*, 45: 233-254. http://www.asgp.pl/45_2
- Heron-Allen, E., Earland, A., 1910, On the Recent and fossil foraminifera of the shore-sands of Selsey Bill, Sussex. - V. Cretaceous foraminifera. *Journal of the Royal Microscopical Society*, 30 (4): 401-426.
<https://doi.org/10.1111/j.1365-2818.1910.tb01342.x>

- Hewaidy, A.E.G., El Ashwah, A., 1993, Faunal and environmental studies on the Upper Cretaceous sequence in El Qusaima area, North-East Sinai, Egypt. *Annals of the Geological Survey of Egypt*, 19: 261-288.
- Hillebrandt, A. von, 1962, Das Paleozän und seine Foraminiferenfauna im Becken von Reichenhall und Salzburg. *Bayerische Akademie der Wissenschaften, Mathematisch-Naturwissenschaftliche Klasse, Abhandlungen*, 108: 1-182.
- Hiltebert, H., 1952, Stratigraphische Fragen des Campan und Maastricht unter besonderer Berücksichtigung der Mikropaläontologie. *Geologisches Jahrbuch*, 64: 595-632.
- Hiltebert, H., 1956, Biostratigraphie der Oberkreide auf Grund von Mikrofossilien. *Paläontologische Zeitschrift*, 30: 19-32.
- Hiltebert, H., 1963, Zur Entwicklung der benthos-Foraminifere *Bolivinoidea*. In *Evolutionary Trends in Foraminifera* (von Koenigswald, G.H.R., Emeis, J.D., Buning, W.L., Wagner, C.L., Eds.). Elsevier Publishing Company, Amsterdam-London-New York, 198-223.
- Hiltebert, H., Koch, W., 1950, Taxonomie und Vertikalverbreitung von *Bolivinoidea*-Arten im Senon Nordwestdeutschlands. *Geologisches Jahrbuch*, 64: 595-632.
- Hiltebert, H., Koch, W., 1955, Biostratigraphie der Grenzsichten Maastricht/Campan in Lüneburg und in der Bohrung Brunhilde. 2 Teil: Foraminiferen. *Geologisches Jahrbuch*, 70: 357-384.
- Hiltebert, H., Koch, W., 1956, Mikropaläontologische Feinhorizontierung von Santon-Profilen durch das Erzlager Lengede-Broistedt. *Paläontologische Zeitschrift*, 30: 33-44.
- Hiltebert, H., Koch, W., 1960, Oberkreide Biostratigraphie Mittels Foraminiferen. *Reports of the 21st International Geological Congress*, Norden, 6: 69-76.
- Hiltebert, H., Koch, W., 1962, Oberkreide der nördlichen Mitteleuropa. In *Leitfossilien der Mikropaläontologie* (Bartenstein H., et al., Eds.). Gebrüder Borntraeger, Berlin-Nikolassee, 299-338.
- Hofker, J., 1952, Zur Fassung der Foraminiferengattung *Bolivinoidea* Cushman 1927. *Geologisches Jahrbuch*, 66: 377-382.
- Hofker, J., 1955, The genus *Bolivinoidea* in the Cretaceous of south Limburg. *Natuurhistorisch Maandblad*, 44: 68-71.
- Hofker, J., 1956a, Die *Pseudotextularia*-Zone der Bohrung Maasbult I und ihre Foraminiferen-Fauna. *Paläontologische Zeitschrift*, 30: 59-79.
- Hofker, J., 1956b, Les Foraminifères de la zone de contact Maastrichtien-Campanien dans l'est de la Belgique et le sud des Pays-Bas. *Annales de la Société Géologique de Belgique*, 80: 191-233.
- Hofker, J., 1957, Foraminiferen der Oberkreide von Nordwestdeutschland und Holland. *Beihefte zum Geologischen Jahrbuch*, 27: 1-464.
- Hofker, J., 1958a, Upper Cretaceous *Bolivinoidea* guide forms. *Micropaleontology*, 4: 329-333. <https://doi.org/10.2307/1484294>
- Hofker, J., 1958b, Foraminifera from the Cretaceous of Limburg, Netherlands. 38. The gliding change in *Bolivinoidea* during time. *Natuurhistorisch Maandblad*, 47: 145-159.
- Hofker, J., 1958c, Les Foraminifères du Crétacé supérieur de Glons. *Annales de la Société Géologique de Belgique*, 81: 467-493.
- Hofker, J., 1960a, The foraminifera of the lower boundary of the Danish Danian. *Meddelelser fra Dansk Geologisk Forening*, 14: 212-242.
- Hofker, J., 1960b, Les Foraminifères du Crétacé supérieur dans le Bassin de Mons. I. Les Foraminifères de la Craie Phosphatée de Ciply. *Annales de la Société Géologique de Belgique*, 83: 165-180.
- Hofker, J., 1960c, Les Foraminifères du Crétacé supérieur dans le Bassin de Mons. II. Les Foraminifères du tuffeau de Saint Symphorien. *Annales de la Société Géologique de Belgique*, 83: 181-195.
- Hofker, J., 1962, Correlation of the Tuff Chalk of Maastricht (Type Maastrichtian) with the Danske Kalk of Denmark (Type Danian), the stratigraphic position of the Type Montian, and the planktonic foraminiferal faunal break. *Journal of Paleontology*, 36: 1051-1089.
- Hofker, J., 1966, Maastrichtian, Danian and Paleocene foraminifera. The foraminifera of the type Maastrichtian in south Limburg, Netherlands, together with the foraminifera of the underlying Gulpen Chalk and the overlying calcareous sediments; the foraminifera of the Danske Kalk and the overlying greensands and clays as found in Denmark. *Paleontographica Supplement*, 10: 1-376.
- Hofmann, T., Švábenická, L. & Hradecká, L., 1999, Biostratigraphy of fissure fillings in the Ernstbrunn Limestone of the Waschberg Zone (Lower Austria). *Abhandlungen der Geologischen Bundesanstalt*, 56: 465-474.
- Holbourn, A., Henderson, A.S. & MacLeod, N., 2013, Atlas of Benthic Foraminifera. Wiley-Blackwell, Chichester, 642 p.
- Howe, R.W., Campbell, R.J. & Rexilius, J.P., 2003, Integrated uppermost Campanian-Maastrichtian calcareous nannofossil and foraminiferal biostratigraphic zonation of the northwestern margin of Australia. *Journal of Micropalaeontology*, 22: 29-62. <https://doi.org/10.1144/jm.22.1.29>
- Huber, B.T., 1992, Upper Cretaceous planktic foraminiferal biozonation for the Austral Realm. *Marine Micropaleontology*, 20: 107-128. [https://doi.org/10.1016/0377-8398\(92\)90002-2](https://doi.org/10.1016/0377-8398(92)90002-2)
- Huber, B.T., Hodell, D.A. & Hamilton, C.P., 1995, Middle-Late Cretaceous climate of the southern low latitudes: stable isotopic evidence for minimal equator-to-pole thermal gradients. *Geological Society of America Bulletin*, 107: 1164-1191. [https://doi.org/10.1130/0016-7606\(1995\)107%3C1164:MLCCOT%3E2.3.CO;2](https://doi.org/10.1130/0016-7606(1995)107%3C1164:MLCCOT%3E2.3.CO;2)
- Ismail, A.A., 2002, The biostratigraphic and sequence stratigraphic applications on the Upper Cretaceous succession of Wadi Feiran, southwestern Sinai, Egypt. *Egyptian Journal of Geology*, 46: 515-533.
- Ismail, A.A., 2012, Late Cretaceous-Early Eocene benthic foraminifera from Esh El Mallaha area, Egypt. *Revue de Paléobiologie*, 31: 15-50.
- Ismail, A.A., El Saadany, M.A., 1995, Some Cretaceous foraminifera from northern Sinai, Egypt. *Middle East Research Center, Ain Shams University, Earth Sciences Series*, 9: 173-203.

- Jaff, R.B.N., Williams, M., Wilkinson, I.P., Lawa, F., Lee, S. & Zalasiewicz, J., 2014, A refined foraminiferal biostratigraphy for the late Campanian-early Maastrichtian succession of northeast Iraq. *GeoArabia*, 19 (1): 161-180. <http://nora.nerc.ac.uk/id/eprint/507074/>
- Jordan, H., Gasse, W., 1986, Bio- und lithostratigraphisch Untersuchungen des glaukonitisch-sandigen Santon und Campan im südwestlichen Münsterland. *Geologie und Paläontologie in Westfalen*, 6: 7-49.
- Kalinin, N.A., 1937, Foraminifera from the Cretaceous of Baktygarny (Aktiubinsk Province). *Etyudy Mikropaleontologii*, 1: 7-61 (In Russian).
- Kaptarenko-Chernousova, O.K., Goliak, L.M., Zernetskyi, B.F., Kraeva, E.I. & Lipnik, E.S., 1963, Atlas of the Characteristic Foraminifera of the Jurassic, Cretaceous and Paleogene of the Platform Part of the Ukraine. *Institut Geologicheskikh Nauk, Seriya Stratigraphiya i Paleontologiya*, 45: 1-200 (In Russian).
- Kaptarenko-Chernousova, O.K., Plotnikova, L.F. & Lipnik, E.S., 1979, *Foraminifera of the Ukrainian Cretaceous*. Naukova Dumka, Kiev, 256 p (In Russian).
- Kavary, E., Frizzell, D.L., 1963, Upper Cretaceous and Lower Cenozoic foraminifera from west central Iran. *University of Missouri School of Mines and Metallurgy Bulletin*, 102: 1-89.
- Keller, B.M., 1939, Foraminifera of the Upper Cretaceous of USSR. *Neftianogo Geologo-Razvedochnogo Instituta*, 116: 7-21 (In Russian).
- Keller, G., 1988, Biotic turnover in benthic foraminifera across the Cretaceous/Tertiary boundary at El Kef, Tunisia. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 66: 153-171. [https://doi.org/10.1016/0031-0182\(88\)90198-8](https://doi.org/10.1016/0031-0182(88)90198-8)
- Khalil, H.T., 1998, Late Cretaceous benthonic foraminiferal biostratigraphy, Sinai, Egypt, with special consideration to the genus *Bolivinoidea*. *Neues Jahrbuch für Geologie und Paläontologie, Monatshefte*, 7: 415-431.
- Khalil, H., Mashaly, S., 2004, Stratigraphy and stage boundaries of the Upper Cretaceous-Lower Paleogene succession in Gabal Musaba Salama area, southwestern Sinai, Egypt. *Egyptian Journal of Paleontology*, 4: 1-37.
- Kiesel, Y., Trümper, E., 1965, Mikrofauna und Stratigraphie im Grenzbereich Kreide/Tertiär in der Bohrung Nennhausen 2. *Mitteilungen des Zentralen Geologischen Instituts*, 1: 155-169.
- Kikoïne, J., 1948, Les Heterohelidae du Crétacé supérieur pyrénéen. *Bulletin de la Société Géologique de France*, 18 (1): 15-35.
- Klasz, I. de, Klasz, S. de, 1993, Important calcareous foraminifera of the type locality of the genus *Quadrabulimina* DE KLASZ, region of Eisenärzt, Upper Bavaria. *Zitteliana*, 20: 311-321.
- Klasz, I. de, Klasz, S. de & Ausseil-Badie, J., 1988, Étude systématique des foraminifères du Danien de la Formation des Madeleines de Dakar (Sénégal). *Cahiers de Micropaléontologie*, 2: 29-112.
- Koch, W., 1977, Biostratigraphie in der Oberkreide und Taxonomie von Foraminiferen. *Geologisches Jahrbuch*, A38: 11-123.
- Koutsoukos, E.A.M. & Klasz, I. de, 2000, Late Cretaceous foraminiferal biogeography (Families bolivinoidea, Buliminellidae, Gavelinellidae, Siphogenerinoidea, turillinoidea) in northeastern Brazilian shelf and central West African basins. *Cretaceous Research*, 21: 381-405. <https://doi.org/10.1006/cres.1999.0206>
- Küpper, I., 1963, Mikropaläontologische Gliederung der Oberkreide des Beckenuntergrundes in den oberösterreichischen Molassebohrungen. *Mitteilungen der Geologischen Gesellschaft in Wien*, 56: 591-651 (Published in 1964).
- Liszka, S., 1955, Foraminifera of the Lower Senonian in the vicinity of Krakow. *Rocznik Polskiego Towarzystwa Geologicznego*, 23: 165-190 (In Polish). http://www.asgp.pl/45_2
- Liszkowa, J., Morgiel, J.J., 1985, Contribution to the knowledge of the foraminifera of the Frydek type facies in the Polish Outer Carpathians. *Kwartalnik Geologiczny*, 29: 65-84.
- Loeblich, A.R. Jr., Tappan, H., 1962, Six new generic names in the Mycetozoida (Trichiidae) and Foraminiferida (Fischerinidae, Buliminidae, Caucasinidae and Pleurostomellidae), and a redescription of *Loxostomum* (Loxostomidae, new family). *Proceedings of the Biological Society of Washington*, 75: 107-113.
- Loeblich, A.R. Jr., Tappan, H., 1964, Sarcodina Chiefly "Thecamoebians" and Foraminifera. In *Treatise on Invertebrate Paleontology. Part C* (Moore, R.C., Ed.). The Geological Society of America and The University of Kansas Press, Boulder and Lawrence, 1-900.
- Loeblich, A.R. Jr., Tappan, H., 1984, Suprageneric classification of the Foraminiferida (Protozoa). *Micropaleontology*, 30: 1-70. <https://doi.org/10.2307/1485456>
- Loeblich, A.R. Jr., Tappan, H., 1987, *Foraminiferal genera and their classification*. Van Nostrand Reinhold Company, New York, 970 p.
- Macfadyen, W.A., 1932, Foraminifera of some late Pliocene and glacial deposits of East Anglia. *Geological Magazine*, 49: 481-496.
- Machaniec, E., Zapałowicz-Bilan, B., 2005, Foraminiferal biostratigraphy and paleobathymetry of Senonian marls (Upper Cretaceous) in the vicinity of Kraków (Januszowice-Korkiew area, Bonarka quarry) - preliminary study. In *Methods and Applications in Micropaleontology* (Tyszk, J., Oliwkiewicz-Miklasinska, M., Gedl, P., Kaminski, M.A., Eds.). *Studia Geologica Polonica*, 124: 285-295.
- Marie, P., 1941, Les foraminifères de la craie à *Belemnites* du Bassin de Paris. *Mémoires du Muséum National d'Histoire Naturelle*, 12: 1-296.
- Marsson, T., 1878, Die Foraminiferen der weißen Schreibkreide der Insel Rügen. *Mitteilungen des Naturwissenschaftlichen Vereins für Neu-Vorpommern und Rügen in Greifswald*, 10: 115-196.
- Martínez-Pardo, R., 1965, *Bolivinoidea draco doreeni* Finlay from the Magellan Basin, Chile. *Micropaleontology*, 11: 360-364. <https://doi.org/10.2307/1484596>
- Martínez R.J.I., 1989, Foraminiferal biostratigraphy and paleoenvironments of the Maastrichtian Colon mudstones of northern South America. *Micropaleontology*, 35: 97-113. <https://doi.org/10.2307/1485462>

- Maslakova, N.I., 1959, Foraminifera. In *Atlas of Upper Cretaceous faunas of northern Caucasus and Crimea* (Moskvina, M.M., Ed.). Gosudarstvennoe Nauchno-tekhnicheskoe Izdatelstvo Neftianoi i Gorno-toplivnoi Literaturi, Moskva, 87-129 (In Russian).
- McGugan, A., 1957, Upper Cretaceous foraminifera from Northern Ireland. *Journal of Paleontology*, 31: 329-348. <http://www.jstor.org/stable/1300633>
- McGugan, A., 1974, Cretaceous foraminifera from Whitepark Bay, Northern Ireland. *Irish Naturalists Journal*, 18: 12-17.
- McNeely, B.W., 1973, Biostratigraphy of the Mesozoic and Paleocene pelagic sediments of the Campeche Embankment Area. In *Initial Reports of the Deep Sea Drilling Project* (Worzel, J.L., Bryant, W., et al., Eds.). United States Government Printing Office, Washington, D.C., 10: 679-695. <https://doi.org/10.2973/dsdp.proc.10.126.1973>
- McNeil, D.H., Caldwell, W.G.E., 1981, Cretaceous rocks and their foraminifera in the Manitoba Escarpment. *The Geological Association of Canada Special Papers*, 77: 1-439.
- Medizza, F., 1967, I generi *Bolivinoidea*, *Aragonia* e *Neoflabellina* (Foraminifera) nelle formazioni cretaceo-eoceniche del Veneto. *Memorie degli Istituti di Geologia e Mineralogia dell'Università di Padova*, 26: 3-44 (Published in 1969).
- Mello, J.F., 1969, Foraminifera and stratigraphy of the upper part of the Pierre Shale and lower part of the Fox Hills Sandstone (Cretaceous) north-central South Dakota. *United States Geological Survey Professional Paper*, 611: 1-117.
- Michalski, M., Vellekoop, J., Dubicka, Z., Peryt, D. & Harasimiuk, M., 2016, Late Maastrichtian cephalopods, dinoflagellate cysts and foraminifera from the Cretaceous-Paleogene succession at Lechówka, southeast Poland: stratigraphic and environmental implications. *Cretaceous Research*, 57: 208-227. <https://doi.org/10.1016/j.cretres.2015.08.012>
- Montanaro Gallitelli, E., 1956, *Bronnimannella*, *Tappanina*, and *Trachelinella*, three new foraminiferal genera from the Upper Cretaceous. *Contributions from the Cushman Foundation for Foraminiferal Research*, 7: 35-39.
- Montanaro Gallitelli, E., 1957, A revision of the foraminiferal family Heterohelcidae. In *Studies in Foraminifera* (Loeblich, A.R. Jr., Ed.). *United States National Museum Bulletin*, 215: 133-154.
- Moore, R.C., Lalicker, C.G. & Fischer, A.G., 1952, *Invertebrate Fossils*. McGraw-Hill Book Company, New York-Toronto-London, 766 p.
- Naidin, D.P., Beniamovski, V.N., 2006, The Campanian-Maastrichtian stage boundary in the Aktulagai section (North Caspian Depression). *Stratigraphy and Geological Correlation*, 14: 433-443.
- Neagu, T., 1979, *Micropaleontologie. Protozoare*. Editura Tehnică, București, 404 p. (In Romanian).
- Neagu, T., 1992, White chalk foraminiferal fauna from southern Dobrogea (Romania). III. Calcareous benthic foraminifera. *Revista Española de Micropaleontología*, 24: 59-115.
- Nederbragt, A.J., 1991, Late Cretaceous biostratigraphy and development of Heterohelcidae (planktic foraminifera). *Micropaleontology*, 37: 329-372. <https://doi.org/10.2307/1485910>
- Nestler, H., 1982, *Die Fossilien der Rügener Schreibkreide*. A. Ziemsen Verlag, Wittenberg Lutherstadt, 108 p.
- Niazi, S.M., Vahidinia, M. & Ashaori, A., 2013, Microbiostratigraphy and paleoecology of the Abtalkh Formation based on foraminifera in the Ghareh-Soo section (southwest Kalat-e-Naderi, northeast Iran). *Sedimentary facies*, 6: 95-114 (In Persian).
- Nomura, R., 1991, Paleoceanography of upper Maastrichtian to Eocene benthic foraminiferal assemblages at sites 752, 753, and 754, Eastern Indian Ocean. In *Proceedings of the Ocean Drilling Program, Scientific Results* (Weissel, J., Pierce, J., et al., Eds.). College Station, 121: 3-29. <https://doi.org/10.2973/odp.proc.sr.121.138.1991>
- Nomura, R., 1995, Paleogene to Neogene deep-sea paleoceanography in the eastern Indian Ocean: benthic foraminifera from ODP Sites 747, 757 and 758. *Micropaleontology*, 41: 251-290. <https://doi.org/10.2307/1485862>
- Nomura, R., Brohi, I.A., 1995, Benthic foraminiferal fauna during the time of the Indian-Asian contact, in southern Balochistan, Pakistan. *Marine Micropaleontology*, 24: 215-238. [https://doi.org/10.1016/0377-8398\(94\)00024-H](https://doi.org/10.1016/0377-8398(94)00024-H)
- Noth, R., 1951, Foraminiferen aus Unter- und Oberkreide des Österreichischen Anteils an Flysch, Helvetikum und Vorlandvorkommen. *Jahrbuch der Geologischen Bundesanstalt*, 3: 1-91.
- Ogg, J.G., Agterberg, F.P. & Gradstein, F.M., 2004, The Cretaceous Period. In *The geologic time scale 2004* (Gradstein, F.M., Ogg, J.G., Smith, A.G., Eds.). Cambridge University Press, Cambridge, p. 344-383.
- Ohmert, W., 2011, Radiolarien-Faunen und Stratigraphie der Pattenau-Formation (Campanium bis Maastrichtium) im Helvetikum von Bad Tölz (Oberbayern). *Zitteliana*, 51: 37-98.
- Olshanetskiy, D.M., 2015, Deep-water benthic foraminifers from the Paleocene and Eocene of the North Pacific region: paleontology, biostratigraphy and paleoceanological reconstructions. *Stratigraphy and Geological Correlation*, 23: 661-715. <https://doi.org/10.1134/S0869593815070023>
- Olsson, R.K., Nyong, E.E., 1984, A paleoslope model for Campanian-lower Maastrichtian foraminifera of New Jersey and Delaware. *Journal of Foraminiferal Research*, 14: 50-68. <https://doi.org/10.2113/gsjfr.14.1.50>
- Omaña, L., 2006, Late Cretaceous (Maastrichtian) foraminiferal assemblage from the inoceramid beds, Ocozocoaulta Formation, central Chiapas, SE Mexico. *Revista Mexicana de Ciencias Geológicas*, 23 (2): 125-132. <http://rmcg.geociencias.unam.mx/revista/index.html>
- Papp, A., Küpper, K., 1953, Die Foraminiferenfauna von Guttaring und Klein St. Paul (Kärnten). 1. Über Globotruncanen südlich Pernerberg bei Klein St. Paul. *Sitzungsberichte der Österreichischen Akademie der Wissenschaften. Mathematisch-Naturwissenschaftliche Klasse*, 162: 31-48.
- Perlmutter, N.M. and Todd, R., 1965, Correlation and foraminifera of the Monmouth Group (Upper Cretaceous) Long Island, New York. *United States Geological Survey Professional Paper*, 483-I: 1-21.

- Peryt, D., 2000. On the age of siliceous chalk in the Piotrawin Quarry, middle Vistula River valley, central Poland. *Biuletyn Państwowego Instytutu Geologicznego*, 393: 81-94 (In Polish).
- Petters, S.W., 1976, Upper Cretaceous subsurface stratigraphy of Atlantic coastal plain of New Jersey. *The American Association of Petroleum Geologists Bulletin*, 60: 87-107. <http://archives.datapages.com/data/bulletins/1974-76/data/pg/0060/0001/0050/0087.htm>
- Petters, S.W., 1977, *Bolivinoides* evolution and Upper Cretaceous biostratigraphy of the Atlantic Coastal Plain of New Jersey. *Journal of Paleontology*, 51: 1023-1036. <http://www.jstor.org/stable/1303775>
- Pérez Estrada, L.M., Bustamante, S.B., Nieves, A.F., Sánchez, R.G. & León, J.H., 2007, Consideraciones bioestratigráficas de la Formación Vía Blanca Basadas en foraminíferos bentónicos pequeños: su relación con otros esquemas biozonales. *Segunda Convención Cubana de Ciencias de la Tierra, Geociencias 2007*, p. 1-10.
- Plotnikova, L.F., 1967, *Shallow water foraminifera of the Upper Cretaceous of the platform part of the USSR (Konksk-Jalinsk Basin and northern part of the Black Sea surroundings)*. Kiev, Naukova Dumka, 107 p. (In Russian).
- Plummer, H.J., 1931, Some Cretaceous Foraminifera in Texas. *The University of Texas Bulletin*, 3101: 109-203.
- Pokorný, V., 1958, *Grundzüge der Zoologischen Mikropaläontologie, Band I*. Berlin, VEB Deutscher Verlag der Wissenschaften, 582 p. (English translation, 1963, Pergamon Press).
- Požaryska, K., 1954, The Upper Cretaceous index foraminifera from Central Poland. *Acta Geologica Polonica*, 4: 249-276 (In Polish). <https://geojournals.pgi.gov.pl/agp/article/view/12140>
- Požaryska, K., 1965, Foraminifera and biostratigraphy of the Danian and Montian in Poland. *Palaeontologia Polonica*, 14: 1-156. <http://www.palaeontologia.pan.pl/>
- Pringgoprawiro, H., 1965, Some significant Upper Cretaceous foraminifera from Groisbach, Morzger Hügel, and Michelstetten, Austria. *Contributions from the Department of Geology, Institute of Technology Bandung*, 60: 21-65.
- Rasheed, D.A., Ravindran, C.N., 1980, The occurrence of *Bolivinoides decoratus* (Jones) from the Ariyalur stage of the Trichinopoly Cretaceous, Tamil Nadu. *Journal of the Geological Society of India*, 21: 206-207.
- Rauzer-Chernousova, D.M., Fursenko, A.V., 1959, *Principles of Paleontology, Part I. Protista*. Akademiya Nauk SSSR, Moskva, 481 p. (In Russian).
- Remin, Z., Dubicka, Z., Kozłowska, A. & Kuchta, B., 2012. A new method of rock disintegration and foraminiferal extraction with the use of liquid nitrogen [LN₂]. Do conventional methods lead to biased paleoecological and paleoenvironmental interpretations? *Marine Micropaleontology*, 86-87: 11-14. <https://doi.org/10.1016/j.marmicro.2011.12.001>
- Reiss, Z., 1954, Upper Cretaceous and lower Tertiary *Bolivinoides* from Israel. *Contributions from the Cushman Foundation for Foraminiferal Research*, 5: 154-164.
- Rescher, K., 1991, Biostratigraphische Gliederung der höheren Oberkreide mit benthonischen Foraminiferen im nordwestlichen Münsterland (Bohrung Metelen 1001). *Facies*, 24: 99-106.
- Revets, S.A., 1996, The generic revision of five families of rotaline foraminifera. Part I. The Bolivinitidae Cushman, 1927. *Cushman Foundation for Foraminiferal research Special Publications*, 34: 1-55.
- Robaszynski, F., Bless, M.J.M., Felder, P.J., Foucher, J.-C., Legoux, O., Manivit, H., Meessen, J.P.M.T. & Van der Tuuk, L.A., 1985, The Campanian-Maastrichtian boundary in the chalk facies close to the type-Maastrichtian area. *Bulletin des Centres de Recherches Exploration-Production Elf-Aquitaine*, 9: 1-113.
- Robaszynski, F., Caron, M., González Donoso, J.M., Wonders, A. A. H. & The European Working Group on Planktonic Foraminifera, 1984, Atlas of Late Cretaceous globotruncanids. *Revue de Micropaléontologie*, 26: 145-305.
- Robaszynski, F., Christensen, W.K., 1989, The upper Campanian-lower Maastrichtian chalks of the Mons basin, Belgium: a preliminary study of belemnites and foraminifera in the Harmignies and Ciply areas. *Geologie en Mijnbouw*, 68: 391-408.
- Rozumeiko, S.V., 1978. *Upper Cretaceous foraminifera from the deposits of the southwestern part of the Eastern European Platform*. Kiev, Naukova Dumka, 176 p. (In Russian).
- Sacal, V., Debourle, A., 1957, Foraminifères d'Aquitaine 2^e partie. Peneroplidae a Victoriellidae. *Mémoires de la Société Géologique de France*, 78: 1-87.
- Said, R., Kenawy, A., 1956, Upper Cretaceous and Lower Tertiary foraminifera from northern Sinai, Egypt. *Micropaleontology*, 2: 105-173. <https://doi.org/10.2307/1484100>
- Salaj, J., 1980, *Microbiostratigraphie du Crétacé et du Paleogene de la Tunisie septentrionale et orientale (Hypostratotypes tunisiens)*. Geologický Ústav Dionýza Štúra, Bratislava, 232 p.
- Salaj, J., 1988, Benthic foraminifera of the *Archaeoglobituncana kefiana* Zone from the basal Maastrichtian in the area of El Kef (Tunisia). *Revue de Paléobiologie, Volume spéciale*, 2: 333-338.
- Salaj, J., Maamouri, A.-L., 1982, *Rugotruncana kefiana* n.sp. (foraminifera) du Sénonien supérieur d'El Kef. *Geologica Carpathica*, 33 (4): 463-479. <http://www.geologicacarthica.com/browse-journal/authors/joseph-salaj/>
- Sandidge, J.R., 1932, Significant foraminifera from the Ripley Formation of Alabama. *The American Midland Naturalist*, 13: 190-202.
- Scheibnerová, V., 1968, Middle and Upper Cretaceous microbiostratigraphy of the Klippen Belt (West Carpathians). *Acta Geologica et Geographica Universitatis Comenianae, ser. Geologica*, 17: 5-98.
- Schijfsma, E., 1946, The foraminifera from the Hervian (Campanian) of southern Limburg. *Mededeelingen van de Geologische Stichting*, 7: 1-174.
- Schönfeld, J., 1990, Zur Stratigraphie und Ökologie bentischer Foraminiferen im Schreiekreide-Richtprofil von Lägerdorf/Holstein. *Geologisches Jahrbuch*, A117: 3-151.

- Sellier de Civrieux, J.M., 1952, Estudio de la microfauna de la seccion-tipo del Miembro Socuy de la Formacion Colon Distrito Mara, Estado Zulia. *Ministerio de Minas e Hidrocarburos Direccion de Geologia*, 2: 231-310.
- Shabestari, G.M., Raisossadat, N., Shakaram, M.A. & Jalili, F., 2014, Biostratigraphy of the Upper Cretaceous deposits in north of Birjand (Shushud section). *Journal of Stratigraphy and Sedimentology Researches*, 30: 47-61.
- Sheldon, E., Gravesen, P. & Nøhr-Hansen, H., 2012, Geology of the Femern Bælt area between Denmark and Germany. *Geological Survey of Denmark and Greenland Bulletin*, 26: 13-16.
- Shimer, H.W., Shrock, R.R., 1944, *Index fossils of North America*. The M.I.T. Press, Cambridge, Massachusetts, 837 p.
- Sliter, W.V., 1968, Upper Cretaceous foraminifera from Southern California and northwestern Baja California, Mexico. *The University of Kansas Paleontological Contributions*, 49: 1-141.
- Sliter, W.V., 1977a, Cretaceous foraminifers from the southwestern Atlantic Ocean, Leg 36, Deep Sea Drilling Project. In *Initial Reports of the Deep Sea Drilling Project* (Barker, P. et al., Eds). United States Government Printing Office, Washington, D.C., 36: 519-573.
<https://doi.org/10.2973/dsdp.proc.36.110.1977>
- Sliter, W.V., 1977b, Cretaceous benthic foraminifers from the western South Atlantic Leg 39, Deep Sea Drilling Project. In *Initial Reports of the Deep Sea Drilling Project* (Supko, P.R., Ed.). United States Government Printing Office, Washington, D.C., 39: 657-697.
<https://doi.org/10.2973/dsdp.proc.39.130.1977>
- Sigal, J., 1952, Ordre des foraminifera. In *Traité de Paléontologie* (Piveteau, J., Ed.). Maisson et Cie, Paris, 1: 133-301.
- Speijer, R.P., van der Zwaan, G.J., 1995, Extinction and survivorship patterns in southern Tethyan benthic foraminiferal assemblages across the Cretaceous/Paleogene boundary. In *Extinction and recovery patterns in benthic foraminiferal paleocommunities across the Cretaceous/Paleogene and Paleocene/Eocene boundaries* (Speijer, R.P., Ed.). *Geologica Ultraiectina*, 124: 19-64.
- Stenestad, E., 1968, Nogle kridtlignende kvartæraflejringer i Nordjylland. *Meddelelser fra Dansk Geologisk Forening*, 18: 285-293.
- Strong, C.P., Edwards, A.R., 1979, Late Haumurian (Maastrichtian) microfossils from Chatham Islands, New Zealand. *N.Z. Journal of Geology and Geophysics*, 22: 613-619.
- Swiecicki, A., 1980, *A foraminiferal biostratigraphy of the Campanian and Maastrichtian Chalks of the United Kingdom*. Ph.D. Thesis, Plymouth University, 1: 1-358, 2: 1-155.
- Sztrákos, K., 2005, Les foraminifères du Paléocène et de l'Éocène basal du sillon nord-pyrénéen (Aquitaine, France). *Revue de Micropaléontologie*, 48: 175-236.
- Szczuchura, J., Pożaryska, K., 1974, Foraminiferida from the Paleocene of Polish Carpathians (Babica Clays). *Palaeontologia Polonica*, 31: 1-142.
<http://www.palaeontologia.pan.pl/>
- Thomas, E., 1990, Late Cretaceous through Neogene benthic foraminifers (Maud Rise, Weddell Sea, Antarctica). In *Proceedings of the Ocean Drilling Program, Scientific Results* (Barker, P.F., Kennett, J.P., et al., Eds.). College Station, 113: 571-594.
<https://doi.org/10.2973/odp.proc.sr.113.123.1990>
- Tjalsma, R.C., Lohmann, G.P., 1983, Paleocene-Eocene bathyal and abyssal foraminifera from the Atlantic Ocean. *Micropaleontology, Special publications*, 4: 1-90.
- Tronchetti, G., 2001, Les foraminifères benthiques des affleurements Campano-maastrichtiens de Tercis les Bains (Landes, France). In *The Campanian-Maastrichtian Stage Boundary. Characterisation and Correlation from Tercis-les-Bains (Landes, SW France) to Europe and Other Continents* (Odin, G.S., Ed.). Elsevier Science, Amsterdam, 314-327.
- Van Hinte, J., 1963, Zur Stratigraphie und Mikropaläontologie der Oberkreide und des Eozäns des Krappfeldes (Kränten). *Jahrbuch der Geologischen Bundesanstalt*, 8: 1-147.
- Van Hinte, J.E., 1965, Remarks on the Kainach Gosau (Styria, Austria). *Proceedings of the Koninklijke Nederlandse Akademie van Wetenschappen (B)*, 68: 72-92.
- Van Hinte, J.E., 1967, *Bolivinoidea* from the Campanian type section. *Proceedings of the Koninklijke Nederlandse Akademie van Wetenschappen (B)*, 70: 254-263.
- Van Hinte, J.E., 1976, A Cretaceous time scale. *The American Association of Petroleum Geologists Bulletin*, 60: 269-287.
<https://doi.org/10.1306%2F83D923ED-16C7-11D7-8645000102C1865D>
- Van Morkhoven, F.P.C.M., Berggren, W.A. & Edwards, A.S., 1986, Cenozoic cosmopolitan deep-water benthic foraminifera. *Bulletin des Centres de Recherches Exploration-Production Elf-Aquitaine*, 11: 1-421.
- Vaptzarova, J., 1974, Foraminifera of the Carpathian-type Maastrichtian of Northwest Bulgaria. *Proceedings of the Geological Institute, series Paleontology*, 23: 31-56 (In Bulgarian).
- Vassilenko, V.P., 1961, Foraminifera of the Upper Cretaceous of Mangyshlak Peninsula. *Trudy VNIGRI*, 171: 1-487 (In Russian).
- Villain, J.-M., 1977, Le Maastrichtien dans sa région type (Limbourg, Pays-Bas). Étude stratigraphique et micropaléontologique. *Palaeontographica*, A157: 1-87.
- Voloshina, A.M., 1974, Review of the Upper Cretaceous buliminid genera from the material from Volno-Podolia. *Paleontologicheskii Sbornik*, 10: 17-22 (In Russian).
- Walaszczyk, I., Dubicka, Z., Olszewska-Nejbert, D. & Remin, Z., 2016, Integrated biostratigraphy of the Santonian through Maastrichtian (Upper Cretaceous) of extra-Carpathian Poland. *Acta Geologica Polonica*, 66: 313-350.
<https://geojournals.pgi.gov.pl/agp/article/view/25564>
- White, M.P., 1929, Some index foraminifera of the Tampico Embayment area of Mexico. Part III. *Journal of Paleontology*, 3, 30-57.
- Wicher, C.A., 1949, On the age of the higher Upper Cretaceous of the Tampico Embayment areas in Mexico, as an example of the worldwide existence of microfossils and the practical consequences arising from this. *Bulletin du Muséum National d'Histoire Naturelle du Pays Serbe*, 2: 76-105.
- Wicher, C.A., 1953, Micropaläontologische beobachtungen in der höheren borealen Oberkreide besonders im Maastricht. *Geologisches Jahrbuch*, 68: 1-26. (Published in 1954).

- Widmark, J.G.V., 1997. Deep-sea benthic foraminifera from Cretaceous-Paleogene boundary strata in the South Atlantic-taxonomy and paleoecology. *Fossils and Strata*, 43: 1-94.
- Williams-Mitchell, E., 1948, The zonal value of the foraminifera in the chalk of England. *Proceedings of the Geologists' Association*, 59: 91-112.
- Witte, L., Schuumann, H., 1996, Calcareous benthic foraminifera across the Cretaceous/Tertiary boundary in the Geulhemmerberg (SE Netherlands). *Geologie en Mijnbouw*, 75: 173-185.
- Witwicka, E., 1958, Cretaceous stratigraphic micropaleontology from a borehole from Chelm. *Biuletyn Instytut Geologiczny*, 121: 177-232 (In Polish).
- Wright, J., 1886, A list of the Cretaceous foraminifera of Keady Hill, County Derry. *Proceedings of the Belfast Naturalists' Field Club*, 2: 327-332.
- Yasuda, H., 1986, Cretaceous and Paleocene foraminifera from northern Hokkaido, Japan. *Science Reports of the Tohoku University, Sendai, Second Series (Geology)*, 57: 1-101.
- Yeremeyeva, A.I., 1957. Some new species of foraminifera from the Cretaceous and Tertiary deposits of the eastern flank of the Urals. *Trudy Gorno-Geologicheskaya Instituta, Akademyia Nauk SSSR*, 28: 5-13 (In Russian).
- Yeremeyeva, A.I., Belousova, N.A., 1961. Stratigraphy and foraminiferal fauna of the Cretaceous and Paleogene sediments of western Urals, pre-Urals and northern Kazakhstan. *Materialy po Geologii i Poleznym Iskopaemim Urala*, 9: 3-189 (In Russian).
- Ziko, A., Darwish, M. & Eweda, S., 1993, Late Cretaceous-Early Tertiary stratigraphy of the Themed area, East Central Sinai, Egypt. *Neues Jahrbuch für Geologie und Paläontologie Monatshefte*, 1993 (H3): 135-149.