

2019

Pepper Mild Mottle Virus: Agricultural Menace Turned Effective Tool for Microbial Water Quality Monitoring and Assessing (Waste)water Treatment Technologies

Erin Symonds
University of South Florida

Karyna Rosario
University of South Florida

Mya Breitbart
University of South Florida

Follow this and additional works at: https://scholarcommons.usf.edu/msc_facpub

 Part of the [Life Sciences Commons](#)

Scholar Commons Citation

Symonds, Erin; Rosario, Karyna; and Breitbart, Mya, "Pepper Mild Mottle Virus: Agricultural Menace Turned Effective Tool for Microbial Water Quality Monitoring and Assessing (Waste)water Treatment Technologies" (2019). *Marine Science Faculty Publications*. 646.

https://scholarcommons.usf.edu/msc_facpub/646

This Article is brought to you for free and open access by the College of Marine Science at Scholar Commons. It has been accepted for inclusion in Marine Science Faculty Publications by an authorized administrator of Scholar Commons. For more information, please contact scholarcommons@usf.edu.

Pepper mild mottle virus: Agricultural menace turned effective tool for microbial water quality monitoring and assessing (waste)water treatment technologies

Erin M. Symonds, Karyna Rosario, Mya Breitbart

Published: April 18, 2019 • <https://doi.org/10.1371/journal.ppat.1007639>

Citation: Symonds EM, Rosario K, Breitbart M (2019) Pepper mild mottle virus: Agricultural menace turned effective tool for microbial water quality monitoring and assessing (waste)water treatment technologies. *PLoS Pathog* 15(4): e1007639. <https://doi.org/10.1371/journal.ppat.1007639>

Editor: Katherine R. Spindler, University of Michigan Medical School, UNITED STATES

Published: April 18, 2019

Copyright: © 2019 Symonds et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was funded by grant OCE-1566562 from the US National Science Foundation. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the US National Science Foundation.

Competing interests: The authors have declared that no competing interests exist.

Introduction

Domestic wastewater pollution in environmental waters or water reuse supplies represents a threat to public health because of high concentrations of diverse pathogens associated with human excreta [1]. Since it is difficult to directly measure waterborne pathogens of concern, microbial water quality monitoring efforts often use surrogates or indicator organisms that are easily detected and whose presence reflects pathogen persistence [2]. Here, we describe an unconventional viral indicator of wastewater pollution, pepper mild mottle virus (PMMoV), a plant pathogen that was first proposed as a water quality indicator in 2009 [3] and promises to improve microbial water quality management worldwide [4].

Why is PMMoV found in human feces and wastewater?

Considering that most recognized viral pathogens causing human gastrointestinal disease contain RNA genomes [5], early metagenomics studies sought to characterize the RNA viral communities (i.e., viromes) in the feces of healthy individuals [6]. Unexpectedly, these viromes were dominated by sequences similar to plant-infecting viruses that are acquired through dietary consumption of plant products [6]. By far, the most abundant RNA virus identified in the feces of healthy individuals was PMMoV [6], which has also been readily identified in untreated wastewater from numerous locations (summarized in [4, 7]; Fig 1). PMMoV infects various species of peppers (*Capsicum* spp.), and its presence in human feces originates from the consumption of infected peppers, which are frequently used in processed pepper products (e.g., hot sauces, curry sauces, dry spices; Fig 1) [6, 8, 9]. Up to 10^8 PMMoV gene copies are found in a single milliliter of hot sauce [9], and healthy individuals can excrete up to 10^9 gene copies in 1 gram of feces [6]. Although the presence of PMMoV in human feces presumably depends on dietary consumption of pepper products, which in turn may depend on age and food preferences of a given individual, PMMoV appears to be widespread and globally distributed, given that this virus has been consistently detected in untreated wastewater from Africa, the Americas, Asia, Australia, and Europe (summarized in [4, 7]). PMMoV can also be detected in treated wastewater effluent and in environments impacted by wastewater discharges (treated or untreated; Fig 1).




Fig 1. The plant pathogen PMMoV has emerged as a useful viral indicator of wastewater pollution because of its presence throughout the domestic wastewater cycle.

PMMoV is excreted from humans in high concentrations after consumption of processed food products, such as hot sauces, that contain infected peppers (gray arrow). Because PMMoV is consistently found at high concentrations in domestic wastewater, it can be used as a marker to trace untreated (purple arrows) or treated (blue arrows) wastewater in the environment. Humans are commonly exposed (green arrows) to water and food resources affected directly or indirectly by wastewater discharges through the consumption of drinking water, fresh agricultural produce, or shellfish raised in polluted environmental waters, as well as the accidental ingestion of environmental water (e.g., lakes, reservoirs, rivers, coastal waters). The viral surrogate PMMoV reflects the presence and persistence of human enteric viruses in such resources, serving as a vital tool for monitoring microbial water quality, (waste)water treatment efficiency, and food safety. *Artwork by Anne Martin (hungrybraindesign.com).* PMMoV, pepper mild mottle virus.
<https://doi.org/10.1371/journal.ppat.1007639.g001>

How can we take advantage of PMMoV as an indicator of wastewater pollution?

Microbial surrogates used to characterize fecal pollution in environmental monitoring programs traditionally use fecal indicator bacteria (FIBs; e.g., *Escherichia coli* and enterococci), a group of bacteria that are native to the guts of animals. FIBs are the most affordable and commonly used fecal pollution indicators in the world. Unfortunately, FIB presence does not always correlate with human pathogens and/or human health risks, especially in the case of human enteric viruses [10–12]. Moreover, the detection of FIBs may not accurately reflect fecal contamination because of their ecology and extraintestinal reservoirs in soil and sediment [10, 13]. Another strategy for tracking fecal pollution is to monitor for specific reference viral pathogens of interest, such as norovirus and adenovirus. However, analytical tools for detecting reference enteric viruses are expensive, laboratory intensive, and often have efficiencies less than 10% [14, 15]. In addition, enteric viruses are typically found in low concentrations (<10⁵ copies per liter) in domestic wastewater in the absence of an outbreak [16] and further diluted upon discharge to environmental waters, which hinders their detection [14, 15].

Because viral enteric pathogens are not practical indicators of fecal pollution, viruses infecting gut bacteria have also been investigated. Viruses that infect FIBs, such as male-specific and somatic coliphages, are the most often used viral surrogates of fecal pollution; however, they do not always correlate with the presence of enteric viruses in the environment or during treatment processes (summarized in [14]). CrAssphage, a virus that infects human gut bacteria (*Bacteroides intestinalis*), was recently highlighted as a promising viral surrogate of human fecal pollution [17, 18]. Studies have found that it is highly human specific and is found in similar concentrations as the microbial source-tracking marker *Bacteroides* HF183 in wastewater and in polluted environmental waters. Future research is needed to understand the potential applications of this novel indicator, how it correlates with human pathogens, and its advantages over existing microbial surrogates of fecal pollution, including PMMoV.

PMMoV is consistently detected in domestic wastewater throughout the world, often at higher concentrations than human enteric pathogens; thus, it has been used as a viral surrogate to identify fecal pollution in the environment (summarized in [4, 7]; Fig 1). PMMoV is a unique viral indicator that offers several advantages over traditional indicators. First, concentrations of up to 10¹⁰ PMMoV gene copies per liter of domestic wastewater circumvent traditional methodological limitations associated with molecular detection of viruses in low concentrations, thus facilitating monitoring efforts. Although PMMoV concentrations vary over space and time, 10⁶ to 10¹⁰ gene copies per liter of domestic wastewater are consistently detected. Second, PMMoV's dietary origin means that its concentration in domestic wastewater does not depend on active viral infection in the human population. This is an important consideration because it is not practical to regularly test for all known enteric pathogens, which exhibit different infection dynamics and seasonality. Third, unlike FIBs, PMMoV has not been detected in environmental waters free of fecal pollution and is typically not found in animals other than humans. However, PMMoV concentrations much lower than those found in domestic wastewater have been reported from birds (chickens, geese, swans, and seagulls), dogs, and cows (summarized in [4, 7]; [19]). Consequently, the use of PMMoV as a human-specific indicator for source-tracking purposes needs further evaluation. Nevertheless, PMMoV has been recognized as a promising viral indicator of domestic wastewater, with detection in environmental waters that is even comparable to nonbiological surrogates of wastewater contamination (e.g., caffeine, pharmaceuticals; summarized in [4, 7]).

Is PMMoV detection indicative of health risks associated with exposure to wastewater pollution?

Ideally, the detection of a surrogate organism should correlate with health risks associated with using water resources impacted by wastewater. The high concentration of PMMoV in treated domestic wastewater effluent and high detection rates in environmental waters originally led to PMMoV being proposed as an ultraconservative wastewater indicator, whose detection may not correlate with enteric pathogens or health risks [3, 20, 21]. However, PMMoV detection and quantification has now been associated with human enteric pathogens and health risks exceeding benchmark limits [22–25]. Furthermore, PMMoV persistence in seawater and river water is comparable to that of human enteric viruses, ranging from 7 to 21 days, depending upon water temperatures [3, 24]. Data combined from multiple studies investigating PMMoV detection in parallel with human pathogens in environmental waters (groundwater, seawater, and surface freshwater) revealed that PMMoV was detected consistently (94.2% of samples) and co-occurred with enteric pathogens in the majority (72.3%) of tested samples [4]. Notably, PMMoV was most often detected along with human pathogens in waters exposed to untreated domestic wastewater and least often in the absence of known pollution sources (11.4% PMMoV-positive samples, with 0% pathogen co-occurrence). Therefore, PMMoV seems to be a conservative tracer, but its detection and concentration correlate with the presence of human enteric pathogens in environmental waters exposed to wastewater contamination.

PMMoV detection and quantification also correlated with health risks associated with recreation in contaminated environmental waters. In coastal waters contaminated with treated and untreated domestic wastewater, PMMoV detection corresponded to health risks greater than the United States Environmental Protection Agency (US EPA) health benchmark for safe recreation [22, 23]. For coastal waters with untreated domestic wastewater pollution, PMMoV concentrations greater than 5.4 × 10³ gene copies per liter corresponded to health benchmark exceedance [23]. In contrast, for coastal waters exposed to secondary treated domestic wastewater pollution, 1.8 × 10³ PMMoV gene copies per liter corresponded to the US EPA health benchmark for safe recreation.

Future research is needed to better understand the correlations between PMMoV, enteric pathogens, and health risks in other environmental water matrices (e.g., freshwater), in different climates, and in the presence of tertiary treated wastewater.

Can PMMoV be used to evaluate the efficiency of water treatment processes?

To protect public health as well as to ensure the microbial safety of drinking water, disinfection treatment processes need to reduce virus concentrations in wastewater prior to direct water reuse or discharge into environmental waters (Fig 1) [12, 26]. It is often difficult to determine the log reduction of viruses achieved by a given treatment technique because many viruses are found in concentrations too low to quantify and human enteric viruses of public health concern, such as norovirus, lack culture-based methods or are difficult to culture. Consequently, molecular methods such as quantitative reverse transcription PCR are commonly used to test virus reductions. In these contexts, PMMoV has proven to be an ideal virus process indicator to evaluate drinking water, wastewater, and water reclamation treatment technologies and facilities because, unlike other viruses, its high concentrations in wastewater allow for the quantification of virus gene copy removal at full-scale treatment plants [27–30]. Moreover, PMMoV reduction levels during treatment of wastewater [16, 30, 31] and drinking water are usually comparable to those of human enteric viruses [27–29]. PMMoV gene copy removal more often has a significant, positive correlation with the gene copy removal of human enteric viruses in comparison with the gene copy removal of other viral indicators (bacteriophages MS2 and φX174) during drinking water treatment (at plants with coagulation-sedimentation, rapid sand filtration, ozonation, and biological activated carbon treatments) [28]. In addition to having practical applications for measuring virus reduction at the full-scale treatment facilities, PMMoV is also useful for quantifying virus reductions by innovative water treatment technologies at smaller scales (summarized in [4]) as well as point-of-use household drinking water treatment [32] and on-farm riverbank filtration systems [33]. Because PMMoV is currently quantified using molecular methods that cannot determine virus infectivity, the incorporation of culture-based analyses (which would require plant growth chambers) [8] and/or selective pretreatment for infectious particles could improve future virus reduction analyses to assess treatment efficiency [34].

How can we better exploit PMMoV to safeguard public health?

In addition to being a viral indicator for water quality monitoring purposes, PMMoV holds potential as a surrogate for enteric viruses in the assessment of food safety (Fig 1). Although more investigations are needed to confirm the application of PMMoV in agricultural contexts, PMMoV has been a useful surrogate for enteric viruses on crops irrigated directly and indirectly with domestic wastewater [33, 35]. Specifically, PMMoV concentrations determined in irrigation water enabled subsequent quantitative microbial risk assessments, which in turn facilitated the evaluation of best practices and practical recommendations regarding safe water reuse strategies for consumers. Furthermore, PMMoV frequently co-occurred with human norovirus in shellfish collected from coastal areas [4] and was consistently detected in shellfish exposed to point source fecal pollution [19], suggesting that PMMoV may be a useful enteric virus surrogate to monitor the microbial quality of shellfish harvested for raw consumption.

From the serendipitous discovery that plant viruses dominate human feces, PMMoV has emerged as a unique surrogate for enteric viral pathogens with proven applications for monitoring microbial water quality, (waste)water treatment efficiency, and food safety (summarized in [4]). Currently, PMMoV is detected and quantified using virus concentration and molecular methods, which are often too time consuming and expensive for use in widespread monitoring. Future research is warranted to develop rapid, lab-free, inexpensive methods for PMMoV detection, such as “dipstick”-type approaches based on immunological assays [14]. Development of more accessible detection methods will enable further exploitation of PMMoV, traditionally considered an agricultural threat, as a vital tool for protecting public health.

References

1. UNICEF; World Health Organization. Progress on sanitation and drinking water: 2015 update and MDG assessment. New York: UNICEF and World Health Organization; 2015.
2. Sinclair RG, Rose JB, Hashsham SA, Gerba CP, Haas CN. A criteria for selection of surrogates used to study the fate and control of pathogens in the environment. *Appl Environ Microbiol*. 2012;78(6):1969–77. pmid:22247166
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
3. Rosario K, Symonds EM, Sinigalliano C, Stewart J, Breitbart M. Pepper mild mottle virus as an indicator of fecal pollution. *Appl Environ Microbiol*. 2009;75(22):7261–7. pmid:19767474
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
4. Symonds EM, Nguyen KH, Harwood VJ, Breitbart M. Pepper mild mottle virus: A plant pathogen with a greater purpose in (waste)water treatment development and public health management. *Water Res*. 2018;144:1–12. pmid:30005176
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
5. Clark B, McKendrick M. A review of viral gastroenteritis. *Curr Opin Infect Dis*. 2004;17(5):461–9. pmid:15353966
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
6. Zhang T, Breitbart M, Lee WH, Run JQ, Wei CL, Soh SWL, et al. RNA viral community in human feces: Prevalence of plant pathogenic viruses. *PLoS Biol*. 2006;4(1):108–18. pmid:16336043
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
7. Kitajima M, Sassi HP, Torrey JR. Pepper mild mottle virus as a water quality indicator. *NPJ Clean Water*. 2018;1(1):19.
[View Article](#) • [Google Scholar](#)
8. Colson P, Richet H, Desnues C, Balique F, Moal V, Grob J-J, et al. Pepper mild mottle virus, a plant virus associated with specific immune responses,

- fever, abdominal pains, and pruritus in humans. *PLoS ONE*. 2010;5(4):e10041. pmid:20386604
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
9. Peng J, Shi B, Zheng H, Lu Y, Lin L, Jiang T, et al. Detection of pepper mild mottle virus in pepper sauce in China. *Arch Virol*. 2015;160(8):2079–82. pmid:26021835
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
 10. Lin J, Ganesh A. Water quality indicators: bacteria, coliphages, enteric viruses. *Int J Environ Health Res*. 2013;23(6):484–506. pmid:23438312
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
 11. Harwood VJ, Levine AD, Scott TM, Chivukula V, Lukasik J, Farrah SR, et al. Validity of the indicator organism paradigm for pathogen reduction in reclaimed water and public health protection. *Appl Environ Microbiol*. 2005;71(6):3163–70. pmid:15933017
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
 12. Gerba CP, Betancourt WQ, Kitajima M. How much reduction of virus is needed for recycled water: A continuous changing need for assessment? *Water Res*. 2017;108:25–31. pmid:27838026
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
 13. Byappanahalli MN, Nevers MB, Korajkic A, Staley ZR, Harwood VJ. Enterococci in the environment. *Microbiol Mol Biol Rev*. 2012;76(4):685–706. pmid:23204362
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
 14. Symonds EM, Breitbart M. Affordable enteric virus detection techniques are needed to support changing paradigms in water quality management. *CLEAN—Soil, Air, Water*. 2015;43(1):8–12.
[View Article](#) • [Google Scholar](#)
 15. Haramoto E, Kitajima M, Hata A, Torrey JR, Masago Y, Sano D, et al. A review on recent progress in the detection methods and prevalence of human enteric viruses in water. *Water Res*. 2018;135:168–86. pmid:29471200
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
 16. Kitajima M, Iker BC, Pepper IL, Gerba CP. Relative abundance and treatment reduction of viruses during wastewater treatment processes—Identification of potential viral indicators. *Sci Total Environ*. 2014;488–489:290–6. pmid:24836386
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
 17. Stachler E, Kelly C, Sivaganesan M, Li X, Bibby K, Shanks OC. Quantitative CrAssphage PCR assays for human fecal pollution measurement. *Environ Sci Technol*. 2017;51(16):9146–54. pmid:28700235
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
 18. Khokhlova EV, Fitzgerald CB, Stockdale SR, Draper LA, Ross RP, et al. ΦCrAss001 represents the most abundant bacteriophage family in the human gut and infects *Bacteroides intestinalis*. *Nat. Commun*. 2018;9(1):4781. pmid:30429469
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
 19. Gyawali P, Croucher D, Ahmed W, Devane M, Hewitt J. Evaluation of pepper mild mottle virus as an indicator of human faecal pollution in shellfish and growing waters. *Water Res*. 2019; 154:370–376. pmid:30822597
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
 20. Kuroda K, Nakada N, Hanamoto S, Inaba M, Katayama H, Do AT, et al. Pepper mild mottle virus as an indicator and a tracer of fecal pollution in water environments: Comparative evaluation with wastewater-tracer pharmaceuticals in Hanoi, Vietnam. *Sci Total Environ*. 2015;506–507:287–98. pmid:25460962
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
 21. Haramoto E, Kitajima M, Kishida N, Konno Y, Katayama H, Asami M, et al. Occurrence of pepper mild mottle virus in drinking water sources in Japan. *Appl Environ Microbiol*. 2013;79(23):7413–8. pmid:24056461
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
 22. Symonds EM, Sinigalliano C, Gidley ML, Ahmed W, McQuaig S, Breitbart M. Faecal pollution along the southeastern coast of Florida and insight into the use of pepper mild mottle virus as an indicator. *J Appl Microbiol*. 2016;121(5):1469–81. pmid:27501154
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
 23. Ahmed W, Hamilton KA, Lobos A, Hughes B, Staley C, Sadowsky MJ, et al. Quantitative microbial risk assessment of microbial source tracking markers in recreational water contaminated with fresh untreated and secondary treated sewage. *Environ Int*. 2018;117:243–9. pmid:29772486
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
 24. Hamza IA, Jurzik L, Überla K, Wilhelm M. Evaluation of pepper mild mottle virus, human picobirnavirus and torque teno virus as indicators of fecal contamination in river water. *Water Res*. 2011;45(3):1358–68. pmid:21074236

[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)

25. Hughes B, Beale DJ, Dennis PG, Cook S, Ahmed W. Cross-comparison of human wastewater-associated molecular markers in relation to fecal indicator bacteria and enteric viruses in recreational beach waters. *Appl Environ Microbiol.* 2017;83(8). pii: e00028–17. pmid:28159789
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)

26. Sano D, Amarasiri M, Hata A, Watanabe T, Katayama H. Risk management of viral infectious diseases in wastewater reclamation and reuse: Review. *Environ Int.* 2016;91:220–9. pmid:26985655
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)

27. Asami T, Katayama H, Torrey JR, Visvanathan C, Furumai H. Evaluation of virus removal efficiency of coagulation-sedimentation and rapid sand filtration processes in a drinking water treatment plant in Bangkok, Thailand. *Water Res.* 2016;101:84–94. pmid:27258619
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)

28. Kato R, Asami T, Utagawa E, Furumai H, Katayama H. Pepper mild mottle virus as a process indicator at drinking water treatment plants employing coagulation-sedimentation, rapid sand filtration, ozonation, and biological activated carbon treatments in Japan. *Water Res.* 2018;132:61–70. pmid:29306700
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)

29. Lee S, Hata A, Yamashita N, Tanaka H. Evaluation of virus reduction by ultrafiltration with coagulation–sedimentation in water reclamation. *Food Environ Virol.* 2017;1–11. pmid:28455611
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)

30. Symonds EM, Verbyla ME, Lukasik JO, Kafle RC, Breitbart M, Mihelcic JR. A case study of enteric virus removal and insights into the associated risk of water reuse for two wastewater treatment pond systems in Bolivia. *Water Res.* 2014;65:257–70. pmid:25129566
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)

31. Schmitz BW, Kitajima M, Campillo ME, Gerba C, Pepper IL. Virus reduction during advanced bardenpho and conventional wastewater treatment process. *Environ Sci Technol.* 2016;50(17):9524–32. pmid:27447291
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)

32. Sangsanont J, The Dan D, Thi Viet Nga T, Katayama H, Furumai H. Detection of pepper mild mottle virus as an indicator for drinking water quality in Hanoi, Vietnam, in large volume of water after household treatment. *J Environ Sci Health Part A Toxic-Hazard Subst Environ Eng.* 2016;51(13):1100–6. pmid:27419274
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)

33. Verbyla ME, Symonds EM, Kafle RC, Cairns MR, Iriarte M, Mercado A, et al. Managing microbial risks from indirect wastewater reuse for irrigation in urbanizing watersheds. *Environ Sci Technol.* 2016;50(13):6803–13. pmid:26992352
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)

34. Lee H-W, Lee H-M, Yoon S-R, Kim SH, Ha J-H. Pretreatment with propidium monoazide/sodium lauroyl sarcosinate improves discrimination of infectious waterborne virus by RT-qPCR combined with magnetic separation. *Environ Pollut.* 2018;233(Supplement C):306–14. pmid:29096303
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)

35. Shrestha S, Shrestha S, Shindo J, Sherchand JB, Haramoto E. Virological quality of irrigation water sources and pepper mild mottle virus and tobacco mosaic virus as index of pathogenic virus contamination level. *Food Environ Virol.* 2018;10(1):107–20. pmid:29098656
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)