

2010

## Monitoring Coral Reefs from Space

C. M. Eakin

C. J. Nim

R. E. Brainard

C. Aubrecht

C. Elvidge

*See next page for additional authors*

Follow this and additional works at: [https://scholarcommons.usf.edu/msc\\_facpub](https://scholarcommons.usf.edu/msc_facpub)

---

### Scholar Commons Citation

Eakin, C. M.; Nim, C. J.; Brainard, R. E.; Aubrecht, C.; Elvidge, C.; Gledhill, D. K.; Muller-Karger, Frank E.; Mumby, P. J.; Skirving, W. J.; Strong, A. E.; Wang, M.; Weeks, S.; Wentz, F.; and Ziskin, D., "Monitoring Coral Reefs from Space" (2010). *Marine Science Faculty Publications*. 1086.

[https://scholarcommons.usf.edu/msc\\_facpub/1086](https://scholarcommons.usf.edu/msc_facpub/1086)

This Review is brought to you for free and open access by the College of Marine Science at Scholar Commons. It has been accepted for inclusion in Marine Science Faculty Publications by an authorized administrator of Scholar Commons. For more information, please contact [scholarcommons@usf.edu](mailto:scholarcommons@usf.edu).

---

**Authors**

C. M. Eakin, C. J. Nim, R. E. Brainard, C. Aubrecht, C. Elvidge, D. K. Gledhill, Frank E. Muller-Karger, P. J. Mumby, W. J. Skirving, A. E. Strong, M. Wang, S. Weeks, F. Wentz, and D. Ziskin

THE OFFICIAL MAGAZINE OF THE OCEANOGRAPHY SOCIETY

# Oceanography

#### CITATION

Eakin, C.M., C.J. Nim, R.E. Brainard, C. Aubrecht, C. Elvidge, D.K. Gledhill, F. Muller-Karger, P.J. Mumby, W.J. Skirving, A.E. Strong, M. Wang, S. Weeks, F. Wentz, and D. Ziskin. 2010. Monitoring coral reefs from space. *Oceanography* 23(4):118–133, doi:10.5670/oceanog.2010.10.

#### COPYRIGHT

This article has been published in *Oceanography*, Volume 23, Number 4, a quarterly journal of The Oceanography Society. Copyright 2010 by The Oceanography Society. All rights reserved.

#### USAGE

Permission is granted to copy this article for use in teaching and research. Republication, systematic reproduction, or collective redistribution of any portion of this article by photocopy machine, reposting, or other means is permitted only with the approval of The Oceanography Society. Send all correspondence to: [info@tos.org](mailto:info@tos.org) or The Oceanography Society, PO Box 1931, Rockville, MD 20849-1931, USA.

# Monitoring Coral Reefs from Space

BY C. MARK EAKIN, CARL J. NIM, RUSSELL E. BRAINARD,  
CHRISTOPH AUBRECHT, CHRIS ELVIDGE, DWIGHT K. GLEDHILL,  
FRANK MULLER-KARGER, PETER J. MUMBY,  
WILLIAM J. SKIRVING, ALAN E. STRONG, MENGHUA WANG,  
SCARLA WEEKS, FRANK WENTZ, AND DANIEL ZISKIN

**ABSTRACT.** Coral reefs are one of the world's most biologically diverse and productive ecosystems. However, these valuable resources are highly threatened by human activities. Satellite remotely sensed observations enhance our understanding of coral reefs and some of the threats facing them by providing global spatial and time-series data on reef habitats and the environmental conditions influencing them in near-real time. This review highlights many of the ways in which satellites are currently used to monitor coral reefs and their threats, and provides a look toward future needs and capabilities.

This nadir true-color image of Australia's Great Barrier Reef was acquired by the Multi-angle Imaging SpectroRadiometer (MISR) instrument on August 26, 2000, and shows part of the southern portion of the reef adjacent to the central Queensland coast. The width of the MISR swath is approximately 380 km, with the reef clearly visible up to approximately 200 km from the coast. *Image courtesy of NASA/GSFC/LaRC/JPL, MISR Team*

## INTRODUCTION

Coral reefs are one of the world's most biologically diverse and productive ecosystems (Porter and Tougas, 2001). They provide abundant ecological goods and services (Moberg and Folke, 1999) and are central to the socio-economic and cultural welfare of coastal and island communities throughout tropical and subtropical oceans by contributing at least \$30 billion (US\$) to the global economy when combined with tourism and recreation, shoreline protection, fisheries, and biodiversity services (UNEP-WCMC, 2006). Unfortunately, a range of human activities adversely impacts these valuable resources. Among the key threats are improper fishing activities, land-based sources of pollution, climate change, ocean acidification, and habitat destruction (Dodge et al., 2008; NOAA Coral Reef Conservation Program, 2009).

Satellites have the capacity to enhance our understanding of coral reef threats by obtaining global information on environmental conditions in near-real time and by providing spatial and time-series data relevant to management that are not practically obtained by in situ observations alone. This paper highlights the various ways remote-sensing data are being used to map and monitor coral reefs. It explains some remote-sensing tools commonly used to measure coral reef parameters of interest, how this information aids coral reef managers, some of the limitations of current technologies, and research gaps. Table 1 summarizes the coral reef parameters that we can currently measure by satellites. Mumby et al. (2004) and Andréfouët (in press) provide additional

reviews of satellite uses, methods, and applications for coral reefs, including discussions about using remote sensing to monitor key physical parameters that influence the conditions of coral reefs.

## MONITORING CORAL REEF ENVIRONMENTAL CONDITIONS

Coral reefs are exposed to a number of stressors from their surroundings. Although most satellites were not designed to observe coral reefs, many remote-sensing instruments provide valuable environmental data that are relevant to reef conditions. Calibration and validation data have been used to strengthen these data in shallow, near-shore environments, allowing coral reef managers and researchers to use this information to both understand the role of stressors and to better manage coral reef resources.

## Physical Parameters

With the growing consequences of global climate change, there is greater need for monitoring impacts in coral reef areas. Increasing anthropogenic concentrations of atmospheric CO<sub>2</sub> have numerous direct and indirect deleterious impacts on coral reefs. Carbon dioxide imparts an important control on the radiative heat balance of Earth's atmosphere, resulting in warming of both the atmosphere and the ocean (IPCC, 2007). Rising upper ocean temperatures, as indicated by sea surface temperature (SST), have increased the frequency and intensity of widespread thermal stress events that can cause mass coral bleaching (Eakin et al., in press), and this bleaching is expected to continue into the future (Hoegh-Guldberg et al., 2007, 2008). Rising SST can also result in an increase in infectious disease outbreaks

---

**C. Mark Eakin** ([mark.eakin@noaa.gov](mailto:mark.eakin@noaa.gov)) is Coordinator, National Oceanic and Atmospheric Administration (NOAA) Coral Reef Watch, Silver Spring, MD, USA. **Carl J. Nim** is NOAA Knauss Fellow, Coral Reef Watch, Silver Spring, MD, USA. **Russell E. Brainard** is Supervisory Oceanographer, NOAA Pacific Islands Fisheries Science Center, Coral Reef Ecosystem Division, Honolulu, HI, USA. **Christoph Aubrecht** is Research Scientist, Austrian Institute of Technology, Vienna, Austria. **Chris Elvidge** is Physical Scientist, NOAA National Environmental Satellite, Data, and Information Service (NESDIS), National Geophysical Data Center, Earth Observations Group, Boulder, CO, USA. **Dwight K. Gledhill** is Associate, NOAA Office of Oceanic and Atmospheric Research (OAR), Atlantic Oceanographic and Meteorological Laboratory, Silver Spring, MD, USA. **Frank Muller-Karger** is Professor, University of South Florida, Institute for Marine Remote Sensing, St. Petersburg, FL, USA. **Peter J. Mumby** is Professor, University of Queensland, School of Biological Sciences, Brisbane St. Lucia, Queensland, Australia. **William J. Skirving** is Contractor, NOAA Coral Reef Watch, Kirwan, Queensland, Australia. **Alan E. Strong** is Contractor, NOAA Coral Reef Watch, Silver Spring, MD, USA. **Menghua Wang** is Oceanographer, NOAA NESDIS Center for Satellite Applications and Research, Camp Springs, MD, USA. **Scarla Weeks** is Research Scientist, University of Queensland, Centre for Spatial Environmental Research, Brisbane St. Lucia, Queensland, Australia. **Frank Wentz** is Research Scientist and Proprietor, Remote Sensing Systems, Santa Rosa, CA, USA. **Daniel Ziskin** is Research Scientist, University of Colorado, Cooperative Institute for Research in Environmental Science, Boulder, CO, USA.

Table 1. Remote sensing platforms and sensors relevant to the monitoring of coral reefs and associated habitats. *Adapted from Mumby et al. (2004) and the Directory of Remote Sensing Applications for Coral Reef Management by the Remote Sensing Working Group of the Global Environment Facility Coral Reef Targeted Research & Capacity Building for Management Program*

Symbology: ✓ indicates well-established, ?✓ indicates fairly well-established, ? indicates experimental, ✓\* indicates data should be used in conjunction with acoustic sonar devices, blank indicates not currently possible.

PLATFORM		SHIP	AIRCRAFT		SATELLITE				
SENSOR TYPE		Acoustic	Imaging Spectrometers (Hyperspectral)	Laser	Microwave	Multispectral (High Spatial Resolution)	Hyperspectral (Medium Spatial Resolution)	Multispectral (Medium Spatial Resolution)	
EXAMPLE(S) OF PLATFORM OR SENSOR		RoxAnn, BioSonics	AVIRIS, CASI, ATM, HyMap	lidar, LADS	Aquarius, SMOS	IKONOS, QuickBird	EO-1 Hyperion	Landsat MSS/TM/ETM, SPOT, IRS	
APPLICATION	Reef location		✓	✓		✓	✓	✓	
	Mangroves		✓			✓	✓	✓	
	Sea grass beds		✓			✓	✓	✓	
	Reef geomorphology/habitats		✓	✓		✓	✓	✓	
	Reef community type		✓	✓		✓*			
	Beta (between habitat) diversity		✓			✓*			
	Connectivity of fish between mangroves and coral reefs							?	
	Coral cover (live vs. dead)			?					
	Reef structural complexity (rugosity)	?			?✓				
	Coral bleaching events			?		?			
	Bathymetry	✓			✓		✓	✓	
	Coral sensitivity to thermal stress								
	Wave exposure								
	Coral bleaching thermal stress								
	Coral disease risk								
	Physical model inversion methods			?			?		
	Sea surface temperature								
	Ultraviolet radiation								
	Photosynthetically active radiation						✓	✓	
	Light attenuation coefficients			✓			✓	✓	✓
	Cloud cover						✓	✓	✓
	Ocean sea level								
	Salinity					✓			
	Chlorophyll <i>a</i> concentration			✓			✓	✓	✓
	Algal blooms			✓				✓	✓
	Suspended sediment concentration			✓			✓	✓	✓
	Wind speed								
	Ocean circulation								
	Coastal circulation (feature tracking)								
	Precipitation								

Table continued on next page...

Acronyms: AVIRIS = Airborne Visible/Infrared Imaging Spectrometer, CASI = Compact Airborne Spectrographic Imager, ATM = Airborne Thematic Mapper, HyMap = Hyperspectral Mapper; lidar = light detection and ranging, LADS = Laser Airborne Depth Sounder, SLFMR = Scanning Low Frequency Microwave Radiometer, SMOS = Soil Moisture and Ocean Salinity, MSS = Multispectral Scanner, TM = Thematic Mapper, ETM = Enhanced Thematic Mapper, SPOT = Satellite Probatoire de l'Observations de la Terre, IRS = Indian Remote Sensing Satellite

Table 1, continued. Remote sensing platforms and sensors relevant to the monitoring of coral reefs and associated habitats. *Adapted from Mumby et al. (2004) and the Directory of Remote Sensing Applications for Coral Reef Management by the Remote Sensing Working Group of the Global Environment Facility Coral Reef Targeted Research & Capacity Building for Management Program*

Symbology: ✓ indicates well-established, ?✓ indicates fairly well-established, ? indicates experimental, ✓\* indicates data should be used in conjunction with acoustic sonar devices, blank indicates not currently possible.

PLATFORM		SATELLITE					
SENSOR TYPE		Multispectral (Low Spatial Resolution)	Meteorological	Radar Scatterometer	Synthetic Aperture Radar	Radar Altimeter	Radiometer
EXAMPLE(S) OF PLATFORM OR SENSOR		SeaWiFS, MODIS, Envisat MERIS	GOES, GMS, Meteosat	ASCAT	ASAR	Jason-1/2	POES, TRMM, AVHRR, ATSR
APPLICATION	Reef location	✓					
	Mangroves	✓					
	Sea grass beds	✓					
	Reef geomorphology/habitats						
	Reef community type						
	Beta (between habitat) diversity						
	Connectivity of fish between mangroves and coral reefs						
	Coral cover (live vs. dead)						
	Reef structural complexity (rugosity)						
	Coral bleaching events						
	Bathymetry						
	Coral sensitivity to thermal stress		?				?
	Wave exposure				?	?	?
	Coral bleaching thermal stress						✓
	Coral disease risk						?
	Physical model inversion methods						
	Sea surface temperature	✓(MODIS)	✓				✓
	Ultraviolet radiation		✓				✓
	Photosynthetically active radiation	✓	✓				✓
	Light attenuation coefficients	✓					✓
	Cloud cover	✓	✓				✓
	Ocean sea level					✓	
	Salinity						
	Chlorophyll <i>a</i> concentration	✓					
	Algal blooms	✓					
	Suspended sediment concentration	✓					✓
Wind speed				✓		✓	
Ocean circulation	✓	✓			✓	✓	
Coastal circulation (feature tracking)	✓	✓				?✓	
Precipitation					✓		

Acronyms: SeaWiFS = Sea-viewing Wide Field-of-view Sensor, MODIS = Moderate Resolution Imaging Spectroradiometer, Envisat MERIS = Environmental Satellite - Medium Resolution Imaging Spectrometer, GOES = Geostationary Operational Environmental Satellite, GMS = Geosynchronous Meteorological Satellite, Meteosat = Meteorological Satellite, ASCAT = Advanced Scatterometer, ASAR = Advanced Synthetic Aperture Radar, POES = Polar Operational Environmental Satellite, TRMM = Tropical Rainfall Measuring Mission, AVHRR = Advanced Very High Resolution Radiometer, ATSR = Along Track Scanning Radiometer

(Bruno et al., 2007). The following section explains how satellite observations are used to monitor a variety of climate-related physical parameters, such as SST, solar radiation, and wind.

### Thermal Stress and Coral Bleaching

Polar-orbiting environmental satellites (POES) provide near-real time observations and have measured SST at 4-km spatial resolution since 1981. The US National Oceanic and Atmospheric Administration's (NOAA's) Coral Reef Watch (CRW) provides a suite of operational global satellite coral bleaching monitoring products at 0.5° latitude-longitude spatial resolution, produced twice weekly (Liu et al., 2006)

from nighttime SSTs (Figure 1a). Over 20 years of NOAA SST data have been reprocessed to produce a consistent data set at 4-km spatial resolution (Kilpatrick et al., 2001). Analysis of the 22-year Pathfinder data set suggests that the water temperatures of most coral reef areas have been increasing at 0.2–0.4°C per decade (Strong et al., 2009). This warming agrees with studies that indicate corals may need to acclimate their thermal tolerance by 0.2–1.0°C per decade to survive repeated coral bleaching events predicted under future climate scenarios (Donner et al., 2005).

The CRW coral bleaching HotSpot, released in 1996 (Strong et al., 1997) and made operational in 2002 (Liu

et al., 2003), was the first coral-specific product developed by NOAA's National Environmental Satellite, Data, and Information Service (NESDIS). Based on the "ocean hot spots" concept introduced by Goreau and Hayes (1994), HotSpots are positive temperature anomalies that exceed the maximum monthly mean (MMM) SST climatology for each 0.5° pixel, thereby identifying currently thermally stressed regions. These HotSpots are accumulated over a moving 12-week window to produce CRW's Degree Heating Week (DHW) index that is highly predictive of bleaching occurrence and severity. Significant coral bleaching is expected to occur one to three weeks after reefs begin




Figure 1. NOAA Coral Reef Watch near-real time satellite global 50-km nighttime product suite for February 9, 2009: (A) sea surface temperatures (SST), (B) coral bleaching alert area product that combines HotSpot and Degree Heating Week products.

to experience DHW values  $\geq 4^{\circ}\text{C}\text{-weeks}$ , and mass bleaching and the onset of coral mortality is expected after DHW values  $\geq 8^{\circ}\text{C}\text{-weeks}$ . CRW later combined the HotSpot and DHW data to produce the Coral Bleaching Alert Area product (Figure 1b)—a singular product, of particular value to managers, that displays areas where bleaching thermal stress currently is expected. In the future, HotSpot and DHW algorithms at 1–4 km resolutions from NOAA and National Aeronautics and Space Administration (NASA) Moderate Resolution Imaging Spectroradiometer (MODIS) sensors (Hu et al., 2009) will be used to evaluate relationships between SST high-resolution spatial structure and coral reef ecosystem responses (Inia Soto, University of South Florida, *pers. comm.*, September 27, 2010).

CRW's "Virtual Stations" also provide time series SST data for over 190 selected reef sites around the world that can be used by coastal resource managers for predicting coral bleaching. Though based on satellite data, Virtual Stations are somewhat analogous to having broad-scale sensors of the surface waters surrounding a reef. CRW uses Virtual Stations to provide automated bleaching alert e-mails that notify subscribers when coral reefs are experiencing conditions conducive to bleaching. These bleaching alerts provide opportunities for local reef managers and researchers to visually assess coral conditions and take management actions if local conditions warrant them.

The spatial resolution of these data and the variability of oceanographic conditions in these areas require calibration and validation of remote-sensing data to improve analyses. Many SST products are calibrated using offshore,




Figure 2. Prototype of the new NOAA Light Stress Damage product combining thermal and light stress as a bleaching predictor.

rather than nearshore, SST observations (Emery et al., 2001). SST data, especially new high-resolution products, need to be verified for use in coral reef environments. Van Hooidonk and Huber (2009) used a meteorological forecast verification method to quantitatively assess the quality of the derived DHW product. Field observations of corals have also been used to verify predicted bleaching events and determine if large-scale high temperatures or local stressors were the cause (McClanahan et al., 2007). Detailed comparisons of satellite remote sensing to actual community changes will permit the development of models with strong management application, such as bleaching susceptibility models for identifying resilient reef areas (Maina et al., 2008).

#### Solar Radiation and Coral Bleaching

A variety of projects over the past decade have examined the use of geostationary satellite data to study aspects of solar radiation over coral reefs (Tovar and Baldasano, 2001; Hansen et al., 2002; Kandirmaz et al., 2004). In 2009, NOAA developed a suite of experimental surface solar radiation products based on data from Geostationary Operational Environmental Satellite (GOES) systems

that were calibrated and validated to provide insolation estimates over oceanic waters (<http://www.osdpd.noaa.gov/ml/land/gsip>). These products measure total daily global ocean surface insolation. These data have allowed CRW and its partners to develop a new set of coral bleaching products that endeavour to provide a combined measure of satellite-derived thermal stress and light as an index of stress on coral photosystems. Figure 2 shows a mock-up of the product that CRW is developing.

#### Bleaching Weather: The Influence of Wind

Local weather patterns shape the potential for coral bleaching. In particular, wind influences local bleaching patterns by affecting temperature, light, water-column mixing, and sediments. As wind speeds decrease, turbulent vertical mixing and evaporative cooling are reduced, sensible heat loss increases, and the likelihood of anomalously high temperatures and light penetration increases (Skirving and Guinotte, 2001; Obura, 2005). Additionally, low winds increase water-column stratification, increase light penetration by enhancing sediment settlement, and facilitate photo-degradation of colored dissolved




Figure 3. NOAA/NESDIS Coral Reef Watch near-real-time satellite 25-km Doldrums experimental product for April 22, 2007, in the Indian Ocean region. The color scale indicates the number of days over which the daily mean National Climatic Data Center Blended Sea Winds remained below  $3 \text{ m s}^{-1}$ .

organic material (Manzello et al., 2006). CRW has developed an experimental Doldrums product using NOAA's National Climatic Data Center (NCDC) Blended Sea Winds Product (Zhang et al., 2006). The experimental Doldrums product identifies regions of mean wind speed below  $3 \text{ m s}^{-1}$  and records the persistence (days) of such conditions (Figure 3; CRW 2007; <http://coralreef-watch.noaa.gov/satellite/doldrums>).

Analyses of long-term patterns of winds may provide additional insights into conditions around coral reefs. A recent analysis of satellite Special Sensor Microwave Imager (SSM/I) observations suggests that precipitation and total atmospheric water have increased equally at a rate of  $\sim 1\%$  per decade over the past two decades (Wentz et al., 2007). A least-squares linear fit of SSM/I wind speed for each  $2.5^\circ$  grid cell was calculated after removing the seasonal variability to provide a decadal trend map of wind speed (Figure 4) and compared with wind trends from the International Comprehensive Ocean-Atmosphere Data Set (ICOADS) from

ship-based observations. Winds averaged from  $30^\circ\text{S}$  to  $30^\circ\text{N}$  increased by  $0.04 \text{ m s}^{-1}$  ( $0.6\%$  per decade), resulting in wet areas becoming wetter (Wentz et al., 2007). Wentz et al. (2007) suggest that two decades may be too short for extrapolating these short-term trends into longer ones; however, continuation of these trends could significantly impact circulation, mixing, and temperature regulation of waters around reefs in the Indo-West Pacific, the heart of coral biodiversity and abundance.

### Chemical Parameters

Though corals predominantly occur in clear, oligotrophic waters, nutrient-rich runoff from land-based sources of pollution (LBSP) often results in algal overgrowth of corals and coral recruits (Adey, 2000). Phosphate pollution can inhibit calcium carbonate ( $\text{CaCO}_3$ ) deposition, slowing coral growth (Muller-Parker and D'Elia, 1997). Increased nutrients from LBSP influence trophic interactions on coral reefs, including phytoplankton blooms that may smother corals (Guzmán et al.,

1990) or enhance recruitment of larval *Acanthaster planci* (crown-of-thorns sea stars) that feed on corals (Brodie, 2005). Suspended sedimentation causes coral reef degradation (Rogers, 1990; Field et al., 2008) by damaging and smothering corals, thus restricting their growth (Fabricius, 2005). Sediment plumes and phytoplankton blooms are recognizable by satellite ocean color sensors. Unfortunately, various factors, including coastal aerosols, mixed signals from the water column, and heterogeneous benthic substrates, make quantitative monitoring of ocean color parameters a challenge in coral reef environments (IOCCG, 2000).

### Land-Based Sources of Pollution

Many coral reefs are naturally exposed to the influence of nutrients and sediment from both large and small rivers (Muller-Karger and Castro, 1994; Soto et al., 2009) and from nonpoint runoff. LBSP are of growing concern as land and coastal zones are developed. LBSP include a wide variety of materials that individually or collectively threaten coral

reefs' health. Of primary need for monitoring the coral reef environment are remotely sensed observations of rainfall events and their associated runoff, which often lead to reduced salinity, increased turbidity and sedimentation, and observations of transported nutrients that contribute to phytoplankton blooms.

Remote sensing of coastal water-quality parameters (e.g., suspended sediment and chlorophyll *a*) using visible radiance is difficult with ocean color sensors due to their technical design, but advances in remote-sensing science have provided some innovative solutions (Muller-Karger et al., 2005). Many global, open-water ocean color products derived from moderate spatial resolution satellite ocean color sensors (e.g., SeaWiFS [Sea-viewing Wide Field-of-view Sensor, MODIS) employ atmospheric correction algorithms that use the near-infrared (NIR) bands (Gordon and Wang, 1994;

IOCCG, 2010). The complexity of turbid, coastal areas prompted researchers to develop a shortwave infrared (SWIR) atmospheric correction algorithm (Wang, 2007) to improve the accuracy of ocean color products in these waters (Wang et al., 2009). One avenue of recent work has enhanced preprocessing techniques to derive coastal coral reef water-quality parameters by combining deep-water algorithms (Lee et al., 2002; Qin et al., 2007) with shallow-water mapping schemes. Although ocean color advances have been made for deriving coastal water-quality parameters in coral reef ecosystems (Wang et al., 2009), many challenges remain.

#### Ocean Acidification

As human activities continue to increase carbon dioxide in the atmosphere ( $CO_{2,atm}$ ), much of this  $CO_2$  is absorbed into surface ocean waters. Surface

ocean  $CO_2$  ( $CO_{2,aq}$ ) then reacts with water to form carbonic acid, thereby lowering pH; hence, the term “ocean acidification” (Caldeira and Wickett, 2003). This process also consumes carbonate ions, reducing the degree of saturation ( $\Omega_{sp}$ ) of seawater with respect to calcium carbonate. This may result in reduced coral growth rates and compromise the structural integrity of coral reefs (Manzello et al., 2008; Veron, 2008; Gledhill et al., 2009). Many experimental studies have suggested a relationship between declining seawater saturation and the rate at which many marine organisms (including many coral species) produce calcium carbonate, albeit the relationship can prove complex when interactions with temperature and nutrients are considered (for a review see Doney et al., 2009). Monitoring ocean acidification, particularly secular declines in  $\Omega_{arg}$  in response to rising




Figure 4. Surface-wind trends for the period July 1987 through August 2006 computed at a spatial resolution of 2.5°. (A) Special Sensor Microwave Imager (SSM/I) satellite wind trends. (B) International Comprehensive Ocean-Atmosphere Data Set (ICOADS) ship wind trends. In the North Pacific and North Atlantic where ICOADS ship observations are more abundant, the two data sets show similar trends. From Wentz et al. (2007)

atmospheric  $\text{CO}_{2,\text{atm}}$ , could yield insights into the expected regional responses of coral reef ecosystems.

In situ measurements of ocean chemistry provide the most accurate means of tracking ocean acidification, but such measurements are inherently limited in space (repeat sampling, moored stations) and/or time (ship surveys). Although current satellites cannot directly measure changes in ocean carbonate chemistry, they can provide synoptic observations of a range of physical and optical parameters that allow us to model these changes. In 2008, CRW and NOAA's Atlantic Oceanographic and Meteorological Laboratory (AOML) developed and released an experimental Ocean Acidification Product Suite (OAPS) (<http://coralreefwatch.noaa.gov/satellite/oa>). The product provides a monthly  $0.25^\circ \times 0.25^\circ$  analysis of modeled sea surface carbonate chemistry ( $\Omega_{\text{arg}}$ ,  $p\text{CO}_{2(\text{sw})}$ , total alkalinity, carbonate ion, and bicarbonate ion) within the oceanic waters of the Greater Caribbean Region (Gledhill et al., 2008), based on a set of regionally specific algorithms, satellite remote-sensing data, and shipboard in situ observations.

The OAPS algorithms merge multiple environmental data sets, including  $\text{CO}_{2,\text{atm}}$  concentrations, sea-level air pressure, temperature, and relative humidity. The dominant controls on surface  $\Omega_{\text{arg}}$  dynamics within these oligotrophic waters are  $p\text{CO}_{2(\text{sw})}$  availability, SST, and salinity. Typically, SST imparts the greatest subannual control on  $\text{CO}_2$  uptake within these oligotrophic waters, for which OAPS uses the currently available  $0.25^\circ$  daily NOAA optimal interpolated (OI) SST product, which depends upon remotely sensed radiometric SST data from in

situ, Advanced Very High Resolution Radiometer (AVHRR), and Advanced Microwave Scanning Radiometer - Earth Observing System (AMSR-E) sensors (Reynolds et al., 2007). While it is hoped that salinity dependency will be partially met through the application of Aquarius satellite sea surface salinity data sets in coming years, the current model relies upon Hybrid Coordinate Ocean Model (HYCOM) + Navy Coupled Ocean Data Assimilation (NCODA)  $1/12^\circ$  Analysis GLBa0.08 salinity fields. Given the coarse spatial and temporal resolution of Aquarius, it is likely that even future OAPS iterations will retain some dependence on HYCOM-produced salinity fields.

By solving for both surface alkalinity and  $p\text{CO}_{2,\text{sw}}$  it is possible to fully describe the carbonic acid system (that is, solve the distribution of the various carbonate species), permitting OAPS to solve for  $\Omega_{\text{arg}}$ . More-sophisticated algorithms are currently under development that would provide for a more general application of OAPS to other tropical regions through the addition of other variables (e.g., winds, chlorophyll *a*).

### Salinity

Sea surface salinity directly influences seawater density and chemical processes such as carbonate chemistry. Until recently, salinity was measured in situ or by airborne microwave radiometers to assess the role of riverine plumes on seagrass and coral reefs (Klemas, 2009). With the launch of the European Soil Moisture and Ocean Salinity (SMOS) satellite in 2009, and NASA's Aquarius satellite in 2010, monthly global averages of sea surface salinity now can be remotely measured with an accuracy of 0.1 and 0.2 psu at 200-km and

150-km spatial resolution, respectively (Robinson, 2004; Martin, 2004; Lagerloef et al., 2008). As mentioned above, salinity at this resolution may be helpful in driving models to provide downscaled salinity to improve our monitoring of alkalinity and, thus, ocean carbon chemistry. It may also help provide information on increased runoff from large rivers as well as production of warm hypersaline waters over extensive shallow banks.

### Coastal Development

Although current satellites cannot identify complex chemical pollutants, it is possible to use satellite observations of human activity to provide a proxy for these other stressors. Understanding the extent of urbanization and human activity can provide a proxy for localized impacts such as pollutants, runoff, fishing, and recreational use of reefs. A prominent indicator of human occupation visible from space is artificial night lighting. Observable features have included biomass burning, massive offshore fisheries, and infrastructure lights. NOAA has processed nighttime lights data acquired by the US Air Force Defense Meteorological Satellite Program (DMSP) Operational Linescan System (OLS) (Aubrecht et al., 2008) in order to integrate the brightness and distance of lights near known coral reef sites as a proxy for human development stresses. Light proximity index (LPI) values have been calculated for the three stressors observable in nighttime lights: human infrastructure, gas flares, and heavily lit fishing boats. The algorithm computes the contribution of lights declining with increased distance from reefs. LPI since 1992 shows increased human settlements near coral reefs




Figure 5. Light proximity index (LPI) percent change over time (1992–2009) from human settlements, lit fishing boats, and offshore gas flares for the reefs of the world. LPI values are normalized by absolute number of reef points to show percent change over time. Colors correspond with stressors and point shapes with the satellite used.

(Figure 5), a reflection of the expanding populations and infrastructural development seen in coastal areas in many parts of the world. In contrast, lit fishing boats activity has declined and gas flares near reefs show a more complex pattern with dips in 1994 and 2001, and a peak in 1997. LPI can provide a valuable tool for coastal management, especially in areas where other data on coastal development may be limited.

### MANAGEMENT APPLICATIONS

Measurement of environmental parameters is helpful to coral reef managers because it alerts them of conditions when coral reefs in their jurisdictions are in need of direct monitoring or management action. There are times, however, when in situ data are not sufficient for monitoring, or reef locations are too remote to gather in situ data. In these cases, high-resolution remote-sensing

data can provide researchers and resource managers with needed data. The availability of high-resolution satellite data has increased in recent years due to launching of more advanced sensors, application of emerging technologies in sensor designs, and new algorithms. These data are used to study reef carbonate dynamics (Moses et al., 2009), bathymetry in coral reef areas (Hogrefe et al., 2008), geomorphology and biodiversity of coral reefs (Andréfouët and Guzman, 2005; Knudby et al., 2007), and coral reef fish species richness, communities, and fisheries (Mellin et al., 2009; Hamel and Andréfouët, 2010). They also allowed managers to more effectively integrate in situ and remotely sensed data (Scopélitis et al., 2010). Given the breadth of literature on this topic, this section will only provide a survey of some important applications, including marine protected area (MPA)

management, habitat characterization, and some important biological parameters of coral reefs.

### The Use of Remote Sensing for Marine Protected Areas

Analyses of coral reefs using high-spatial-resolution (30-m pixel size or less) data span a broad scale of applications, including MPA design and evaluation (Green et al., 2009; Dalleau et al., 2010), study of ecosystem associations, (e.g., coral reefs with seagrass beds and mangroves), and investigations of the ecology of coral reefs and the organisms that live in them (e.g., fish). Although the bulk of these projects are conducted at local scales, there are calls to automate these techniques for use at broader scales (Andréfouët, 2008). NOAA’s analysis of MPA characteristics and design in the northwestern Hawaiian Islands and in Puerto Rico has used remotely sensed data in conjunction with in situ measurements of coral habitat and spectral reflectance characteristics to produce benthic habitat maps for shallow-water coral reefs. These baseline habitat maps provide resource managers and researchers with essential information for planning MPAs, monitoring changes, and evaluating MPA effectiveness in both the Pacific (Friedlander et al., 2007, 2008) and the Caribbean (García-Sais et al., 2008).

### Habitat Characterization

Mapping has proven to be a valuable tool for understanding the interconnections between coral reefs and associated habitats, such as the role mangroves play as juvenile reef fish nurseries (Mumby, 2006). Such ecosystem-based perspectives can aid the designation and management of MPAs.

High-spatial-resolution mapping of mangrove and seagrass species has been explored (Myint et al., 2008; Phinn et al., 2008) and regional seagrass mapping using Landsat data has been assessed for broader applications across the Caribbean (Wabnitz et al., 2008).

A wide variety of remote-sensing platforms and algorithms are available for habitat mapping. A diverse literature exists on habitat mapping using satellite sensors such as Satellite Pour l'Observation de la Terre (SPOT) and Landsat Thematic Mapper (TM) at a spatial resolution of tens of meters (LeDrew et al., 2000), and on using high-spatial-resolution sensors such as the commercial Earth-observation satellites IKONOS and QuickBird (Mumby and Edwards, 2002; Andréfouët et al., 2003; Mishra et al., 2006). Even greater accuracies can be obtained by including data from airborne sensors like the Compact Airborne Spectrographic Imager (CASI) either as the prime data source (Mumby et al., 1998; Bertels et al., 2008) or in combination with satellite data (Rowlands et al., 2008).

Landsat data to identify geomorphological characteristics of coral reefs on a global scale. Data obtained in this effort have already informed numerous studies on strategic MPA placement, reef condition assessments (Burke and Maidens, 2004), and both climate forcing of reef growth on the Great Barrier Reef and climate change influences on biogeochemical budgets in French Polynesian atolls (Andréfouët et al., 2006).

Seafloor bathymetry and benthic rugosity are important variables to coral reef researchers and managers alike. Pseudo-bathymetry has been derived from IKONOS data in a number of studies (Lyzenga et al., 2006), modified by applying nonlinear inversion models (Su et al., 2008), and used to fill the gap between terrestrial digital elevation models (DEMs) and sonar-acquired bathymetry (Hogrefe et al., 2008). In other studies using light detection and ranging (lidar), rugosity maps were developed to illustrate massive stony coral colonies on patch reefs (Brock et al., 2006) and to investigate statistical relationships between rugosity and reef

(Pittman et al., 2007; Knudby et al., 2008; Knudby et al., 2010). Mellin et al. (2009) advocated a “hierarchy of habitat” for predicting coral reef fish habitats using geomorphology, benthic assemblages, rugosity, and depth as key habitat variables.

### Biological Parameters of Coral Reef Ecosystems

Coral reef cover is a parameter of great interest to researchers and managers. To date, there have been three approaches to quantifying changes in the benthic composition of coral reefs using coral and macroalgae cover as key indicators of reef health. A number of scientists have used time-series data to detect changes in overall reflectance that can be attributed to major changes in benthic state. For example, Dustan et al. (2001) analyzed a Landsat time series from Florida and found a change in “temporal texture” associated with the die-off of long-spined sea urchins. These indirect methods are useful, but the data have been difficult to interpret because multiple changes in benthic characteristics may have similar effects. A second approach uses high-resolution optical data to measure changes in reef spectra. Researchers either interpret these data directly (Lesser and Mobley, 2007) or extract characteristic spectral derivatives from them (Holden and LeDrew, 1998) to classify live and recently dead coral (Mumby et al., 2004, 2001). For these studies, it is important that the reefs have a clear-water environment and are devoid of brown macroalgae, which are frequently spectrally confused with coral (Hedley and Mumby, 2002). The wider efficacy of these methods for evaluating coral cover is still under investigation (Hedley et al., 2004; Joyce et al., 2004).

## “SATELLITE TECHNOLOGIES HAVE BECOME ESSENTIAL TOOLS FOR MONITORING CORAL REEF HEALTH AND THE INCREASING THREATS CORAL REEFS FACE AROUND THE GLOBE.”

Perhaps the most ambitious project to characterize the habitats of coral reefs worldwide was the Millennium Coral Reef Mapping Project (MCRMP) (Andréfouët et al., 2002), which used

fish communities (Kuffner et al., 2007). Other studies combined bathymetry with habitat maps derived from remote-sensing satellites to drive models that predict coral reef fish distribution

Figure 6. Image differencing, or spectral brightening, of Nikunau Island, Kiribati. The difference image (far right) shows a gold color where presumably bleaching has occurred.



A third approach to mapping corals has used sonar to discriminate branching corals, such as *Acropora* spp., from other reef structures (Purkis et al., 2006).

Mass coral bleaching has been successfully detected using remotely sensed observations in cases where the extent of bleaching is pronounced in high-coral-cover reef habitats; Figure 6). High spatial resolution is key to reliable remote detection of coral bleaching (Andréfouët et al., 2002). Elvidge et al. (2004) developed a method for detecting the brightening of reef areas when corals bleach using pairs of high-resolution multispectral images. The technique involves detection of spectral brightening observed when comparing satellite images acquired before the bleaching event with images collected during the bleaching event. This technique has continued to be refined and applied to multiple satellite sensors (Daniel Ziskin, University of Colorado at Boulder, *pers. comm.*, September 27, 2010). However, images like those in Figure 7 only provide a general index of reef bleaching and cannot separate bleaching of corals from bleaching of crustose coralline algae in reef systems, limiting their value for reefs where the community structure is poorly known. Additionally, as long as

these approaches require the purchase of costly satellite imagery, they will only be practical for remote areas that cannot be reached by divers.

### THE FUTURE OF MONITORING CORAL REEFS FROM SPACE

Improvements in three major areas are increasing the potential for using remotely sensed data to monitor coral reefs: resolution, spectral bands, and algorithms. Whether it is remote sensing of stressors, environmental parameters, or habitats and their changes, increased spatial and temporal resolution provide new ways that coral reefs can be monitored. Unfortunately, there is often a trade-off between spatial and temporal resolutions, as finer-resolution imagery is frequently obtained through narrower swaths and a corresponding increase in time between repeated observations. However, improved technology, such as off-nadir viewing, and image processing are helping to resolve this issue. Higher-resolution applications also require a larger data set of in situ observations for calibrating and validating changes, especially in highly variable nearshore environments.

At the same time, a greater portion of the electromagnetic spectrum is

being used in remote sensing. Enhanced optical systems, such as multispectral to hyperspectral sensors covering more of the visible and nearby bands, will be used to detect changes in benthic habitats and will do a better job of separating benthic features from water-column properties. Because coral reefs normally exist in shallow, clear waters, many optical bands can “see” the bottom. This attribute is good for habitat measurements, but it creates problems for separating changes in the benthos from changes in the water column. Both enhanced spatial and spectral resolution are essential to resolving these problems and will provide the quantitative data needed to answer many resource-monitoring challenges that currently exist for coral reef ecosystems.


None of these improvements are sufficient, however, without considerable work to develop, calibrate, and validate new and improved algorithms. Because satellites can only measure a small set of parameters directly, most of what we monitor from satellites is the result of complex algorithms that derive the parameters of interest. This is also where the combination of multiple instruments and spectral bands show great promise in enhancing our

## A CLASSROOM ACTIVITY USING SATELLITE SEA SURFACE TEMPERATURES TO PREDICT CORAL BLEACHING

Don't miss the related Hands-On Oceanography activity from Oceanography 22-2. This activity illustrates how temperature influences coral bleaching and how remote sensing is used to monitor coral health. It can be presented with different levels of complexity to upper middle school through early college students.

Download the full activity from

[HTTP://WWW.TOS.ORG/HANDS-ON](http://www.tos.org/hands-on)



remote-sensing capabilities.

Although satellite remote sensing is already an essential tool for assisting global efforts to map and monitor coral reefs, as described above, many opportunities and challenges remain. Will future technological advances enable us to monitor the abundance and distribution of coral reef fish, near-real-time changes in benthic habitats, indices of coral reef biodiversity, or other indicators of ecosystem health and ecological responses to climate change and ocean acidification? Where else do we need to stretch the envelope of remote-sensing capabilities to best address these challenges? Resource managers are asking when scientists will provide tools to alert them when changes occur in the cover of seagrass on the ocean bottom. They would like to see hypoxic or “black water” events that kill fish. These

managers would also like to know when changes in water quality cause harmful algal blooms or the spread of bacterial mats across the seafloor. They would like to know when diseases are spreading through ecosystems. Some of these capabilities are likely to be available over time scales of a few years while others will likely take decades and require significant advances in the development of new sensors and algorithms.

Some of these needs will be answered by advancing the use of remotely sensed and in situ data to initiate and support numerical models. These models estimate environmental parameters (as was done for ocean acidification) or impacts, such as changes in community composition. In many cases, these models integrate multiple data sets to provide new estimates of unmeasured parameters. These estimates may serve as solutions to

scientific questions that inform management needs or may be a bridge until new sensors are available. The development and launch of new satellite sensors is a slow process, and agencies involved need to better incorporate the needs of resource managers into their instrument and satellite development processes. At the same time, we need to identify key parameters that must be collected with high quality and continuously to understand long-term changes to coral reefs and the water quality and oceanographic conditions influencing them.

Coral reefs are important and valuable ecosystems. Satellite technologies have become essential tools for monitoring coral reef health and the increasing threats coral reefs face around the globe. We need to continue to advance the tools available and the science behind them in order to make these tools as useful as possible. As the anthropogenic threats to coral reefs continue to mount, we need to continue our diligent use of all tools at our disposal to keep these valuable resources healthy.

### ACKNOWLEDGEMENTS

Thanks to the many colleagues and collaborators who have contributed to the development of satellite remote-sensing technology and its application to coral reef ecosystems. In particular, we thank Serge Andréfouët and additional reviewers for their thoughtful evaluation of manuscripts. We are also grateful to the funding agencies and organizations that have supported the work, including the National Oceanic and Atmospheric Administration, the National Aeronautics and Space Administration, the Australian Research Council, and the World Bank/Global Environment Facility. The manuscript contents are

solely the opinions of the authors and do not constitute a statement of policy, decision, or position on behalf of NOAA or the US government. 

## REFERENCES

- Adey, W.H., T.A. McConnaughey, A.M. Small, and D.M. Spoon. 2000. Coral reefs: Endangered, biodiverse, genetic resources. Pp. 33–42 in *Seas at the Millennium*. C.R.C. Sheppard, ed., Elsevier.
- Andréfouët, S. 2008. Coral reef habitat mapping using remote sensing: A user vs producer perspective: Implications for research, management and capacity building. *Spatial Science* 53(1):113–129.
- Andréfouët, S. In press. Remote sensing. Chapter in *Encyclopedia of Modern Coral Reefs: Structure, Form and Process*. D. Hopley, ed., Springer-Verlag.
- Andréfouët, S., and H.M. Guzman. 2005. Coral reef distribution, status and geomorphology: Biodiversity relationship in Kuna Yala (San Blas) archipelago, Caribbean Panama. *Coral Reefs* 24(1):31–42.
- Andréfouët, S., R. Berkemans, L. Odriozola, T. Done, J. Oliver, and F. Muller-Karger. 2002. Choosing the appropriate spatial resolution for monitoring coral bleaching events using remote sensing. *Coral Reefs* 21(2):147–154.
- Andréfouët, S., P. Kramer, D. Torres-Pulliza, K.E. Joyce, E.J. Hochberg, R. Garza-Perez, P.J. Mumby, B. Riegl, H. Yamano, W.H. White, and others. 2003. Multi-site evaluation of IKONOS data for classification of tropical coral reef environments. *Remote Sensing of Environment* 88(1–2):128–143.
- Andréfouët, S., F.E. Muller-Karger, J.A. Robinson, C.J. Kranenburg, D. Torres-Pulliza, S.A. Spraggins, and B. Murch. 2006. Global assessment of modern coral reef extent and diversity for regional science and management applications: A view from space. Pp. 1,732–1,745 in *Proceedings of 10<sup>th</sup> International Coral Reef Symposium*. Okinawa, Japan, June 28–July 2, 2004. Japanese Coral Reef Society, Tokyo, Japan. Available online at: [http://www.reefbase.org/resource\\_center/publication/icrs.aspx?icrs=ICRS10](http://www.reefbase.org/resource_center/publication/icrs.aspx?icrs=ICRS10) (accessed September 22, 2010).
- Aubrecht, C., C.D. Elvidge, T. Longcore, C. Rich, J. Safran, A.E. Strong, C.M. Eakin, K.E. Baugh, B.T. Tuttle, A.T. Howard, and E.H. Erwin. 2008. A global inventory of coral reef stressors based on satellite observed nighttime lights. *Geocarto International* 23(6):467–479.
- Bertels, L., T. Vanderstraete, S. Van Coillie, E. Knaeps, S. Sterckx, R. Goossens, and B. Deronde. 2008. Mapping of coral reefs using hyperspectral CASI data: A case study: Fordata, Tanimbar, Indonesia. *International Journal of Remote Sensing* 29(8):2,359–2,391.
- Brock, J.C., C.W. Wright, I.B. Kuffner, R. Hernandez, and P. Thompson. 2006. Airborne lidar sensing of massive stony coral colonies on patch reefs in the northern Florida reef tract. *Remote Sensing of Environment* 104:31–42.
- Brodie, J., K. Fabricius, G. De'ath, and K. Okaji. 2005. Are increased nutrient inputs responsible for more outbreaks of Crown of Thorns starfish? An appraisal of the evidence. *Marine Pollution Bulletin* 51:266–278.
- Bruno, J.F., E.R. Selig, K.S. Casey, C.A. Page, B.L. Willis, C.D. Harvell, H. Sweatman, and A.M. Melendy. 2007. Thermal stress and coral cover as drivers of coral disease outbreaks. *PLoS Biology* 5(6):e124.
- Burke, L., and J. Maidens. 2004. *Reefs at Risk in the Caribbean*. World Resources Institute, Washington, DC, 80 pp. Available online at: <http://www.wri.org/publication/reefs-risk-caribbean> (accessed September 22, 2010).
- Caldeira, K., and M.E. Wickett. 2003. Anthropogenic carbon and ocean pH. *Nature* 425:365, doi:10.1038/425365a.
- Dalleau, M., S. Andréfouët, C.C. Wabnitz, C. Payri, L. Wantiez, M. Pichon, K. Friedman, L. Vigliola, and F. Benzoni. 2010. Use of habitats as surrogates of biodiversity for efficient coral reef conservation planning in Pacific Ocean islands. *Conservation Biology* 24(2):541–552.
- Dodge, R.E., C. Birkeland, M. Hatziozols, J.A. Kleypas, S.R. Palumbi, O. Hoegh-Guldberg, R. Van Woessik, J.C. Ogden, R.B. Aronson, B.D. Causey, and F. Staub. 2008. A call to action for coral reefs. *Science* 322:189–190.
- Doney, S.C., V.J. Fabry, R.A. Feely, and J.A. Kleypas. 2009. Ocean acidification: The other CO<sub>2</sub> problem. *Annual Review of Marine Science* 1:169–192.
- Donner, S.D., W.J. Skirving, C.M. Little, M. Oppenheimer, and O. Hoegh-Guldberg. 2005. Global assessment of coral bleaching and required rates of adaptation under climate change. *Global Change Biology* 11(12):2,251–2,265.
- Dustan, P., E. Dobson, and G. Nelson. 2001. Landsat Thematic Mapper: Detection of shifts in community composition of coral reefs. *Conservation Biology* 15(4):892–902.
- Eakin, C.M., J.A. Morgan, S.F. Heron, T.B. Smith, G. Liu, L. Alvarez-Filip, B. Baca, E. Bartels, C. Bastidas, C. Bouchon, and others. In press. Caribbean corals in crisis: Record thermal stress, bleaching, and mortality in 2005. *PLoS One*.
- Elvidge, C.D., J.B. Dietz, R. Berkemans, S. Andréfouët, W. Skirving, A.E. Strong, and B.T. Tuttle. 2004. Satellite observation of Keppel Islands (Great Barrier Reef) 2002 coral bleaching using IKONOS data. *Coral Reefs* 23(3):461–462.
- Emery, W.J., S. Castro, G.A. Wick, P. Schluessel, and C. Donlon. 2001. Estimating sea surface temperature from infrared satellite and in situ temperature data. *Bulletin of the American Meteorological Society* 82(12):2,773–2,785.
- Fabricius, K.E. 2005. Effects of terrestrial runoff on the ecology of corals and coral reefs: Review and synthesis. *Marine Pollution Bulletin* 50:125–146.
- Field, M.E., S.A. Cochran, J.B. Logan, and C.D. Storlazzi, eds. 2008. *The Coral Reef of South Moloka'i, Hawai'i: Portrait of a Sediment-Threatened Fringing Reef*. US Geological Survey Scientific Investigations Report 2007-5101, US Geological Survey, 190 pp. Available online at: <http://pubs.usgs.gov/sir/2007/5101> (accessed September 22, 2010).
- Friedlander, A., G. Aeby, R. Brainard, E. Brown, K. Chaston, A. Clark, P. McGowan, C. Morishige, J. Rooney, H. Slay, and others. 2008. The state of coral reef ecosystems of the Main Hawaiian Islands. Pp. 219–253 in *The State of Coral Reef Ecosystems of the United States and Pacific Freely Associated States: 2008*. J.E. Waddell and A.M. Clarke, eds, NOAA Technical Memorandum NOS NCCOS 73, NOAA/NCCOS Center for Coastal Monitoring and Assessment's Biogeography Team, Silver Spring, MD.
- Friedlander, A.M., E.K. Brown, and M.E. Monaco. 2007. Coupling ecology and GIS to evaluate efficacy of marine protected areas in Hawaii. *Ecological Applications* 17(3):715–730.
- García-Sais, J., R. Appeldoorn, T. Battista, L. Bauer, A. Bruckner, C. Caldwell, L. Carrubba, J. Corredor, E. Diaz, C. Liljestrom, and others. 2008. The state of coral reef ecosystems of Puerto Rico. Pp. 75–116 in *The State of Coral Reef Ecosystems of the United States and Pacific Freely Associated States: 2008*. J.E. Waddell and A.M. Clarke, eds, NOAA Technical Memorandum NOS NCCOS 73, NOAA/NCCOS Center for Coastal Monitoring and Assessment's Biogeography Team, Silver Spring, MD.
- Gledhill, D.K., R. Wanninkhof, and C.M. Eakin. 2009. Observing ocean acidification from space. *Oceanography* 22(4):49–59.
- Gledhill, D.K., R. Wanninkhof, F.J. Millero, and M. Eakin. 2008. Ocean acidification of the Greater Caribbean Region 1996–2006. *Journal of Geophysical Research* 113, C10031, doi:10.1029/2007JC004629.
- Gordon, H.R., and M. Wang. 1994. Retrieval of water-leaving radiance and aerosol optical thickness over the oceans with SeaWiFS: A preliminary algorithm. *Applied Optics* 33(3):443–452.
- Goreau, T.J., and R.L. Hayes. 1994. Coral bleaching and ocean "hot spots." *Ambio* 23:176–180.
- Green, A., S.E. Smith, G. Lipsett-Moore, C. Groves, N. Peterson, S. Sheppard, P. Lokani, R. Hamilton, J. Almany, J. Aitsi, and L. Bualia. 2009. Designing a resilient network of marine protected areas for Kimbe Bay, Papua New Guinea. *Oryx* 43(4):488–498.
- Guzmán, H.M., J. Cortés, P.W. Glynn, and R.H. Richmond. 1990. Coral mortality associated with dinoflagellate blooms in the eastern Pacific (Costa Rica and Panama). *Marine Ecology Progress Series* 60:299–303.
- Hamel, M.A., and S. Andréfouët. 2010. Using very high resolution remote sensing for the management of coral reef fisheries: Review and perspectives. *Marine Pollution Bulletin* 60:1,397–1,405.

- Hansen, L.B., N. Kamstrup, and B.U. Hansen. 2002. Estimation of net shortwave radiation by the use of remote sensing and a digital elevation model: A case study of a high arctic mountainous area. *International Journal of Remote Sensing* 23(21):4,699–4,718.
- Hedley, J.D., and P.J. Mumby. 2002. Biological and remote sensing perspectives of pigmentation in coral reef organisms. *Advances in Marine Biology* 43:277–317.
- Hedley, J.D., P.J. Mumby, K.E. Joyce, and S.R. Phinn. 2004. Spectral unmixing of coral reef benthos under ideal conditions. *Coral Reefs* 23:60–73.
- Hoegh-Guldberg, O., P.J. Mumby, A.J. Hooten, R.S. Steneck, P. Greenfield, E. Gomez, C.D. Harvell, P.F. Sale, A.J. Edwards, K. Caldeira, and others. 2007. Coral reefs under rapid climate change and ocean acidification. *Science* 318(5857):1,737–1,742, doi:10.1126/science.1152509.
- Hoegh-Guldberg, O., P.J. Mumby, A.J. Hooten, R.S. Steneck, P. Greenfield, E. Gomez, C.D. Harvell, P.F. Sale, A.J. Edwards, K. Caldeira, and others. 2008. Response to coral adaptation in the face of climate change. *Science* 320 (5874):315–316, doi:10.1126/science.320.5874.315.
- Hogrefe, K.R., D.J. Wright, and E.J. Hochberg. 2008. Derivation and integration of shallow-water bathymetry: Implications for coastal terrain modeling and subsequent analyses. *Marine Geodesy* 31:299–317.
- Holden, H., and E. LeDrew. 1998. Spectral discrimination of healthy and non-healthy corals based on cluster analysis, principal components analysis and derivative spectroscopy. *Remote Sensing of Environment* 65:217–224.
- Hu, C., F. Muller-Karger, B. Murch, D. Myhre, J. Taylor, R. Luerssen, C. Moses, C. Zhang, L. Gramer, and J.C. Hendee. 2009. Building an automated integrated observing system to detect sea surface temperature anomaly events in the Florida Keys. *IEEE Transactions on Geoscience and Remote Sensing* 47(6):1,607–1,620.
- IOCCG (International Ocean Colour Coordinating Group). 2000. *Remote Sensing of Ocean Colour in Coastal and Other Optically-Complex Waters*. S. Sathyendranath, ed., Report of the Ocean Color Coordinating Group No. 3. IOCCG, Dartmouth, Canada, 140 pp. Available online at: <http://www.ioccg.org/reports/report3.pdf> (accessed September 22, 2010).
- IOCCG. 2010. *Atmospheric Correction for Remotely-Sensed Ocean-Colour Products*. M. Wang, ed., Reports of International Ocean-Color Coordinating Group No. 10. IOCCG, Dartmouth, Canada, 78 pp. Available online at: <http://www.ioccg.org/reports/report10.pdf> (accessed September 22, 2010).
- IPCC (Intergovernmental Panel on Climate Change). 2007. *Climate Change 2007: Synthesis Report, Contribution of Working Groups I, II and III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. Core Writing Team, R.K. Pachauri and A. Reisinger, eds, IPCC, Geneva, Switzerland, 104 pp. Available online at: [http://www.ipcc.ch/publications\\_and\\_data/publications\\_ipcc\\_fourth\\_assessment\\_report\\_synthesis\\_report.htm](http://www.ipcc.ch/publications_and_data/publications_ipcc_fourth_assessment_report_synthesis_report.htm) (accessed September 22, 2010).
- Joyce, K.E., S.R. Phinn, C.M. Roelfsema, D.T. Neil, and W.C. Dennison. 2004. Combining Landsat ETM plus and Reef Check classifications for mapping coral reefs: A critical assessment from the southern Great Barrier Reef, Australia. *Coral Reefs* 23(1):21–25.
- Kandirmaz, H.M., L. Yegingil, V. Pestemalci, and N. Emrahoglu. 2004. Daily global solar radiation mapping of Turkey using Meteosat satellite data. *International Journal of Remote Sensing* 25(11):2,159–2,168.
- Kilpatrick, K.A., G.P. Podesta, and R. Evans. 2001. Overview of the NOAA/NASA Advanced Very High Resolution Radiometer Pathfinder algorithm for sea surface temperature and associated matchup database. *Journal of Geophysical Research* 106:9,179–9,197.
- Klemas, V.V. 2009. Remote sensing of coastal resources and environment. *Environmental Research, Engineering and Management* 2(48):11–18.
- Knudby, A., E. LeDrew, and A. Brenning. 2010. Predictive mapping of reef fish species richness, diversity and biomass in Zanzibar using IKONOS imagery and machine-learning techniques. *Remote Sensing of Environment* 114:1,230–1,241.
- Knudby, A., E. LeDrew, and C. Newman. 2007. Progress in the use of remote sensing for coral reef biodiversity studies. *Progress in Physical Geography* 31(4):421–434.
- Knudby, A., C.M. Newman, and E. LeDrew. 2008. Remote sensing for studies of the spatial distribution of coral reef fishes. Pp. 631–635 in *Proceedings of the 11<sup>th</sup> International Coral Reef Symposium*. Ft. Lauderdale, FL.
- Kuffner, I.B., J.C. Brock, R. Grober-Dunsmore, V.E. Bonito, T.D. Hickey, and C.W. Wright. 2007. Relationships between reef fish communities and remotely sensed rugosity measurements in Biscayne National Park, Florida, USA. *Environmental Biology of Fishes* 78(1):71–82.
- Lagerloef, G., F.R. Colomb, D. Le Vine, F.J. Wentz, S. Yueh, C. Ruf, J. Lilly, J. Gunn, Y. Chao, A. Decharon, and others. 2008. The Aquarius/SAC-D Mission: Designed to meet the salinity remote-sensing challenge. *Oceanography* 21(1):68–81. Available online at: [http://www.tos.org/oceanography/issues/issue\\_archive/21\\_1.html](http://www.tos.org/oceanography/issues/issue_archive/21_1.html) (accessed September 27, 2010).
- LeDrew, E., M. Wulder, and H. Holden. 2000. Change detection of satellite imagery for reconnaissance of stressed tropical corals. Paper presented at IEEE Geoscience and Remote Sensing Society, 2000, Honolulu, HI, July 24–28, 2000.
- Lee, Z., K.L. Carder, and R.A. Arnone. 2002. Deriving inherent optical properties from water color: A multiband quasi-analytical algorithm for optically deep waters. *Applied Optics* 41(27):5,755–5,772.
- Lesser, M.P., and C.D. Mobley. 2007. Bathymetry, water optical properties, and benthic classification of coral reefs using hyperspectral remote sensing imagery. *Coral Reefs* 26:819–829.
- Liu, G., A.E. Strong, and W. Skirving. 2003. Remote sensing of sea surface temperatures during 2002 Barrier Reef coral bleaching. *Eos, Transactions, American Geophysical Union* 84(15):137.
- Liu, G., A.E. Strong, W.J. Skirving, and L.F. Arzayus. 2006. Overview of NOAA Coral Reef Watch Program's near-real-time satellite global coral bleaching monitoring activities. Pp. 1,783–1,793 in *Proceedings of the 10<sup>th</sup> International Coral Reef Symposium*. Okinawa, Japan, June 28–July 2, 2006.
- Lyzenga, D.R., N.R. Malinas, and F.J. Tanis. 2006. Multispectral bathymetry using a simple physically based algorithm. *IEEE Transactions on Geoscience and Remote Sensing* 44:2,251–2,259.
- Maina, J., V. Venus, T.R. McClanahan, and M. Ateweberhan. 2008. Modelling susceptibility of coral reefs to environmental stress using remote sensing data and GIS models. *Ecological Modelling* 212:180–199.
- Manzello, D., J.C. Hendee, D. Ward, and Z. Hillis-Starr. 2006. An evaluation of environmental parameters coincident with the partial bleaching event in St. Croix, US Virgin Islands. Pp. 709–717 in *Proceedings of the 10<sup>th</sup> International Coral Reef Symposium*, Okinawa, Japan, June 28–July 2, 2004.
- Manzello, D.P., J.A. Kleypas, D.A. Budd, C.M. Eakin, P.W. Glynn, and C. Langdon. 2008. Poorly cemented coral reefs of the eastern tropical Pacific: Possible insights into reef development in a high-CO<sub>2</sub> world. *Proceedings of the National Academy of Sciences of the United States of America* 105(30):10,450–10,455.
- Martin, S. 2004. *An Introduction to Ocean Remote Sensing*. Cambridge University Press, Cambridge, UK, 454 pp.
- McClanahan, T., M. Ateweberhan, C. Ruiz Sebastián, N. Graham, S. Wilson, J. Bruggemann, and M. Guillaume. 2007. Predictability of coral bleaching from synoptic satellite and *in situ* temperature observations. *Coral Reefs* 26(3):695–701.
- Mellin, C., S. Andréfouët, M. Kulbicki, M. Dalleau, and L. Vigliola. 2009. Remote sensing and fish-habitat relationships in coral reef ecosystems: Review and pathways for systematic multi-scale hierarchical research. *Marine Pollution Bulletin* 58:11–19.
- Mishra, D., S. Narumalani, D. Rundquist, and M. Lawson. 2006. Benthic habitat mapping in tropical marine environments using QuickBird multispectral data. *Photogrammetric Engineering & Remote Sensing* 72(9):1,037–1,048.
- Moberg, F., and C. Folke. 1999. Ecological goods and services of coral reef ecosystems. *Ecological Economics* 29:215–233.

- Moses, C.S., S. Andréfouët, C.J. Kranenburg, and F.E. Muller-Karger. 2009. Regional estimates of reef carbonate dynamics and productivity using Landsat 7 ETM+, and potential impacts from ocean acidification. *Marine Ecology Progress Series* 380:103–115.
- Muller-Karger, F.E., and R.A. Castro. 1994. Mesoscale processes affecting phytoplankton abundance in the southern Caribbean Sea. *Continental Shelf Research* 14(2/3):199–221.
- Muller-Karger, F.E., C. Hu, S. Andréfouët, R. Varela, and R. Thunell. 2005. The color of the coastal ocean and applications in the solution of research and management problems. Pp. 101–127 in *Remote Sensing of Coastal Aquatic Environments: Technologies, Techniques and Applications*. R.L. Miller, C.E. Del Castillo, and B.A. McKee, eds, Springer, Dordrecht.
- Muller-Parker, G., and C.F. D'Elia. 1997. Interactions between corals and their symbiotic algae. Pp. 96–105 in *Life and Death of Coral Reefs*. C. Birkeland, ed., Chapman & Hall, New York.
- Mumby, P.J. 2006. Connectivity of reef fish between mangroves and coral reefs: Algorithms for the design of marine reserves at seascape scales. *Biological Conservation* 128:215–222.
- Mumby, P.J., and A.J. Edwards. 2002. Mapping marine environments with IKONOS imagery: Enhanced spatial resolution can deliver greater thematic accuracy. *Remote Sensing of Environment* 82(2–3):248–257.
- Mumby, P.J., J.R.M. Chisholm, C.D. Clark, J.D. Hedley, and J. Jaubert. 2001. A bird's-eye view of the health of coral reefs. *Nature* 413(6851):1.
- Mumby, P.J., E.P. Green, C.D. Clark, and A.J. Edwards. 1998. Digital analysis of multi-spectral airborne imagery of coral reefs. *Coral Reefs* 17(1):59–69.
- Mumby, P.J., J.D. Hedley, J.R.M. Chisholm, C.D. Clark, H. Ripley, and J. Jaubert. 2004. The cover of living and dead corals from airborne remote sensing. *Coral Reefs* 23(2):171–183.
- Mumby, P.J., W. Skirving, A.E. Strong, J.T. Hardy, E.F. LeDrew, E.J. Hochberg, R.P. Stumpf, and L.T. David. 2004. Remote sensing of coral reefs and their physical environment. *Marine Pollution Bulletin* 48(3–4):219.
- Myint, S.W., C.P. Giri, L. Wang, Z. Zhu, and S.C. Gillette. 2008. Identifying mangrove species and their surrounding land use and land cover classes using an object-oriented approach with a lacunarity spatial measure. *GIScience and Remote Sensing* 45(2):188–208.
- NOAA Coral Reef Conservation Program. 2009. *NOAA Coral Reef Conservation Program Goals & Objectives 2010–2015*. Silver Spring, MD, 40 pp. Available online at: [coralreef.noaa.gov/aboutcrp/strategy/currentgoals/.../3threats\\_go.pdf](http://coralreef.noaa.gov/aboutcrp/strategy/currentgoals/.../3threats_go.pdf) (accessed September 27, 2010).
- Obura, D.O. 2005. Resilience and climate change: Lessons from coral reefs and bleaching in the western Indian Ocean. *Estuarine, Coastal and Shelf Science* 63(3):353.
- Phinn, S., C. Roelfsema, A.G. Dekker, V.E. Brando, and J. Anstee. 2008. Mapping seagrass species, cover and biomass in shallow waters: An assessment of satellite multi-spectral and airborne hyper-spectral imaging systems in Moreton Bay (Australia). *Remote Sensing of Environment* 112(8):3,413–3,425.
- Pittman, S.J., J.D. Christensen, C. Caldow, C. Menza, and M.E. Monaco. 2007. Predictive mapping of fish species richness across shallow-water seascapes in the Caribbean. *Ecological Modelling* 204:9–21.
- Porter, J.W., and J.I. Tougas. 2001. Reef ecosystems: Threats to their biodiversity. Pp. 73–95 in *Encyclopedia of Biodiversity*. S.A. Levin, ed., Academic Press, San Diego, CA.
- Purkis, S.J., S.W. Myint, and B.M. Riegl. 2006. Enhanced detection of the coral *Acropora cervicornis* from satellite imagery using a textural operator. *Remote Sensing of Environment* 101:82–94.
- Qin, Y., V.E. Brando, A.G. Dekker, and D. Blondeau-Patissier. 2007. Validity of SeaDAS water constituents retrieval algorithms in Australian tropical coastal waters. *Geophysical Research Letters* 34:1–4.
- Reynolds, R.W., T.M. Smith, C. Liu, D.B. Chelton, K.S. Casey, and M.G. Schlax. 2007. Daily high-resolution-blended analyses for sea surface temperature. *Journal of Climate* 20:5,473–5,496.
- Robinson, I.S. 2004. *Measuring the Oceans from Space: The Principles and Methods of Satellite Oceanography*. Springer-Verlag, New York, 669 pp.
- Rogers, C.S. 1990. Responses of coral reefs and reef organisms to sedimentation. *Marine Ecology Progress Series* 62:185–202.
- Rowlands, G.P., J.A. Goodman, B.M. Riegl, P.G. Renaud, and S.J. Purkis. 2008. Habitat mapping in the Farasan Islands (Saudi Arabia) using CASI and QuickBird imagery. Pp. 642–646 in *Proceedings of the 11<sup>th</sup> International Coral Reef Symposium*. Session number 17, Ft. Lauderdale, FL, July 7–11, 2008.
- Scopélitis, J., S. Andréfouët, S. Phinn, L. Arroyo, M. Dalleau, A. Cros, and P. Chabanet. 2010. The next step in shallow coral reef monitoring: Combining remote sensing and *in situ* approaches. *Marine Pollution Bulletin*, doi:10.1016/j.marpolbul.2010.07.033.
- Skirving, W., and J. Guinotte. 2001. The sea surface temperature story on the Great Barrier Reef during the coral bleaching event of 1998. Pp. 301–314 in *Oceanographic Processes of Coral Reefs: Physical and Biological Links in the Great Barrier Reef*. E. Wolanski, ed., CRC Press, Boca Raton, FL.
- Soto, I., S. Andréfouët, C. Hu, F.E. Muller-Karger, C.C. Wall, J. Sheng, and B.G. Hatcher. 2009. Physical connectivity in the Mesoamerican Barrier Reef System inferred from 9 years of ocean color observations. *Coral Reefs* 28:415–425.
- Strong, A.E., C.S. Barrientos, C. Duda, and J. Sapper. 1997. Improved satellite techniques for monitoring coral reef bleaching. Pp. 1,495–1,498 in *Proceedings of the 8<sup>th</sup> International Coral Reef Symposium Vol. 2*. H.A. Lessios and I.G. Macintyre, eds, Smithsonian Tropical Research Institute, Panama.
- Strong, A.E., G. Liu, C.M. Eakin, T.R.L. Christensen, W.J. Skirving, D.K. Gledhill, S.F. Heron, and J.A. Morgan. 2009. Implications for our coral reefs in a changing climate over the next few decades: Hints from the past 22 years. Pp. 1,324–1,328 in *Proceedings of the 11<sup>th</sup> International Coral Reef Symposium*. Ft. Lauderdale, FL, July 7–11, 2008.
- Su, H., H. Liu, and W.D. Heyman. 2008. Automated derivation of bathymetric information from multi-spectral satellite imagery using a non-linear inversion model. *Marine Geodesy* 31:281–298.
- Tovar, H.F., and J.M. Baldasano. 2001. Solar radiation mapping from NOAA AVHRR data in Catalonia, Spain. *Journal of Applied Meteorology* 40:1,821–1,834.
- UNEP-WCMC (United Nations Environment Programme-World Conservation Monitoring Centre). 2006. *In the Front Line: Shoreline Protection and Other Ecosystem Services from Mangroves and Coral Reefs*. UNEP-WCMC, Cambridge, UK, 33 pp.
- van Hooijdonk, R., and M. Huber. 2009. Quantifying the quality of coral bleaching predictions. *Coral Reefs* 28:579–587.
- Veron, J.E.N. 2008. Mass extinctions and ocean acidification: Biological constraints on geological dilemmas. *Coral Reefs* 27:459–472.
- Wabnitz, C.C., S. Andréfouët, D. Torres-Pulliza, F.E. Muller-Karger, and P.A. Kramer. 2008. Regional-scale seagrass habitat mapping in the Wider Caribbean region using Landsat sensors: Applications to conservation and ecology. *Remote Sensing of Environment* 112(8):3,455–3,467, doi:10.1016/j.rse.2008.01.020.
- Wang, M. 2007. Remote sensing of the ocean contributions from ultraviolet to near-infrared using the shortwave infrared bands: Simulations. *Applied Optics* 46(9):1,535–1,547.
- Wang, M., S. Son, and L.W. Harding Jr. 2009. Retrieval of diffuse attenuation coefficient in the Chesapeake Bay and turbid ocean regions for satellite ocean color applications. *Journal of Geophysical Research* 114, C10011, doi:10.1029/2009JC005286.
- Wang, M., S. Son, and W. Shi. 2009. Evaluation of MODIS SWIR and NIR-SWIR atmospheric correction algorithms using SeaBASS data. *Remote Sensing of Environment* 113:635–644.
- Wentz, F.J., L. Ricciardulli, K. Hilburn, and C. Mears. 2007. How much more rain will global warming bring? *Science* 317(233):233–235.
- Yamano, H., and M. Tamura. 2004. Detection limits of coral reef bleaching by satellite remote sensing: Simulation and data analysis. *Remote Sensing of Environment* 90:86–103.
- Zhang, H.M., J.J. Bates, and R.W. Reynolds. 2006. Assessment of composite global sampling: Sea surface wind speed. *Geophysical Research Letters* 33:1–5.