

2-2018

The Time has come for Dimensional Personality Disorder Diagnosis

Christopher J. Hopwood
University of California, Davis

Roman Kotov
Stony Brook Univeristy

Robert F. Krueger
University of Minnesota

Thomas A. Widiger
University of Kentucky

Robert R. Althoff
University of Vermont

See next page for additional authors

Follow this and additional works at: https://scholarcommons.usf.edu/psy_facpub



Part of the [Psychology Commons](#)

Scholar Commons Citation

Hopwood, Christopher J.; Kotov, Roman; Krueger, Robert F.; Widiger, Thomas A.; Althoff, Robert R.; Ansell, Emily B.; Bach, Bo; Bagby, R. M.; Blais, Mark A.; and Bornovalova, Marina Alexa, "The Time has come for Dimensional Personality Disorder Diagnosis" (2018). *Psychology Faculty Publications*. 2428.
https://scholarcommons.usf.edu/psy_facpub/2428

This Article is brought to you for free and open access by the Psychology at Scholar Commons. It has been accepted for inclusion in Psychology Faculty Publications by an authorized administrator of Scholar Commons. For more information, please contact scholarcommons@usf.edu.

Authors

Christopher J. Hopwood, Roman Kotov, Robert F. Krueger, Thomas A. Widiger, Robert R. Althoff, Emily B. Ansell, Bo Bach, R. M. Bagby, Mark A. Blais, and Marina Alexa Bornovalova



Published in final edited form as:

Personal Ment Health. 2018 February ; 12(1): 82–86. doi:10.1002/pmh.1408.

The time has come for dimensional personality disorder diagnosis

CHRISTOPHER J. HOPWOOD¹, ROMAN KOTOV², ROBERT F. KRUEGER³, DAVID WATSON⁴, THOMAS A. WIDIGER⁵, ROBERT R. ALTHOFF⁶, EMILY B. ANSELL⁷, BO BACH⁸, R. MICHAEL BAGBY⁹, MARK A. BLAIS¹⁰, MARINA A. BORNOVALOVA¹¹, MICHAEL CHMIELEWSKI¹², DAVID C. CICERO¹³, CHRISTOPHER CONWAY¹⁴, BARBARA DE CLERQ¹⁵, FILIP DE FRUYT¹⁵, ANNA R. DOCHERTY¹⁶, NICHOLAS R. EATON², JOHN F. EDENS¹⁷, MIRIAM K. FORBES³, KELSIE T. FORBUSH¹⁸, MICHAEL P. HENGARTNER¹⁹, MASHA Y. IVANOVA⁶, DANIEL LEISING²⁰, W. JOHN LIVESLEY²¹, MARK R. LUKOWITSKY²², DONALD R. LYNAM²³, KRISTIAN E. MARKON²⁴, JOSHUA D. MILLER²⁵, LESLIE C. MOREY¹⁷, STEPHANIE N. MULLINS-SWEATT²⁶, J. HANS ORMEL²⁷, CHRISTOPHER J. PATRICK²⁸, AARON L. PINCUS²⁹, CAMILO RUGGERO³⁰, DOUGLAS B. SAMUEL²³, MARTIN SELLBOM³¹, TIM SLADE³², JENNIFER L. TACKETT³³, KATHERINE M. THOMAS²³, TIMOTHY J. TRULL³⁴, DAVID D. VACHON³⁵, IRWIN D. WALDMAN³⁶, MONIKA A. WASZCZUK², MARK H. WAUGH³⁷, AIDAN G. C. WRIGHT³⁸, MATHEW M. YALCH³⁹, DAVID H. ZALD⁴⁰, and JOHANNES ZIMMERMANN⁴¹

¹University of California, Davis, Davis California, USA ²Stony Brook University, Stony Brook, New York, USA ³University of Minnesota, Minneapolis, MN, USA ⁴University of Notre Dame, South Bend, IN, USA ⁵University of Kentucky, Lexington, KY, USA ⁶University of Vermont, Burlington, VT, USA ⁷Syracuse University, Syracuse, NY, USA ⁸Region Zealand Psychiatry, Roskilde, Denmark ⁹University of Toronto, Scarborough, Toronto, ON, Canada ¹⁰Harvard Medical School, Boston, MA, USA ¹¹University of South Florida, Tampa, Florida, USA ¹²Southern Methodist University, Dallas, TX, USA ¹³University of Hawai'i, Honolulu, HI, USA ¹⁴College of William & Mary, Williamsburg, VA, USA ¹⁵University of Ghent, Ghent, Belgium ¹⁶University of Utah, Salt Lake City, UT, USA ¹⁷Texas A&M University, College Station, TX, USA ¹⁸University of Kansas, Lawrence, KS, USA ¹⁹Zurich University of Applied Sciences, Zurich, Switzerland ²⁰Technische Universität Dresden, Dresden, Germany ²¹University of British Columbia, Vancouver, BC, Canada ²²Albany Medical College, Albany, NY, USA ²³Purdue University, West Lafayette, IN, USA ²⁴University of Iowa, Iowa City, IA, USA ²⁵University of Georgia, Athens, GA, USA ²⁶Oklahoma State University, Stillwater, OK, USA ²⁷University of Groningen, Groningen, the Netherlands ²⁸Florida State University, Tallahassee, FL, USA ²⁹Pennsylvania State University, State College, PA, USA ³⁰University of North Texas, Dallas, TX, USA ³¹University of Otago, Otago, New Zealand ³²University of New South Wales, Kensington, New South Wales, Australia ³³Northwestern University, Evanston, IL, USA ³⁴University of Missouri, Columbia, MO, USA ³⁵McGill University, Montreal, Quebec, CA ³⁶Emory University, Atlanta, GA, USA ³⁷University of Tennessee, Knoxville, TN, USA ³⁸University of Pittsburgh, Pittsburgh, PA, USA ³⁹Marian University, Indianapolis, IN,

USA ⁴⁰Vanderbilt University, Nashville, TN, USA ⁴¹Psychologische Hochschule Berlin, Berlin, Germany

The committee revising the ICD-11 Mental or Behavioural Disorders section ‘Personality Disorders and Related Traits’ has proposed replacing categorical personality disorders with a severity gradient ranging from personality difficulties to severe personality disorder and five trait domains: negative affectivity, dissocial, disinhibition, anankastic and detachment ¹. While acknowledging that there are multiple potential pathways for moving toward a more evidence-based and clinically useful scheme for classifying personality dysfunction, we applaud and support the proposed transition from a categorical model of personality disorder types, which has proven to be empirically problematic and of limited clinical utility, to a dimensional model of personality disorder that has considerable connection to scientific evidence and potential for clinical application.

There is no evidence supporting the hypotheses that personality disorders are categorical ^{2,3} or that there are 10 (or any other number of) discrete types of personality disorder ⁴. Well-established problems with categorical personality disorder diagnosis such as low reliability, diagnostic comorbidity and within-disorder heterogeneity complicate research and treatment ⁵. There are no validated interventions for most of the categorical personality disorders, and although several psychotherapies from different theoretical perspectives have been developed for borderline personality disorder that have evidence of moderate efficacy, none have proven to be relatively more effective than any of the others ⁶. Evidence for treatment mechanisms is sparse, and there is no evidence that existing approaches have specific efficacy for borderline personality disorder as opposed to general efficacy for a variety of psychiatric difficulties ⁷.

In contrast, there is a vast body of empirical literature supporting dimensional models of personality disorder that are closely aligned with the proposed model ^{8–11}, in addition to the emerging body of work on the specific dimensions proposed for ICD-11 ^{12–19}. The ICD-11 proposal has two elements. The severity dimension has ties to the psychodynamic tradition ^{20,21}, which has historically been at the forefront of personality disorder classification, and aligns with a number of empirical efforts to quantify general personality dysfunction (e.g. ^{22–27}). Research demonstrates that much of the predictive and prognostic value in personality disorder data can be derived from such a dimension ²⁸.

The personality trait model proposed for ICD-11 resembles other dimensional models of personality such as the Five-Factor Model or the DSM-5 Alternative Model for Personality Disorders ^{16,29,30}. Although there are some important differences between the ICD-11 proposal and these other models that will be adjudicated by future research, the more important point at this stage is that evidence consistently supports the validity of dimensional trait models for describing individual differences in personality. In contrast to the categorical model of personality disorder types, there is a large literature on the genetic underpinnings, cross-cultural validity, course, correlates and measurement of broad personality traits ^{11,31}. Dimensional models also address issues such as comorbidity and

heterogeneity in a direct and empirically tractable manner⁸; recapture but empirically reorganize the information provided by personality disorder types³²; and have considerable potential for guiding and tracking treatment^{33,34}. We would highlight that research has repeatedly shown that the borderline personality disorder construct in particular can be accounted for by empirically derived dimensions of personality traits and functioning^{35–40}.

Nevertheless, some people in the field continue to argue in favour of personality diagnosis by categorical types. We are concerned about the implications of retaining a categorical system that has been so thoroughly shown to be empirically and clinically problematic. It is very difficult to justify allocating resources toward continued research on an approach that has proven to be fundamentally flawed, as opposed to a dimensional model that points to exciting new avenues for research on aetiology, mechanisms and treatment (e.g.⁴¹). We are likewise concerned about the implications that retaining a demonstrably problematic model has for patients' lives. It would be very unsettling to be told that one's problems are due to a specific medical condition, only to learn later that the supposed condition had been abandoned by the medical community. It is probably already confusing for patients, who might discover via an internet search on their personality diagnosis that much of the field does not believe such a disorder actually exists. It would be far preferable to be straightforward with our patients about what we know and do not know regarding personality and its related problems than to label them with legacy diagnoses that will not stand the test of time.

Reasonable concerns have been expressed about challenges associated with the transition from a categorical to a dimensional model of personality disorder. Such concerns need to be balanced against several field surveys that show that a majority of clinicians and researchers support the transition to a more dimensional, evidence-based framework^{42–44}. We acknowledge that the transition to a dimensional model needs to be thoughtful with regard to issues such as third-party reimbursement. Moreover, we recognize that legal, community mental health and other systems will need to be educated regarding how to translate from the old system to the new. However, we do not believe that these practical issues provide a compelling rationale for retaining a system that does not effectively capture individual differences in patients' personality difficulties. In contrast, moving forward with an evidence-based framework for diagnosing personality disorders has significant potential to stimulate research that can lead to new treatments and aetiological models that will ultimately reduce the burden of personality disorders on patients, families and society. The changes proposed for ICD-11 also provide a generative model for conceptualizing the meta-structure of psychopathology. Indeed, there are clear phenotypic and genetic links between the dimensions proposed for ICD-11 and a number of mental health conditions beyond personality disorders^{45,46}.

Past scientists believed that the sun revolved around the earth, the brain was organized according to the principles of phrenology, and spirits were responsible for psychiatric problems. It is a testament to science that these views gave way to a more accurate model of nature. The new perspectives that replaced them contributed to major advancements in astronomy, neuroscience and mental health. Likewise, the evidence is clear that personality disorders do not exist as 10 discrete types. The categorical model has become a hindrance to

research and practice. As an example, see the unfortunate outcome of the *DSM-5* revision process, in which a model that is not supported by evidence or the majority of the field was retained as the official diagnostic scheme despite the viable alternative proposed by the Personality and Personality Disorders Work Group, published in Section III of *DSM-5*. It is time for the field to transition to a model that fits research data and clinical reality. The ICD-11 proposal connects psychiatric classification of personality disorder manifestations with scientific evidence. The proposed changes would enhance diagnostic efficiency and patient care while spurring research that can further improve the assessment and treatment of psychopathology. As clinicians and researchers who have dedicated our careers to understanding and helping people with personality pathology, we urge the ICD-11 PD work group to remain committed to an evidence-based revision of personality disorder diagnosis.

References

1. Tyrer P, Crawford M, Mulder R. Reclassifying personality disorders. *Lancet*. 2011; 377:1814–5. [PubMed: 21353696]
2. Clark LA. Assessment and diagnosis of personality disorder: perennial issues and emerging conceptualization. *Annu Rev Psychol*. 2007; 58:227–58. [PubMed: 16903806]
3. Trull TJ, Durrett CA. Categorical and dimensional models of personality disorder. *Annu Rev Clin Psychol*. 2005; 1:355–80. [PubMed: 17716092]
4. Anderson J, Snider S, Sellbom S, Krueger RF, Hopwood CJ. A comparison of the DSM-5 section II and section III personality disorder structures. *Psychiatry Res*. 2014; 216:363–72. [PubMed: 24656519]
5. Widiger TA, Samuel DB. Diagnostic categories or dimensions? A question for the diagnostic and statistical manual of mental disorders, fifth edition. *J Abnorm Psychol*. 2005; 114:494–504. [PubMed: 16351373]
6. Cristia IA, Gentili C, Cotet CD, Palomba D, Barbui C, Cuijpers P. Efficacy of psychotherapies for borderline personality disorder: a systematic review and meta-analysis. *JAMA Psychiat*. 2017; 74:319–28.
7. Bateman AW, Gunderson J, Mulder R. Treatment of personality disorder. *Lancet*. 2015; 385:735–43. [PubMed: 25706219]
8. Krueger RF, Markon KE. The role of the DSM-5 personality trait model in moving toward a quantitative and empirically based approach to classifying personality and psychopathology. *Annu Rev Clin Psychol*. 2014; 10:477–501. [PubMed: 24329179]
9. Markon KE, Krueger RF. Delineating the structure of normal and abnormal personality: an integrative hierarchical approach. *J Pers Soc Psychol*. 2005; 88:139–57. [PubMed: 15631580]
10. Morey LC, Hopwood CJ, Gunderson JG, Skodol AE, Shea MT, Yen S, et al. Comparison of alternative models for personality disorders. *Psychol Med*. 2007; 37:983–94. [PubMed: 17121690]
11. Widiger TA, Trull TJ. Plate tectonics in the classification of personality disorder. *American Psychologist*. 2007; 62:71–83. [PubMed: 17324033]
12. Kim Y-R, Blashfield R, Tyrer P, Hwang S, Lee H. Field trial of a putative research algorithm for the diagnosis of ICD-11 personality disorders in psychiatric patients: 1. severity of personality disturbance. *Personality and Mental Health*. 2014; 8:67–78. [PubMed: 24532556]
13. Kim Y-R, Tyrer P, Kim H-S, Kim S-G, Hwang ST, Lee G-Y, et al. Preliminary field trial of a computer algorithm for diagnosing ICD-11 personality disorders in psychiatric patients: 2. proposed trait domains. *Personality and Mental Health*. 2015; 9:298–307. [PubMed: 26472077]
14. Kim YR, Tyrer P, Lee HS, Kim SG, Connan F, Kinnaird E, et al. Schedule for personality assessment from notes and documents (SPAN-DOC): preliminary validation, links to the ICD-11 classification of personality disorder, and use in eating disorders. *Personality and Mental Health*. 2016; 10:106–17. [PubMed: 27120421]

15. Mulder RT, Horwood J, Tyrer P, Carter J, Joyce PR. Validating the proposed ICD-11 domains. *Personality and Mental Health*. 2016; 10:84–95. [PubMed: 27120419]
16. Oltmanns JR, Widiger TA. A self-report measure of the ICD-11 dimensional trait model proposal: the personality inventory for ICD-11. *Psychol Assess*. in press.
17. Tyrer P, Crawford M, Sanatnia R, Tyrer H, Cooper S, Muller-Pollard C, et al. Preliminary studies of the ICD-11 classification of personality disorder in practice. *Personality and Mental Health*. 2014; 8:254–63. [PubMed: 25200623]
18. Tyrer P, Tyrer H, Yang M, Guo B. Long-term impact of temporary and persistent personality disorder on anxiety and depressive disorders. *Personality and Mental Health*. 2016a; 10:76–83. [PubMed: 26754031]
19. Tyrer P, Wang D, Tyrer H, Crawford M, Cooper S. Dimensions of dependence and their influence on the outcome of cognitive behavior therapy for health anxiety: randomized controlled trial. *Personality and Mental Health*. 2016b; 10:96–105. [PubMed: 27120420]
20. Bender DS, Morey LC, Skodol AE. Toward a model for assessing level of personality functioning in DSM-5, part I: a review of theory and methods. *J Pers Assess*. 2011; 93:332–46. [PubMed: 22804672]
21. Kernberg, OF. *Severe Personality Disorders: Psychotherapeutic Strategies*. New Haven, CT: Yale University Press; 1984.
22. Bornstein RF. Reconceptualizing personality disorder diagnosis in the DSM-V: the discriminant validity challenge. *Clinical Psychology: Science and Practice*. 1998; 5:333–43.
23. Hopwood CJ, Malone JC, Ansell EB, Sanislow CA, Grilo CM, McGlashan TH, et al. Personality assessment in DSM-V: empirical support for rating severity, style, and traits. *J Pers Disord*. 2011; 25:305–20. [PubMed: 21699393]
24. Livesley WJ. Suggestions for a framework for an empirically based classification of personality disorder. *Can J Psychiatry*. 1998; 43:137–47. [PubMed: 9533966]
25. Morey LC. Development and initial evaluation of a self-report form of the DSM-5 level of personality functioning scale. *Psychol Assess*. in press.
26. Morey LC, Berghuis H, Bender DS, Verheul R, Krueger RF, Skodol AE. Toward a model for assessing level of personality functioning in DSM-5, part II: empirical articulation of a core dimension of personality pathology. *J Pers Assess*. 2011; 93:346–53.
27. Parker G. The etiology of personality disorders: a review and consideration of research models. *J Pers Disord*. 1997; 11:345–69. [PubMed: 9484696]
28. Wright AGC, Hopwood CJ, Skodol AE, Morey LC. Longitudinal validation of general and specific structural features of personality pathology. *J Abnorm Psychol*. 2016; 125:1120–34. [PubMed: 27819472]
29. Bach B, Sellbom M, Kongerslev M, Simonsen E, Krueger RF, Mulder R. Deriving ICD-11 personality disorder domains from DSM-5 traits: initial attempt to harmonize two diagnostic systems. *Acta Psychiatr Scand*. in press.
30. Widiger TA, Simonsen E. Alternative dimensional models of personality disorder: finding a common ground. *J Pers Disord*. 2005; 19:110–30. [PubMed: 15899712]
31. Soto CJ, Tackett JL. Personality traits in childhood and adolescence: structure, development, and outcomes. *Current Directions in Psychological Science*. 2015; 24:358–62.
32. Miller JD, Few LR, Lynam DR, MacKillop J. Pathological personality traits can capture DSM-IV personality disorder types. *Personality Disorders: Theory, Research, and Treatment*. 2015; 6:32–40.
33. Hopwood CJ, Zimmermann J, Pincus AL, Krueger RF. Connecting personality structure and dynamics: toward a more evidence-based and clinically useful diagnostic scheme. *J Pers Disord*. 2015; 29:431–48. [PubMed: 26200845]
34. Roberts BW, Luo J, Briley DA, Chow P, Su R, Hill PL. A systematic review of personality trait change through intervention. *Psychol Bull*. 2017; 143:117–41. [PubMed: 28054797]
35. Bornoalova MA, Hicks BM, Patrick CJ, Iacono WG, McGue M. Development and validation of the Minnesota borderline personality disorder scale (MBPD). *Assessment*. 2011; 18:234–52. [PubMed: 21467094]

36. Miller JD, Morse JQ, Nolf K, Stepp SD, Pilkonis PA. Can DSM-IV borderline personality disorder be diagnosed via dimensional personality traits? Implications for the DSM-5 personality disorder proposal. *J Abnorm Psychol.* 2012; 121:944–50. [PubMed: 22428791]
37. Sellbom M, Sansone RA, Songer DA, Anderson JL. Convergence between DSM-5 section II and section III diagnostic criteria for borderline personality disorder. *Australian and New Zealand Journal of Psychiatry.* 2014; 48:325–32. [PubMed: 24253360]
38. Sharp C, Wright AGC, Fowler JC, Freuh C, Allen JG, Oldham J, et al. The structure of personality pathology: both general ('g') and specific ('s') factors? *J Abnorm Psychol.* 2015; 124:387–98. <http://www.personalityprocesses.com/s/Sharp-Wright-et-al-in-press-Strucutre-of-PP.pdf>. [PubMed: 25730515]
39. Trull TJ, Widiger TA, Lynam DR, Costa PT Jr. Borderline personality disorder from the perspective of general personality functioning. *J Abnorm Psychol.* 2003; 112:193–202. [PubMed: 12784828]
40. Wright AGC, Hopwood CJ, Zanarini MC. Associations between changes in normal personality traits and borderline personality disorder symptoms over 16 years. *Personality Disorders: Theory, Research, and Treatment.* 2015; 6:1–11.
41. Cuthbert BN. The RDoC framework: facilitating transition from ICD/DSM to dimensional approaches that integrate neuroscience and psychopathology. *World Psychiatry.* 2014; 13:28–35. [PubMed: 24497240]
42. Bernstein DP, Iscan C, Maser J. Boards of Directors of the Association for Research in Personality Disorders and Interntional Society for the Study of Personality Disorders. Opinions of personality disorder experts regarding the DSM-IV personality disorder classification system. *J Pers Disord.* 2007; 21:536–51. [PubMed: 17953505]
43. Keeley J, Reed G, Roberts M, Evans S, Medina-Mora M, Robles R, et al. Developing a science of clinical utility in diagnostic classification systems field study strategies for ICD-11 mental and behavioral disorders. *American Psychologist.* 2016; 71:3–16. [PubMed: 26766762]
44. Morey LC, Skodol AE, Oldham JM. Clinician judgments of clinical utility: a comparison of DSM-IV-TR personality disorders and the alternative model for DSM-5 personality disorders. *J Abnorm Psychol.* 2014; 123:398–405. [PubMed: 24886013]
45. Kotov R, Gamez W, Schmidt F, Watson D. Linking big personality traits to anxiety, depressive, and substance use disorders: a meta-analysis. *Psychol Bull.* 2010; 136:768–821. [PubMed: 20804236]
46. Kotov R, Krueger RF, Watson D, Achenbach TM, Althoff RR, Bagby M, et al. The hierarchical taxonomy of psychopathology (HiTOP): a dimensional alternative to traditional nosologies. *J Abnorm Psychol.* in press.